

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search


Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/





RECORDS OF BIG GAME

THIRD EDITION

Rowland Ward







· ,

	•	
		·









Yours faithfully Romandward

\$ 38.00 C

.

ુડ



Yours faithfully Romandward

RECORDS OF BIG GAME

WITH

THEIR DISTRIBUTION, CHARACTERISTICS, DIMENSIONS, WEIGHTS, AND

MEASUREMENTS OF HORNS ANTLERS, TUSKS, & SKINS

THIRD EDITION

By ROWLAND WARD, F.Z.S.

AUTHOR OF 'THE SPORTSMAN'S HANDBOOK,' ETC.



WITH ABOUT 217 ILLUSTRATIONS

LONDON
ROWLAND WARD, LIMITED
"THE JUNGLE," 166 PICCADILLY, W.
1899

ر د 798.

HARVARD UNIVERSITY LIBRARY

THIS BOOK IS DEDICATED

TO THE


SPORTSMEN OF THE WORLD

WITHOUT WHOSE DETERMINED PLUCK AND LOVE OF TRAVEL THESE RECORDS

COULD NOT HAVE BEEN PRODUCED, NOR COULD THE AUTHOR

. HAVE SPENT A LIFETIME IN THE ART HE HAS

TRIED TO ADVANCE



PREFACE TO THIRD EDITION

In the present edition, the finest known specimens of antlers, horns, tusks, and skins are, so far as possible, recorded. It has been found impracticable in some instances to verify the measurements of trophies in distant parts of the world; and such records must accordingly be taken on the responsibility of their respective owners or other persons who have been good enough to measure them. It is much to be regretted that one pair of hands and a steel-tape are not responsible for the measurements of all the actual "records." the many difficulties in connection with this compilation is due to the circumstance that different measurements of the same specimen are often sent me; this sometimes arising from the use of common tapes or string, which are absolutely unreliable unless checked at the time by a steel measure. In this connection I may mention that I shall at all times be pleased to cause such measurements to be carefully verified at my establishment in Piccadilly. With the horns of freshly killed hollow-horned ruminants an allowance for shrinkage should be made when comparing with older trophies. average specimen of an Ovis ammon horn, for instance, will frequently shrink half an inch in length and proportionately in girth after it has left the field.

With the *Cervidæ* many difficulties have arisen as to comparison; and I may point out that although length of antlers is invariably put at the top of the list, other particulars such as *number of tines*, *general symmetry*, *spread*, and *weight of antler*, are in many instances the making of a good trophy. Small specimens are frequently noted in order to include measurements from different localities, as well as to record horns of certain species, such as *Ovis poli*, etc., in which recently shot specimens do not approach those of the record example.

I have to thank many sportsmen and naturalists all over the world for the help they have afforded me.

To Mr. A. O. Hume, and Prince Henry of Liechtenstein, my special thanks are due.

In the present volume great care has been taken with regard to the accuracy of the numbers; and, considering that there are so many thousand measurements, it will be readily understood the task attempted has been one of no ordinary difficulty.

The new illustrations, numbering over fifty in the present volume, are nearly all produced direct from photographs of notable examples.

A new feature in this edition is the introduction of a short description of the leading characteristics and the exact geographical distribution of each species and race. This, it is hoped, will render the volume of additional value as a work of reference to the sportsman.

ROWLAND WARD.

PREFACE TO SECOND EDITION

SINCE the publication of the first edition of this work under the title of "Horn Measurements," I have been successful in collecting much new and valuable data bearing upon the measurements of horns and other statistics of Big Game. For this information I am to a considerable extent indebted to numerous sportsmen, and to the curators of some of the most important museums of the world, who have been kind enough to measure all the finest specimens in the collections under their charge. In addition to this valuable help, for which I desire to express my grateful acknowledgment, I have had a record carefully prepared of all the most remarkable specimens registered, as well as those that have from time to time passed through our hands. I am indebted to Mr. H. A. Bryden for much of the descriptive matter of the South African game. There will also be found embodied with all these new records numerous illustrations of typical heads, skulls, and horns drawn especially for this new edition.

In addition to this supplementary information, the new edition will be found to contain more exact localities than have hitherto been recorded, and we continue in constant correspondence with sportsmen in almost every part of the world with a view of still further completing our records.

	•			
·		•	•	
			•	

PREFACE TO FIRST EDITION

My object in producing this book is to start a record of Horn Measurements of the Great Game of the World. I only regret that it was not commenced at an earlier date, as in that case it would have been more complete. In my earlier life I had but little help, and often worked thirty hours at a stretch; my work necessitated attention to specimens that demanded immediate treatment, and my love for reproducing life-like studies prevented me from keeping records.

The measurements presented here have been taken principally by one hand, and, for that reason, I value them the more. The dimensions from acknowledged authorities I naturally am not responsible for. This work is not designed to be in any way a scientific treatise, but is prepared for sportsmen and scientific men who are interested to see comparable measurements at a glance. I think these have not been produced in like form before. These records can be added to, and I shall feel indebted to sportsmen who will contribute any authentic record measurements. It must be borne in mind that many trophies which have passed through my hands are now scattered all over the world, and the dimensions of them, for the time, are

not recorded here. Some that I give now are fine measurements, some are only ordinary statistics, many are of new species altogether; these last being mainly a result of the quite recent opening up of Africa. The advice noted as to the way to measure must be dealt with strictly, for many persons measuring by different methods produce untrustworthy record. We have tried to be fair in producing the statistics, taking them rather under than over the mark; and the tape has not been pressed into corrugations of horns, but carried outside, over all inequalities.

R. W.

ABBREVIATIONS AND SIGNS

- Owner's measurements or other known authority. d Male. ... Unrecorded. ... Unrecorded.

R, Right horn or antler. L, Left horn or antler.

Measurements are on the outside of the longest horn from base to tip; with Deer from the bottom edge of burr, or coronet, to the highest tip point, except where notified to the contrary.

Circumference is at the base; with Deer above brow tine; in the Red Deer and Wapiti group between bez and trez.

Length is expressed in inches, when not otherwise noted.

Weights taken in the field should be accepted as approximate, and, unless mentioned, are of adult males.

Heights are in most cases taken at the shoulder of adult males (see p. 480).

P.Z.S. = Proceedings of the Zoological Society of London.

	ě			
				.
			•	

LIST OF ILLUSTRATIONS

PAGE	PAGE
Abyssinian Buffalo Skull and Horns . 395	Cape Hartebeest Horns 117
Addax Skull and Horns 278 African Elephant Head 446	Caspian Red Deer Antlers . 14, 32
African Elephant Head 446	Red Deer Head
,, ,, Tusks 448, 449	Chamois Horns 314
,, Rhinoceros Head 434	Chamois Horns
" ,, Skull and Horns . 433	Chiru Head
American Bighorn Head 365	
,, Bison Skull and Horns 405	Clarke's Gazelle Head 254
Anoa Head 402	
Anoa Head	Coke's Hartebeest Head 115
" Tahr Skull and Horns 330	Congo Buffalo Skull and Horns 397
Arna Head	
Arui Head	;
	Dorcas Gazelle Head 224
Banting Head 416	Duiker Horns 146
,, and Gaur Skulls and Horns . 417	East Caucasian Tur Head 358
Barasingh Head 36	East Caucasian Tur Head
Bear 473	
Beira Antelope Skulls and Horns 258	,, ,, (Cow)
Beisa Head	,, Skull and Antlers
Bharal Head 360	European Bison Skulls and Horns . 403
Blackbuck Head 205	,, Muflon Head 386
,, Horns 202	
Blue Duiker Head 149	Extinct Giant Fallow Deer Skull 59
Roar Head	Fallow Deer Head 56
,, Tusk 422	
Bongo Horns	
,, Tusk	
Brindled Gnu Head 139	Gaur Head 410
,, ,, Horns 136	
Bubaline Hartebeest Horns 106	
Buffon's Kob Horns 183	Gemsbuck Horns
Burchell's Rhinoceros Horns 438	
,, ,, Skull and Horns . 441	Goa Head 217
Burmese Banting Head 416	Grant's Gazelle Head 259
Bushbuck Head 282, 286	Skull and Horns 243
	Gray Rhebok 190
Cape Buffalo Horns 391	Gray's Waterbuck 181

PAGE	PAG
Greater Kudu Head 295	Neumann's Hartebeest Head
,, ,, Horns 299	,, ,, Skull and Horns 11
Grysbuck Head 167	Nilgai Skull and Horns 28 Nilgiri Tahr Head
Hangul Head	Nilgiri Tahr Head
Himalayan Goral 317	Nyala Horns
Ibex Head	Nyasaland Gnu Head 13
Serow Head	
Skull and Horns 220	Oribi Head 160
Tahr Head 228	
Hog-Deer Head	Pekin Sika Head 5.
Hog-Deer Head	Pelzeln's Gazelle Head, male 23.
Hybrid Hartebeest Skull and Horns 120	,, Skull and Horns, female 23.
Trybrid Hartebeest Skull and Horns . 120	Père David's Deer Antlers 80
Impala Head 209	l =
Impala Head 209 ,, Skull and Horns 206	Peters's Gazelle Skulls and Horns 240
Indian Elephant Skull and Tusks . 442	Phillips's Dik-dik Skull 150 Prong-horn Head
,, Gazelle Head 228	Prong-horn Head 104
,, Rhinoceros Heads 430	Przewalski's Gazelle Skull and Horns . 219
•	Puku Head
Jackson's Hartebeest Head 119	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
,, ,, Skull and Horns 120	Red Deer Antlers (Caucasus) 32
Japanese Sika Antlers51Javan Rusa Skull and Antlers68	(Exmoor)
Javan Rusa Skull and Antlers 68	in the Costle at Marite
Johnston's Gnu Head 137	Caucasis
	Old Familia
Kamschatkan Bighorn Skull and Horns 369	,, ,, (Old English) 23
Kirk's Dik-dik	,, ,, (Park) 22
Kirk's Dik-dik	,, ,, (Scotch)
,, Skull and Horns 170	Reedbuck Head
Korrigum Hartebeest Head 126	Reindeer Antlers
	", Head
Lesser Kudu Head 303	Roan Antelope Head 269
Loder's Gazelle Head 236	Rocky Mountain Goat Head 327
Lichi or Lechwe Head 184	Roe Deer Head (Siberian) 86
., Horns 186	,, Skull and Antlers 88
Lesser Kudu Head	
Livingstone's Antelope Skull and Horns 165	Sable Antelope Head
	,, Skull and Horns 260
Malayan Sambar Antlers 65	Saiga Head
Marco Polo's Sheen Head	Horns
Markhor Horns (Astor) 224	Salt's Dik-dik Head
Markhor Horns (Astor)	Sambar Antlers
(Pir Panial) 222 226 227	Sacraby Head
(Sulaman) 239, 330, 337	Sambar Antlers 61 Sassaby Head
,, ,, (Suicinail) 330	Concernition Duffulo Clariff and II and
y, nead y, 340	Senegambian Buffalo Skull and Horns . 396
,, (Suleman)	,, Eland Horns 312
Mesopotamian Fallow Deer Head . 58	Shapu Head
Mongolian Gazelle Skulls and Horns . 220	Shou Skull and Antlers 40, 55
Mountain Reedbuck Head 198	Siberian Argali Skull and Horns 375
Mule-Deer Head 93, 95	,, Skulls and Horns . 370
Muntjac Antlers	Sinaitic Ibex Head 348 Sind Ibex Skull and Horns
,, Head 83	Sind Ibex Skull and Horns 351
Musk Ox Head	Sing-Sing Waterbuck Skull and Horns . 178

		PAGE		PAG
Situtunga Horns		292	Uganda Kob Skull and Horns	18
Sæmmerring's Gazelle Head, &		247	Urial Head	38
Spanish Ibex Head		354		
Speke's Gazelle Head, & .		233	Virginian Deer Antlers	9
,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,,		231	· · · · · · · · · · · · · · · · · · ·	7
Springbuck Head		223	W 11 1 0 11 17 1	
"Horns		214	Waller's Gazelle Head	25
Steinbuck Head		168	Wapiti Head	4
Swamp-Deer Antlers		75	Wart-Hog Head	42
Swayne's Dik-dik Head .		156	" Tusks	42
,, Hartebeest Head .		112	Waterbuck Head	17
			,, Horns	17
Takin Frontlet and Horns, ?		325	West African Harnessed Antelope Horns	-
,, Skull and Horns, & .		324	,, Hartebeest Horns	10
Thameng Antlers		78	West Caucasian Tur Skull and Horns .	35
Thomson's Gazelle Head, &		241	White-bearded Gnu Head	14
,, ,, Horns, ♀		240	,, ,, Skull and Horns .	14
Tibetan Argali Head		377	White-tailed Gnu Head	14
Tiger		480	,, ,, Horns	14
, Skin		459		
Topi Hartebeest Head		127	Yak Head	40
Tora ,, ,, .		110	,, Skull and Horns .	
., ,, Horns .		109	Yarkand Stag Skull and Antlers	39

•

.



ADDITIONS AND CORRECTIONS


OCTOBER 1899

Page.	-									Owner.
5.	Rei	NDEER	OR CA	RIBOU, a	add					
	62	51	40	50]	21 + 17	Cana	da.	•	•	Sir Edmund G. Loder, Bart.
76.						ch specin				Major C. S. Cumberland.
103.		THERN			-	nen (16 fe				F. C. Selous.
I 20.	•		HARTEI	BEESTE.	—16⅓ in	ch specim	en sho	uld b	e a	Mr. Justice Hopley.
221.						specimen 				A. J. Grant.
246.	PET	ers' G	ZELLE.	—Add t	o Heigh	t at Shoul	der, 26	5 to 3	3 in	ches.
275.	_	NGE-EA			irst speci	men shou	ld read	i—	}	R. B. P. Cator.
	33½		7	13		•	••		,	
297.	Gre	ATER F	ζυ ου. —	-Add be	fore first	specimen	—		ì	F. C. Selous.
	64	41	1	I	23 N	Iashona la	nd.			1. C. Scious.

lateral hoofs are unusually large and spreading; and there is a patch



RECORDS OF BIG GAME



Antlers of Male Woodland Reindeer. From a specimen in the British Museum.

REINDEER or CARIBOU (Rangifer tarandus).

DISTINGUISHED from all other deer by the presence of antlers in both sexes; those of males being complex, with the brow tines palmated and often unsymmetrical, and the bez, or second tine, also generally expanded. The muzzle is entirely covered with hair, the ears and tail are short, the throat has a fringe of long hair, and the coat is very thick and of a nearly uniform clove-brown colour, with some white in the region of the tail, and on the under parts and legs. The false or lateral hoofs are unusually large and spreading; and there is a patch

of long white hair covering a gland on the hock, but none on the hind cannon-bone. Height at shoulder reaching to 4 feet 10 inches (Newfoundland); weight of a full-grown Scandinavian stag 30 stone, clean (Abel Chapman); antlers average about 30 lbs. per pair.

Reindeer inhabit the circumpolar regions of both hemispheres, in Europe including Scandinavia, Lapland, and Northern Russia; their



Antlers of Male Woodland Reindeer from Nova Scotia. From a specimen in the British Museum.

southern limit varies from 52° to 54° N. latitude, while they extend to between 80° and 81° northwards.

Six varieties, or local races, of reindeer are recognised: (1) the Scandinavian reindeer (R. tarandus typicus) of Northern Europe and Asia, which is a rather small animal with moderately expanded antlers; (2) the Spitzbergen reindeer (R. tarandus spetzbergensis), characterised by the peculiar form of the nasal bones of the skull; (3) the woodland

reindeer (R. tarandus caribou) of the forest districts of Arctic America, distinguished by its large bodily size and the short, much-palmated antlers; (4) the Newfoundland reindeer (R. tarandus terræ-novæ),



Side view of Antlers of Male Barren-Ground Reindeer. From a specimen in the British Museum.

which is closely allied to the last, but with even more complex antlers and some differences in coloration; (5) the Greenland reindeer (R. tarandus grænlandicus), which is apparently very similar to the sixth variety; and (6) the barren-ground reindeer (R. tarandus arcticus), from the open country north of the forests in America, a very distinct

animal, characterised by its small bodily size, and the great length and simple form of the antlers, in which, except on the brow tine, there is scarcely any palmation. Reindeer heads are probably some of the most difficult to measure, owing to the different curves the top points frequently present. Females have smaller and lighter antlers than males.



Head of Male Newfoundland Reindeer. From Dr. J. A. Allen.

Abel Chapman, writing in the *Field*, says two or three year old bucks run between 16 and 20 stone; but one big stag he shot was made out (perhaps erroneously) to scale well over 30 stone, clean. The measurements of the antlers have been separated for the convenience of sportsmen.

REINDEER or CARIBOU (Rangifer tarandus)—continued.


Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.		Owner.
62	5½	39	49 1	20+17	Canada		Mrs. Macintosh.
578	5‡	13 3		13+7	Arctic Region.	•	British Museum.
57‡	51	33½	44½ (outside)	20 + 16	Do. .		Sylvester Browne.
548	48	201		11+13	N. America .		The late Captain Sir John Franklin, British Museum.
-54	6 3	211	32‡	12+11	?		Paris Museum.
$52\frac{1}{2}$	5‡	15	26½	17+12	N. Canada .		J. Talbot Clifton.
52	4#	$28\frac{1}{2}$	38	7+7	Do		David T. Hanbury.
151 8	47	40 ⁷ 8	48 3	17 + 14	Hudson Bay .	•	Sir Edmund G. Loder, Bart.
49	41	248	328	7+9	N. America .	•	Hon. Walter Rothschild.
² 48	6 <u>1</u>	11 2	25½	17+15	Newfoundland		Sir Edmund G. Loder, Bart.
47 ⁸	5 1	32 1	•••	19+12	N. America .		A. Murray, British Museum.
46 1	6 <u>1</u>	161	28	32	British Columbia		J. Turner-Turner.
45½	5½	314	331	13+12	Newfoundland		F. C. Williamson.
45½	41	19‡	31	15+13	N. Canada .	•	David T. Hanbury.
448	5 <u>1</u>	34	37		Newfoundland		C. H. Akroyd.
44	5 1	18‡	30	18+13	Do		Vice - Admiral Sir William Kennedy.
44	54	19	26½	17 + 14	Do		Lieut. A. E. Allgood, R.N.
434	5 8	40	417	16+11	Do		St. George Littledale.
$-42\frac{1}{2}$	71	33 1	38 1	22+18	Do		R. Gordon Smith.
-42½		•••	49 1 (outside)	24	Do	•	General R. L. Dashwood.
42	$6\frac{1}{2}$		181	33	Alaska		Earl of Lonsdale.
42	51/2	36½	391	20+12	Newfoundland		Q. C. Colmore.
41	51/2	201	28	15+9	Do		LieutCol. Hon. W. Coke.
41	6	35	33	20	British Columbia		Major C. C. Ellis.
40	6	311	351/2	19+14	Newfoundland	•	Vice - Admiral Sir William Kennedy.
39	58		23‡	17+14	Do		Sir Victor Brooke's Collection.
39	6	31	30½	15+10	Do		Captain C. E. Stracey.
39	7	28	31	20+19	Do		Vice - Admiral Sir William Kennedy.
		1 W	oodland C	aribou.		3 Ba	arren-ground.

REINDEER or CARIBOU (Rangifer tarandus)—continued.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.	Owner. :
39	7	29	314	18+16	?	Duke of Westminster.
39	5≇		•••	11+11	British Columbia	Sir Peter Walker, Bart.
39	44	26	26 <u>2</u>	15+15	Newfoundland	D. F. Moir, R.N.
-39	7 8	33	31 2	29+22	Do	Lord Thurlow.
38	54	191	$21\frac{1}{2}$	17+14	British Columbia	T. P. Kempson.
38	6	36	351	39	Newfoundland	Lieut. F. C. Osborne, R.N.
371	6 1	23	23	11+10	British Columbia	W. S. Power.
371	54	12	20	14+12	Canada	J. W. Osborne.
37½	5 1	•••	29	•••	?	Otho Shaw.
37‡	5∰		24	27	Newfoundland	Dr. Wm. Tait.
37	51	21½	25	12+12	British Columbia	H. G. Walker.
-37	•••		40 (outside)	43	Newfoundland	General R. L. Dashwood.
37	41	142	25	15+13	Do	Captain H. H. Grenfell, R.N.
36 1	51	28	25	8.9	British Columbia	H. G. Walker.
36 1	51	19	22	10+9	Newfoundland	V. L. A. Campbell, R.N.
36 <u>1</u>	6 <u>3</u>	30	314	19+18	Do	A. Wilson, R.N.
36 1	6 1	26 1	28½	17+15	British Columbia	Sir Peter Walker, Bart.
36	58	44		15+12	Do	J. V. Colby.
36	5	23	271	12+11	Do	Captain F. Molyneux.
351	6 2	184	29½	13+11	New Brunswick	P. N. Graham.
351	5 1	321	34½	12+10	Newfoundland	G. C. W. Crispin, R.N.
3 3 ½	54	23	26	15+11	British Columbia	Sir Peter Walker, Bart.
-32½	81	12‡	18	17+10	Canada	James J. Harrison.
321	5 1	32½	•••	12+12	?	Major R. Hallowes.
-32	51/2			14	British Columbia	Theodore Roosevelt.

REINDEER or CARIBOU (Rangifer tarandus)—continued.

Length on out- side curve.		Tip to Tip.	Widest inside.	Points.	Locality.	Owner.
¹ 60	5₩	385	411	22+15	?	Sir V. Brooke's Collection.
-59 1	7	38	44	15+16	Norway	J. Whitaker.
-59	41/2	421	46 (outside)	7+5	Do	H. J. Elwes.
-58	6	30½	37	33	Jotunheim, Norway.	S. Ratcliff.
-551	6 <u>1</u>	38	431	18+8	Sundal Fjelds, Nor- way	Capt. Gerard Ferrand.
54½	48	33 1	40 <mark>8</mark>	15+13	Norway	J. H. Thomas.
54	5	22	413	16+11	Do	Kenneth M'Douall.
-51	$6\frac{1}{10}$	•••			Rundane, Norway .	Capt. John Marriott.
-51	54	29		25	Norway	Abel Chapman.
50 8	48	24	29 (outside)	16+13	3	H.R.H. the Duke of Saxe- Coburg and Gotha.
-49½	48	20 <u>8</u>	344	13+12	Norway	H. J. Elwes.
49	41	25	36 1	18+15	Do	J. H. Barnard.
♀47	48	23	29	17+9	Do	Kenneth M'Douall.
46 <u>1</u>	6	211	26 <u>1</u>	37	Do	Hon. Walter Rothschild.
-46	5	29	38	42	Jotunheim, Norway.	S. Ratcliff.
44	41	201	321	12+9	Norway	Col. C. B. Harvey.
-42	6	26	40 3	22+13	Do	R. Rankin.
41 8	4 8	•••	24½	12+9	Spitzbergen	W. D. James.
41	41/2	26 1	30 8	12 + 12	Do.	Sir W. Martin Conway.
-40½	41/2	29	30	11+8	Norway	Lewis J. Cadell.
382	4	25 1	•••	7+5	Do	British Museum.
38	5₺	21½		15+13	Do	Do.
37	41	18	20½	15+11	E. Spitzbergen .	Arnold Pike.
36	41	24	26	17+11	Do	Do.
-36	6 1	34	33½	23+20	Lapland	Dublin Museum.
⁹ -34 ¹ / ₂	6₹	31		33	Norway	Abel Chapman.
-31	51	23	20	19+18	Do	J. Benett-Stanford.
		1 Pe	rished antle	rs.	² A very o	ld buck.



Skull and Antlers of Male Elk.
From an Alaskan specimen in the possession of the Duke of Westminster.

ELK or MOOSE (Alces machlis).

The largest member of the deer tribe, distinguished by its ungainly form, long limbs, broad, produced, and flabby muzzle (all of which, except a small triangular patch below the nostrils, is covered with hair), the presence of a pendulous hairy organ (the so-called "bell") on the throat of the males, and the form and position of the antlers in that sex. These latter are set on the skull with their bases at right angles to the middle line of the face, the beams having neither brow nor bez tines, but expanding after a short distance into a broad palmation, carrying a number of snags on the outer border; in young elk each antler is divided in a fork-like manner into a small front and a larger hind portion. The main hoofs are long and pointed, and the lateral pair large: there is a gland and tuft of hair both on the hock and hind cannon-bone, the latter being situated high up. The tail is very short. From birth to old age elk are uniformly coloured; the general tint of the hair, which is long, coarse, and somewhat brittle, varying

from yellowish gray to deep blackish brown, and being usually darker in American than in European examples. The height varies from 5 feet 9 inches at the shoulder in Scandinavian examples (Sir H. Pottinger) to as much as 6 feet 6 inches in American specimens (General R. L. Dashwood); the weight from 900 to 1400 lbs., that of the antlers being about 60 lbs.

Elk inhabit the forests and marshy districts of Scandinavia, Eastern Russia, Siberia, Northern Russia, and thence eastwards through Siberia



Antlers of Male Elk.

north of about latitude 50° N. to Amurland; while in America (where they are invariably known as moose) at the present time they are found in Alaska, Montana, Nova Scotia, and New Brunswick. Although many sportsmen say they can be distinguished by the antlers and the colour of the skin, the European and American elk appear so similar that they do not seem worthy of being regarded as even distinct local varieties, but for convenience in comparing the European and American measurements are separated.

An elk killed at Meswiez, Lithuania, by Count Scheibler measured

							Feet.	Inches.
Length	tip	of nose	to	tip o	of tail		9	$\frac{1}{2}$
••					root		8	6

					Feet.	Inches.
Crest to nose.		•			2	7
Height at withers			•		5	9
" " quarters					6	$5\frac{1}{2}$
Girth quarters.						193
Round thigh .						$7\frac{1}{2}$
Below knee .					3	$6\frac{1}{2}$
Round neck near th	he e	ars			4	9
Estimated weight	15	oo lbs.	Se	e antl	ers, 27 1	, etc.
Circum-	Jak					


Length to longest tine.	Circum- ference above burr.	Tip to Tip.	Greatest width.	Breadth of Palm.	Points.	Locality.	Owner.
1-55½	7 ½		70 1	29 ?		Alaska	W. W. Hart.
-55	11 ?	37	69	21	21	Do	F. B. Tolhurst.
-49	•••		78 <u>1</u>	18	40	Yukon	W. F. Sheard.
²-48	10		69	15	32	Do	Dall De Weese.
-474	9‡	371	72	16‡	20+17	Do	Duke of Westminster.
441	101	49½	66	14	17+14	Kenai Mts	Viscount Powerscourt.
-44			611	•••		Canada	General R. L. Dashwood.
43‡	81	40	59 1	14½	14+13	Manitoba	G. H. M. Banks.
43	91	22½	471	12	11+11	Canada	Viscount Powerscourt.
-42 3	10	35‡	56	13‡	12+11	N. America .	T. W. Wood, Jun.
-41	81	•••		214	27	Chesincook, M.E.	Col. Hoselton.
-41		•••	54½	•••	12+11	Canada	General R. L. Dashwood.
-41	•••	•••	65	24	14+13	Manitoba	Otho Shaw.
40	8	38	60	13	11+10	Canada	Viscount Powerscourt.
40	8	35	55½	13	12+11	Maine	J. S. Braithwaite.
39 8	61		51 8	15#	13+12	N. America .	British Museum.
-39 1	81	•••	391	13\$		Do	Otho Shaw.
$39\frac{1}{2}$	7	38	54½	16	13+12	Canada	Sylvester Browne.
391	7 3	37	56]	131	14+11	British Columbia	Sir Peter Walker, Bart.
39 1	7 5	33½	498	111	12+10	N. America .	J. Carr Saunders.
391	7 1	36 1	512	9 1	8+7	E. slopes of the Rockies	J. C. L. Knight-Bruce.
39	81	381	57	12½	15+12	N. America .	Sir Edmund G. Loder, Bt.

 $^{^1\,}$ Height at shoulder, 8 ft. 2 in.?; skull horns, 68 lbs.; estimated weight, 2600 lbs. $^2\,$ Height at shoulder stated to be 7 ft. 8 in.?

ELK or MOOSE (Alces machlis)—continued.

Length to longest tine.	Circum- ference above burr.	Tip to Tip.	Greatest width.		Points.	Locality.	Owner.
381	7	•••	52 7	107	11+10	N. America .	British Museum.
		•••	67		23	Yukon	J. H. Whitehouse.
-38½	91/2	44	66	14	28	New Brunswick	S. Decatur.
38 1	6 <u>1</u>	34	49½	9 3	9+9	N. America .	Sir Victor Brooke's Collection.
38	7 §	30½	47	12	12+12	British Columbia	Sir Peter Walker, Bart.
38	6 <u>1</u>		59 1	93	11+11	?	H.R.H. the Duke of Saxe- Coburg and Gotha.
38	8	211	57½	14	18+15	Manitoba	C. H. Akroyd.
-38	11	50	61	14	26	New Brunswick	J. Bodkin.
-37 3	8	•••	51 1	111	14+13	Canada	F. Ashby.
$37\frac{1}{2}$	81	•••	49½	11	21	?	Ernest Farquhar.
371	7 1 8	35	50 §	115	13+10	New Brunswick	Hon. Charles Ellis.
-35 3	71/2	331	474	11	12+11	Canada	James J. Harrison.
$-35\frac{1}{2}$	81	•••	42] 8	•••	10+9	Do	Paris Museum.
34½	8	39 1	49 1	101	9+9	New Brunswick	P. N. Graham.
34 1	6	42	51 1	$9\frac{1}{2}$	10+9	Canada	G. Marchetti.
-33½	7	28	44	11	13+13	N. America .	Dublin Museum.
33	6	30	44	10	10+10	Manitoba	Major C. S. Cumberland.
$32\frac{1}{2}$	7	294	44	8	11+9	N. America .	Charles Makin.
-31\$	7		•••		14	Do	A. Rogers.
311/2		•••	51½	121	•••	?	Earl of Lonsdale.
-30	51/2	•••	40½	13	22	Montana	Theodore Roosevelt.
283	78	35	49 8	121	11+11	New Brunswick	Hon. Charles Ellis.
27	5 7	28	42	6₹	6+7	Eastern slopes of	Hon. F. Thellusson.
26 <u>1</u>	6 3	30	40	6	6+6	the Rockies N. of Manitoba	P. B. Vander-Byl.

ELK or MOOSE (Alces machlis)—continued.



Antlers of Male Elk.

Length to longest stine.	Circum- ference above burr.	Tip to Tip.		Breadth of Palm.	Points.	Locality.	Owner.
37 3	8	358	57\$	1112	11+13	Norway	Thomas Bate.
33	$8\frac{1}{2}$	36 1	513	15‡	10+10	Do	Capt. Gerard Ferrand.
-33	9	34	35		9+10	E. Prussia .	H.R.H. the Duke of Saxe- Coburg and Gotha.
$32\frac{1}{2}$	$7\frac{1}{2}$	33	46	I I ½	10+10	Sweden	Capt. Gerard Ferrand.
32	6		438	91	7+8	Russia	Sir Edward Caley, British Museum.
32	83	37½	48	11½	10+9	Nr. St. Peters- burg	Prince Demidoff.
-32	7	•••	52	9	9+9	Norway	H. J. Elwes.
	7	35	42½		9+7	Do	J II. J. Liwes.
-31	7	28	45		17	Do	S. Ratcliff.
¹ 30 ³	$6\frac{1}{2}$	43	46	103	15	Do	Abel Chapman.
30½	8	27	40½	91	10+7	Do	LieutCol. G. D. F. Sulivan.
30 1	7 1	29½	441	10	9+8	Do	Sir Victor Brooke's Collection.
29	6	30½	42½	$7\frac{1}{2}$	11+9	Do	Sir H. Pottinger, Bart.
29	$6\frac{1}{2}$	311/2	39 1 ₂	$6\frac{1}{2}$	10+9	Do	Do.
29	71/2		$43\frac{1}{2}$	101	24	Do	Sir Peter Walker, Bart.
$28\frac{1}{2}$	71/2	243	42	9½	10+9	Do	Sir H. Pottinger, Bart.
				1 Esti	mated weig	ht, 90 stone.	

ELK or MOOSE (Alces machlis)—continued.

Length to longest tine.	Circum- ference above burr.	Tip to Tip.	Greatest Width.	Breadth of Palm.	Points.	Locality.	Owner.
-28½	6	33 1	43 1	11	12+10	Lithuania .	Prince Radziwill.
28	6 1	32	47	9	8+8	Norway	Sir H. Pottinger, Bart.
28	7 1	311	38	10	12+10	Sweden	Viscount Powerscourt.
-27½	7	25½	33	82	5+5	Lithuania .	Count Scheibler.
-27	6 <u>3</u>	30	39		20	Norway	S. Ratcliff.
27	6 <u>1</u>	40	43½	•••	8+7	Do	J. H. Barnard.
27	6 <u>1</u>	20½	38	9	8+6	Nr. St. Peters-	Prince Demidoff.
-26 8	81	28 <u>1</u>	412	9 l	8+8	burg Norway	Sir H. Pottinger, Bart.
261	6‡	25	38	6	9+7	Russia	LieutCol. Hon. W. Coke.
261	6 <u>1</u>		39 8	6 <u>7</u>	5+4	S. Russia	British Museum.
26	6	271	421	7	7+6	Norway	Lord Delamere.
26	6 <u>1</u>	29	37	5	6+5	Do. . · .	Sir H. Pottinger, Bart.
231/2	5 3		35%	5	5+3	Do	E. N. Buxton.
23	6	271	36	6	7+6	Sweden	Sir H. Pottinger, Bart.



Antlers of Caspian Red Deer. From a Hungarian specimen in the collection of Viscount Powerscourt. Counting from the skull upwards, the first tine is the brow, the second the bez, and the third the trez, above which come the surroyals, or crown. The main shaft is termed the beam.

RED DEER (Cervus elaphus).

The red deer of Western Europe is the typical representative of the genus *Cervus*, in which the antlers of the males are set on the skull at an oblique angle to the middle line of the forehead, and always have a brow tine, while they are generally more or less nearly cylindrical, although sometimes palmated. There is always a large bare portion on the muzzle, the face is long, the ears are generally large, and the tail is comparatively short, often extremely so. Although there is almost always a gland and tuft on the hind cannon-bone, usually situated high up, there is none on the hock itself. The coat may be spotted.

In the red deer the antlers are subcylindrical and complex, generally with a bez tine, and always with a trez, the number of points exceeding five, and the crown frequently forming a cup. The tail is relatively long and pointed, and there is a distinct light-coloured patch on the buttocks, which includes the tail; the general colour of the adult summer coat being reddish brown, and that of the winter dress grayish brown, while the young are profusely spotted.

Red deer, in the widest sense of the term, are inhabitants of Europe, North Africa, Asia Minor, and Northern Persia. In the typical red deer (C. elaphus typicus) of Western, Northern, and Central Europe, the antlers attain their maximum degree of complexity, sometimes having twenty or even more points, although, in many Scotch examples the bez tine is wanting. In a park red deer killed at Spetchley Park the weight was 419 lbs. gross, and 216 when cleaned; while in one shot at Knowsley many years ago the clean weight was no less than 424 lbs. The height at the shoulder now reaches about 4 feet. H.R.H. the Duke of Braganza saw a continental stag shot which scaled 41 stone 10 lbs. two years ago, and last year H.R.H. shot a 10-pointer with a spread of 55 inches.

The Corsican red deer (C. elaphus corsicanus), of Corsica and Sardinia, is a very small race, with the bez tine of the antlers wanting, and the general colour of the upper parts dark brown in summer and blackish in winter. Nearly allied is the North African red deer (C. elaphus barbarus), which is of rather larger dimensions, with a grayish-brown streak down the middle of the back, and small irregular whitish spots on the flanks and sometimes on the back; traces of such spots being occasionally observable in the summer coat of does of the typical race. The bez tine seems to be very generally wanting.

The Caspian red deer (C. elaphus maral) is a large variety, described on page 33.

Lord Tweedmouth furnishes the following dimensions of his fine Scotch stag:—

Widest span over all, $39\frac{1}{4}$ inches; span inside below cups, 34 inches; span outside below cups, 37 inches.

Right antler.—Length, 39 inches; length of brow, $10\frac{1}{4}$ inches; of bez, 10 inches; of trez, 13 inches. Length of tines in cup, 10, 7, $4\frac{1}{4}$ inches. Circumference at coronet, $8\frac{1}{2}$ inches; between bez and trez, $7\frac{1}{4}$ inches; above trez, 6 inches.

Left antler.—Length, 38 inches; of brow, 10 inches; of bez, 8 inches; of trez, 11 inches. Length of tines in cup, 8, 6, and 4 inches. Circumference at coronet, 9 inches; between bez and trez, $7\frac{1}{2}$ inches; above tray, $6\frac{1}{4}$ inches.

Shot, October 9, 1880. Weight, 21 stone 9 lbs.; clean, but stag was much run.



Antlers of Red Deer killed by Lord Burton, with fully developed cups.

a.—SCOTCH RED DEER.

Length on out- side curve.	Circum- ference between bez and trez.	Tip to Tip.	Widest inside.	Spread.	Points.	Weight.	Locality.	Owner.
-4I	7	$22\frac{1}{2}$	35		7+7	St. 1DS.	Inverness (1794)	W. H. Walker.
1-40	6		30		10	14.7	Glentilt .	Duke of Atholl.
-39	71		34	39 1	6+6	21.9	Guisachan .	Lord Tweedmouth.
-378		20 3	26½	•••	7+7	•••	?	Sir Humphrey de Trafford, Bart.
378	5	142	26		6+7	•••	?	H.R.H. the Duke of Saxe- Coburg and Gotha.
374	5	143	24 ²	•••	6+5	•••	?	Do.
1-361	•••		35	12	•••		Glenmoriston	Col. W. Gordon-Cumming.
36 1	5	34	39½	41	5+6		Glenstrath- farrar	Viscount Powerscourt (shot by the late Roualeyn Gordon-Cumming).
36 <u>1</u>	6 1	3112	32½	•••	6 +6		?	Sir Humphrey de Trafford, Bart.
36 1	7	22 3	28		7+7		?	Do.
-36			36	42	13		Monymusk	The late Capt. Johnstone Grant.
36	51/2	16 1	28	36	7+7	•••	(1795)	Duke of Portland.

¹ Recorded by J. G. Millais.

SCOTCH RED DEER-continued.

Length on outside curve.	Circum- ference between bez and trez.	Tip to Tip.	Widest inside.	Outside spread.	Points.	Weight.	Locality.	Owner.		
36	41	21½	27		6+6		Caenlochan .	Henry Tate.		
351	43	27½	34	36 8	4+3	•••	Do.	Do.		
351	5½		25½		6+6	17 6	Sutherland .	Abel Chapman.		
35 1	6			29	12	19	Auchnashel- lach	The late Lord Alexander		
35¥	41/2	17½	30	•••	5+6	17 6	Glenisla, For-	Paget. Viscount Powerscourt.		
$-35\frac{1}{2}$	7	•••	26½		20	•••	farshire Glenquoich .	Lord Burton. (See illustration.)		
35½	43	271	34	36 8	4+3		Caenlochan .	Henry Tate.		
-35 1	48	31 17	30		6+6		Deanich, Ross- shire	G. Percy V. Aylmer.		
35	4 1	22	28		7+6	•••	Talladh-a-Bhe- ithe, Rannoch	E. Weller-Poley.		
-35	•••	•••	•••	34	10	16 I	N. Morar .	Capt. T. W. Gill.		
-35		•••	29	411	7+6	•••	?	Sir Arthur Grant, Bart.		
1-35	54	26	33½	$37\frac{1}{2}$	12	•••	Guisachan .	Lord Tweedmouth.		
² -35	41/2	•••	•••	$27\frac{1}{2}$	8	•••	Cairn Thomais Gaick	John Hargreave.		
-34 1	•••	•••	•••	311	3+2	16 12	Cluanie .	Hon. T. A. Brassey.		
34½	5 1	37½	341		12	•••	Braemore .	Sir John Fowler, Bart.		
34½	41	18 1	27		6+5		N. Morar .	J. R. Hutchison.		
² -34½	5	•••		341	12	16 7 (clean)	Glentilt .	Duke of Atholl.		
34½	6	173	25	314	5+4	15 12	Caenlochan .	Mrs. Henry Tate.		
34‡	418	16‡	248		5+5	•••	Auchnasheen	F. Devas.		
-34	$5\frac{1}{2}$			30	Switch	•••	Letterewe .	Hon. T. A. Brassey.		
34	41	172	26 1	29	5+4		Caenlochan .	W. A. L. Fletcher.		
34			30	•••	11	17 2	N. Morar .	Capt. T. W. Gill.		
34	4	28 <u>1</u>	303		5+4		Glenmuick .	J. Ponsonby.		
34	41	12	23½	•••	5+4	•••	?	C. A. Grenfell.		
34	4	16 8			5+4	•••	Blair-Atholl	Captain A. W. E. Count Gleichen.		
-34	48	19	28	•••	6+5	•••	Rothiemurchus, Inverness	A. Basil Brooke.		
² -34	•••	•••			11	•••	Braemar .	Col. Gordon-Cumming.		
² -34	41	•••	•••	36½	12	•••	Glenbruar, Perthshire	Sir W. Ogilvie Dalgleish.		
			1 Abov	e trez.		² Re	² Recorded by J. G. Millais.			

C

SCOTCH RED DEER-continued.

Length on outside curve.	Circum- ference between bez and trez.	Tip to Tip.		Outside spread.	Points.	Weight.	Locality.	Owner.
² -34	48			30	10		Morar	W. Stirling.
² 34	7			26	14		Glenartney .	Earl of Ancaster.
33 3	5 ½	191	25½		5+5		?	H. S. O'Brien.
-33½	5		34½		5+5	20	Dalness .	J. G. Millais.
33½	4	16 <u>1</u>	25	29	5+5		Caenlochan .	H. C. Pilkington.
-30½	$4\frac{1}{2}$		27		12		Kintail	Sir Edmund G. Loder, Bart.
33	41	81	20 3	. 244	4+4	•••	Ben Alder .	Julius Wernher.
33	41	24 3	28 1	33½	5+4	15 12	Do.	F. C. Selous.
33	43	21½	28 8	33	4+4		Kintail	R. P. Page.
-32 3	5	25 3	29 1		6+6	•••	Ardverikie .	W. H. Walker.
323	41/2		28		5+4	•••	3	Sir Charles Tennant, Bart.
32½	41	16 1	26 <u>‡</u>	•••	5+5	•••	Auchnashel- lach	G. Webster.
32 1	4 1	203	28		5+5	•••	Morar	J. R. Hutchison.
32	4	25‡	25	28	5+5	•••	Invercauld .	Otto Beit.
32	5		22½	•••	5+4	19	Rannoch .	Sir W. G. Pearce, Bart.
¹ 3 ²	41/2	31	29½	40	7+9	•••	Rhidorrach .	Viscount Powerscourt.
32	4	24½	29	•••	4+4	•••	Braulen .	J. K. Fowler.
-32	•••	•••	•••	34	10	•••	Kintail	Sir Edmund G. Loder, Bart.
314	3 3	•••	26½	•••	4+4	•••	Rhidorrach .	Captain F. Cookson.
312	4 1	18	24 <u>3</u>		6+5	14	Inchgrundle.	Countess of Dudley.
311	$4\frac{1}{2}$	16	21	231	6+5	•••	Shank	R. K. Micklethwait.
311	4 3	22	28 ‡	314	3+3	•••	Caenlochan .	E. L. Fletcher.
311/2	4	121	21½	25 1	6+4	•••	Do.	W. W. Gossage.
311/2	5	•••	26 <u>1</u>	•••	12	17	Auchnasheen	J. F. Laycock.
311	4	22	26 1	$32\frac{1}{2}$	7+6	•••	Kintail	R. P. Page.
311/2	4	$24\frac{1}{2}$	29	•••	5+4	•••	Braulen .	J. Talbot Clifton.
311	4‡	21	25½	•••	4 + 4	15 3	Invermark .	W. S. M. Burns.
311/2	48	201	25 3	•••	3+3		Dalnacardoch	Hon. T. A. Brassey.

Above trez.

² Recorded by J. G. Millais.

SCOTCH RED DEER—continued.

	Circum- ference between bez and trez.	Tip to Tip.	Widest inside.	Outside spread.	Points.	Weight.	Locality.	Owner.
31 8	48	8 <u>\$</u>	24	•••	5+5		Glenmuick .	C. Lawrence.
-311	51/2	25½	29	•••	Switch	•••	Achdalien .	M. K. North.
314	48	16	24 1	•••	7+5	•••	Rhidorrach .	Capt. F. Cookson.
314	48	24 7	32		5+3		Ross-shire .	E. Lort-Phillips.
314	4	•••	40}		6+6	•••	?	Dr. Fancourt Barnes.
311	4	15	25		5+4	•••	Auchnashel- lach	W. Maxwell Lyte.
31	4\$	24	25	•••	5+5		Morar	J. R. Hutchison.
-31	51/2	$32\frac{1}{2}$	•••	38	5+4		Inverlochy .	Mrs. J. E. Platt.
-31	5 1	201	24	•••	10		?	C. V. A. Peel.
31	47	19‡	234	33 1	8+6		S. Harris .	Earl of Dunmore.
31	45	$24\frac{1}{2}$	28 1		5+4		Caenlochan .	F. W. Robinson.
-31	$4\frac{1}{2}$	20	15		8+7		?	J. Benett-Stanford.
31	41/2	16 1	$23\frac{1}{2}$	29	6+5	•••	Inversanda .	J. Hamilton Leigh.
30 7	48	25 7	28 1		5+6		Rhidorrach .	W. R. Cookson.
30 3	5	191	24		6+5	•••	Arisaig .	J. R. Hutchison.
30 2	41	13	23 1		5+4	14	Invergarry .	Guy Stephenson.
30½	3 1	24 1	31	33 1	3+3	•••	Inversanda .	J. Hamilton Leigh.
30½	41	131	$23\frac{1}{2}$	•••	6+6	•••	Letterewe .	Gordon Wood.
30½	4‡	174	24 <u>3</u>		7+6	•••	Rhidorrach .	J. Talbot Clifton.
30 1	41/2	181	24		4+3	14	Glen Tana .	Major Hon. A. H. Henni- ker.
30 1	4	25½	28‡		4+3		Auchnashel- lach	Capt. H. Reynolds, R.N.
30½	44	22]	29	31	3+3		?	L. Breitmeyer.
301	41	20	$25\frac{1}{2}$		6+5		?	H.R.H. le Duc d'Orléans.
-3 0	48		25½		5+5	15	Blackmount	J. G. Millais.
-29 8	4	$32\frac{1}{2}$			5+5	16 2	Do	Do.
29½	$4\frac{1}{2}$	24 1	29	•••	4+5	•••	?	Marquis Camden.
-29 1	6 <u>1</u>		27 ⁷ 8	•••		16 11	Blackmount .	Marquis of Breadalbane.
-29	4	•••	29	31	6+6	•••	Inverness .	H. Seton-Karr.

The widths of the six widest heads in the Duke of Fife's collection are as follows:—

The six longest antlers of this celebrated collection are respectively— $37 \quad 36 \quad 35\frac{1}{2} \quad 35\frac{1}{2} \quad 35$

The antlers of one of the red deer in Mr. Lucas's Park at Warnham Court, Sussex, in 1889, had 34 points; 1890, 34 points; 1891, 37 points; 1891, 47 points and weighed 17 lbs.; 1893, 45 points, $16\frac{1}{9}$ lbs.

For an interesting account of many fine specimens the reader is referred to *British Deer and their Horns*, by J. G. Millais.

b.—IRISH RED DEER.

Length on outside curve.	Circum- ference between bez and trez.	Tip to Tip.	Widest inside.	Points.	Weight.	Locality.	Owner.
1-42½				12	22 7	Colebrooke	Sir Douglas Brooke, Bart.
-4I	6	171	23	10+9	23 3	Do	Do.
-40	51/2	28	29½	8+8	25 5	Do	Do.
39 8	5‡	$14\frac{1}{2}$	26§	7+7	26 11	Do	Do.
38	58	20	30§	10+8	25 O (clean)	Do	Do.
35	43		30	9	(clean)	Ireland	Hon. A. Charteris.
35	5 1	198	26 3	6+5	20 2	Powerscourt Park	Viscount Powerscourt.
35	5‡	211	29	5+5	24 st. as he fell 18 st. clean	Do	Do.
34½	5	I 2 ½	22½	5+5	24 st. as he fell 18 st. clean	Do	Do.
34	41/2	25½	28 1	6+6	26 st. as he fell 20 st. clean	Do	Do.
34	5	20½	26	6+6	20 st. clean	Do	Do.
34	44		31 1/2	6+5	26 8	Muckross	Ralph Sneyd.
34	41	13 1	25	5+5		Colebrooke:	Capt. J. M. Rogers.
33 2	5	•••	26	5+5	22 0	Muckross	Hon. Mrs. Bourke.
31	4 3	11	20 ³	7+7	•••	Do	Ralph Sneyd.
-31	43	. 181	22	5+7	•••	?	Dublin Museum.
30 1	4		26½	5+5	19 10	Muckross	Geoffrey Carr-Glyn.
293	4 3	211	$24\frac{1}{2}$	6+ 6		Colebrooke	Montrose Cloete.

¹ See Millais's British Deer and their Horns.



Antlers of Exmoor Stag.

c.—WEST OF ENGLAND RED DEER.

Length on out- side curve.	Circum- ference between bez and trez.	Tip to Tip.	Widest inside.	Points.	I	ocal	ity.	(Owner. Sir John Heathcoat-Amory, Bart.;
¹ 41	51	211	321	5+5	Exmoor	٠.		-{	killed in 1897 with the Devon and Somerset Stag-hounds.
39	53	152	281	6 + 6	Do.				R. A. Sanders.
381	5 ² 5 ¹ 5	178	31]	6+6	Quanto	ck I	Iills		Viscount Ebrington; killed in 1885.
371	5	221	30 1	7+7	Exmoor				C. Nelder; killed in 1803.
36 3	6	21½	28 2	6+7	Do.	•	•	•	Sir A. Acland - Hood, Bart.; killed in 1893.
35½	5	20 ž	27 ½	6+6	Do.				Earl Fortescue; killed in 1812.
35	5 5	154	27	6 + 5	Do.				Com. G. F. Inglefield, R.N.
35	48	25 <u>i</u>	323	5 + Š	Do.		•	•	Sir C. T. D. Acland, Bart.; killed in 1893.
348	5∰	23 1	311	6 + 6	Do.				Viscount Ebrington; killed in 1881.
² -33	5# 6½		29	6+6	Do.	٠	•	•	The late Sir T. D. Acland, Bart.; killed in 1788.
3 -33	5 ‡		29 1	9+7	Do.				,, ,, 1792.
-33	5 8	39	32	7+7	Do.				,, ,, ,, 1877.
33	ξŝ	13	25	8+7	Do.				Viscount Ebrington; killed in 1881.
32	5 1	30∄	30‡	7+4	Do.				Earl Fortescue; killed in 1814.
-33 33 32 32	5 1 5 1 4 1 2	152	26	6+6	Do.				Col. J. F. Hornby.
30]	5	24½	29½	6+6	Do.				Earl Fortescue.

¹ Weight 333 lbs., clean. Length of brow-tine, 17 inches. 2 Weight, 17 stone 2 lbs. 3 See Red Deer in Fur and Feather Series.



Head of Stoke Park Red Deer.

d.—ENGLISH PARK RED DEER.

Length on outside curve.	Circum- ference between bez and trez.	Tip to Tip.	Widest inside.	Spread.	Points.	Weight.	Locality.	Owner.
$-42\frac{1}{2}$	6‡			40	12			Earl of Ilchester.
411	5	1 223	32		7+6		Langley Park	J. G. Millais.
-40				41	7+7		. 5	Sir Greville Smyth, Bart.
$-37\frac{1}{2}$	7 ½	29½	27 ½	36½	10+9		Welbeck .	Duke of Portland.
-37	74			37	23	31 st. (clean)	Warnham .	T. Lucas.
-36	6 <u>1</u>	24	30	•••	6+6	•••	Vaynol, North Wales	G. W. D. Assheton- Smith.
36	1 5½	19	281		9+11		Woburn .	Duke of Bedford.
-35	6	141	24		6+6		Vaynol, North Wales	G. W. D. Assheton- Smith.
$34\frac{1}{2}$	7	33	26₹		20 + 20		Warnham .	W. H. Lucas.
341	1 5 1	181	26		8 + 8		Woburn .	Duke of Bedford.
34	$4\frac{1}{2}$		23		6+6	••	Stowe	H.R.H. la Comtesse de Paris.
34	5∰	213	281		7 + 7		Dorset	Earl of Ilchester.
34	6 <u>3</u>			48 (about)	45		Warnham .	C. T. Lucas.
$33\frac{1}{2}$	4 🖁	•••	33		6+6		Stowe	H.R.H. le Duc d'Orléans.
33½	5‡	211	28 1		9+9		Woburn .	Duke of Bedford.
33	7 above			451	44		Warnham .	C. T. Lucas.

Some of the above measurements are recorded by J. G. Millais in Britisli Deer and their Horns.

¹ White Stag.



Skull and Antlers of Old English Red Deer. (Found in cutting the Manchester Ship Canal.)

e.—ANCIENT BRITISH RED DEER.

Length on outside curve.	Circum- ference between bez and trez.	Tip to Tip.	Widest inside.	Spread.	Points.	Where di	ug up.		Owner.
$-47\frac{1}{2}$	8	35			5+8	Mancheste Canal exc		hip	Sir R. M. Brooke, Bart.
40	71/2	22‡	$28\frac{1}{2}$	43‡	12+9	Comberme		·	Duke of Westminster.
$38\frac{1}{2}$	5	18	30	39 1	8+6	Ireland			Viscount Powerscourt.
36½	5	24	25	35½	8+8	Do.			Do.
36	5½	23	27	$37\frac{1}{2}$	10+9	Kerry, Irel	land		Do.
¹ 36	57		•••	421	23	South Irela	and		Sir Douglas Brooke, Bart.
$35\frac{1}{2}$	5‡	26½	32 8	423	13+11	Do.			Sir Victor Brooke's Collection.
35‡	5 1	31‡	273	•••	12+10	Ireland			Viscount Powerscourt.
-35	7		21	24	9+11	England			C. G. Burrow.
$34\frac{1}{2}$	5	22	30		10+10	Ireland			Viscount Powerscourt.
33	44	27	281	35 1	9+8	Do.	•	•	Do.

¹ Recorded by J. G. Millais.

f.—NORWEGIAN RED DEER.

"Hitteren may still be considered the headquarters of the red deer in Norway. These animals (unlike the elk, which would appear to be diminishing) are increasing in numbers, and are now found in districts where they were previously unknown. Of 138 killed in the whole country last year fifteen were shot in South Bergenhus, twelve in Romsdal, and thirty-one in North Bergenhus Amt" (SNOWFLY, Field, 11th December 1897).

Length on outside curve.	Circum- ference between bez and trez.	Tip to Tip.	Widest inside.		Points.	Weight.	Loca	lity.		Owner.
-34	•••	•••	•••		7	•••	Norway			J. II. Thomas.
312	4 1		25½	•••	6+5		Do.			A. Brassey.
311	41/2		24½	•••	5+5		Do.			II. Seton-Karr.
311/2	4 1	•••	28		4+3		Do.			E. M. Denny.
31	4‡	23	25	291	6+6		Do.			J. H. Thomas.
30	5		29		12	20 st.	Do.			E. M. Denny.
-30	41		28	33	5+4	•••	Do.			II. Seton-Karr.
30	44	29 1	303	•••	5+4		Do.			Do.
29½	47	18 1	231		5+5		Do.			J. H. Thomas.
29 1	48	2 6	30 1		6+5		Do.	•		G. L. Denmar.
28 1	4	27	25		4+4		Do.			H. F. Kemp.
-2 8	51	Sin	gle An	ller	5	•••	Do.	•		H. Seton-Karr.
$27\frac{1}{2}$	5	26	26		5+6		Do.			J. II. Thomas.
26 1	4‡	19‡	214		4+4		Do.	•		Do.
26	41/2	•••	30	$3^{2\frac{1}{2}}$	6+5	20 st. 9 lbs.	Do.	•		H. Seton-Karr.
24 ³	31/2	21	201		2+2		Do.	•		A. Henderson.
				g.—8	PAN]	ISH R	ED DEI	ER.		
1-40	•••		36 1	•••	17	Sierra	Morena,	Spair	1	Abel Chapman.
$-37\frac{1}{2}$			34½	•••	15	Sierra	Morena			Do.
. 36	51/2	37	•••		15	South	Spain .			The late Lord Lilford.
$-32\frac{1}{2}$	•••	•••	•••		13	Plains	of Anda	lucia .	•	W. J. Buck.
-29	5‡	•••	25		12	Andal	lucia .		,	Abel Chapman.
-28½	5‡		26 1		13 1 A n	Do nountain h		•		Do.

h.—other continental red deer.

Some of the specimens in this list from Eastern Europe probably belong to the Maral, or Caspian, race.

Owner.	H.R.H. D. Miguel, Duke of Braganza.	Count Gèza Andrassy.	E. N. Buxton.	Count Gèza Andrassy.	Viscount Powerscourt (a).	Do.	J. J. S. Whitaker.	Duke of Ratibor.	Prince Lulu Rohan.	Count Joseph Hoyos.	Prince Philip of Saxe-Coburg	Prince Henry of Liechten-	Prince Hugo Windischgraetz.	LieutCol. L. Marshall.	Prince Henry of Liechten-	H.R.H. the Duke of Saxe-	Viscount Powerscourt (c).	Do.	
Locality.	Galicia .	Zemplen	Galicia .	Zemplen	Hungary .	Do	La Mandria	Pilis Moun-	Radauc .	Hungary .	Do	Carpathians	~	Carpathians	Do.	Gotha .	Germany .	Hungary .	
Weight of Stag.	206 kilos or 32 stone	5 to. 250 kilos	36 stone,	cut up 236 kilos	:	:	:	:	:	354 lbs.	÷	÷	:	:	:	:	:	÷	
Spread.	÷	413	45	47	45.25	20	÷	55.9	÷	483	:	43	:	373	÷	84	\$1\$	28	
Widest inside.	:	32	32.75	38	42.5	39.5	\$:	398	:	÷	;	:	8	:	27	414	43	
Tip to Tip.	:	114	÷	27	38.5	23.5	42	30	;	:	:	:	:	£	:	:	33‡	38	ģ
Circum- ference between Bez and Trez.	0	8.75	7.25	7.20	5.5	7-25	9	:	:	:	:	:	:	₹9	i	:	6 3	7	1 Due un
Circumfer- ence below Crown.	15½	8.20	:	6	:	:	;	:	:	98	:	:	:	:	:	:	:	:	
Circumfer- ence above Middle Point.	:	7.28	:	7	:	:	:	:	:	8 ₹9	:	:	:	:	:	:	÷	:	
Circum- ference above Burr.	0	8.47	:	ĩ 8	÷	:	:	90-9	:	974	:	:	:	:	84	6	:	:	
Circum- ference of Burr.	† 11	10-04	÷	11	:	:	:	:	:	ro ‡ or	$10_{ m I^3_0}$:	:	:	:	:	:	:	
Length along Curve.	531 ⁹ 4	53.54	52	51	15	Şı	50	50-4	493	49 <u>\$</u>	49	49	48.8	483	48	84	48	84	
Number of Weight in Ibs. Points. Avoirdupois.	ov er 2 0	23.36	20.5	23	:	:	:	21.3	31 and an		21	:	25.3	:	:	:	:	:	
Number of Points.	- 18	91 -	- 18	-14		1 14	- 14	- 21	- 14	- 18	- 12	91 –	- 20	14	- 19	61 -	20	118	

1 Dug up.

	Owner.	. H.R.H. D. Miguel, Duke of	Prince Philip of Saxe-Coburg	Viscount Powerscourt.	. Count Ferdinand Trautt-	A. von André.	. Viscount Powerscourt (a).	. Count Fery Nádasdy.	. Prince Victor Ratibor.	. Count Max Hoyos.	. Count Mittrovszky.	. Count Bela Szechenyi.	ns Prince Henry of Liechten-	. Viscount Powerscourt (e).	ny Do. (d).	. H.R.H. the Duke of Saxe-	Viscount Powerscourt.	is Prince Altenburg.	Count Erbach.	. Major-Gen. Sir Arthur Ellis, K.C.V.O.	. Sir Edmund G. Loder, Bart.
	Locality.	Galicia	Hungary	Do.	Do.	Galicia	Germany	Hungary	Do.	Do.	Do.	Do.	Carpathians	Hungary	N. Germany	Gotha	Hungary	Carpathians	Bukowina,	Hungary .	Do.
	Weight of Stag.	37 stone	:	i	242 kilos	÷	:	÷	÷	479 lbs.	:	:	÷	:	:	:	:	:	:	:	:
ontinued.	Spread.	32	÷	30	291 ₈	÷	÷	:	÷	42 Å	:	÷	÷	53	:	8	₹05	37	543	÷	44
DEER-continued.	Widest inside.	. 25	÷	9	254	313	384	÷	:	39‡	:	÷	36	424	31	36	$39\frac{1}{2}$	₹ 6z	35	38‡	÷
RED	, Tip to Tip.	13	:	30	354	124	23	:	÷	52¥	:	:	:	38	194	:	32	:	:	263	:
CONTINENTAL	Circum- ference between Bez and Trez.	7.3	:	5‡	÷	‡ 9	6 3	:	:	:	:	:	:	ę į	73	:	53	7.4	፥	\$ 9	:
	Circumfer- ence below Crown.	12	7.87	:	84	:	:	7.87	68.9	8+8	7.28	8.46	÷	:	÷	÷	:	:	:	÷	:
OTHER	Circumfer- ence above Middle Point.	7.4	7.87	:	78	:	:	7.09	8.07	₹9	6-891	69-9	÷	÷	:	÷	:	÷	:	:	:
	Circum- ference above Burr.	₹6	9-25	:	814	÷	÷	7.87	9-45	9 1	9.25	9-05	:	:	:	6	:	:	:	:	:
	Circum- ference of Burr.	01	11.21	:	ΙΙĄ	÷	:	9.84	11.02	914	18-11	10.63	÷	:	6	:	፥	:	01	÷	:
	Length along Curve.	48	47.84	47 §	474	47	47	46.46	46.46	46	46.06	46.06	46	46	46	46	46	46	46	458	45 <u>å</u>
	Number of Weight in lbs. Points. Avoirdupois.	23	20-57	:	:	:	:	20.79	20.68	:	16-61	22.55	÷	:	÷	:	:	:	:	÷	:
	Number of Points.	- 14	91 -	1.5	91 -	13	∞	- 20	91 -	- 22	91 -	- 20	- 14	91	13	81 -	15	- 13	81 -	01 -	÷

. Viscount Powerscourt (/).

÷ 65

454 454

Count George Erdody.	Count Belà Szechenyi.	Prince Demeter Ghika.	Prince Henry of Liechten-	Viscount Powerscourt (b).	Do.	H.R.H. the Duke of Saxe.	E. N. Buxton.	Viscount Powerscourt (g).	Major-Gen. Sir Arthur Ellis,	Count Tassilo Festetics.	Count Belà Szechenyi.	Count Michael Esterhazy.	Jeno Kund.	Viscount Powerscourt.	Do. (h).	A. von André.	Count Rudolph Erdödy.	F. Pausinger.	Count Tassilo Festetics.	A. von André.	Count Tassilo Festetics.	Prince Philip of Saxe-Coburg	Viscount Powerscourt (/).	Do. (i).
Do	Do.	Roumania.	Hungary .	Germany .	Ωο.	Gotha .	Galicia .	Carpathians	Hungary .	Do.	Do	Do	Do.	Germany .	Do.	Hungary .	Do	Do	Do.	Do	Do.	Do.	Germany .	Do
:	:	:	:	:	:	÷	:	÷	:	÷	÷	÷	:	:	:	:	÷	:	:	÷	:	:	:	:
:	:	:	÷	:	43	÷	46-5	፥	:	:	÷	:	÷	\$2\$:	:	· . :	:	:	36	:	:	:	:
:	:	34‡	34‡	37	36 §	33	39	284	32	÷	:	:	:	39‡	33	39	:	:	25.12	291	:	:	38‡	33
;	:	23₿	26 ig	23	23₺	38	:	143	17	:	:	:	:	31	23‡	243	:	:	36-06	15	:	:	374	284
:	:	:	s	F 9	54	:	9	74	7.4	:	:	:	:	74	₹ 9	53	:	:	7.64	† 9	:	÷	F 9	† 9
2.48	7.48	:	1410	i	÷	:	:	:	:	12.21	8.07	2.68	68.9	:	÷	:	9.64	7.28	12.23	5#	8.27	7.28	÷	:
7.28+	69.9	:	7	÷	:	:	÷	:	:	9-05	6-49†	7.68†	7.28	÷	÷	:	7.09†	7.09†	÷	† *9	7.28	60-1	:	:
8.07	6.49	:	₩	:	:	6	:	:	÷	7.87	9-45	9-84	10-24	:	÷	÷	8.46	8.27	8.03	:	8.07	8.66	:	:
10.04	9.84	<u>¥</u> 6	12	÷	:	÷	÷	÷	:	9-05	10-63	11.41	11.41	÷	:	÷	10-04	10-43	10-02	=	10-04	10-43	:	:
45.27	45.27	45‡	45å	45	45	45	45	443	44	44.49	44.49	44.09	44.09	4	4	4	43.7	43.7	43.6	43½	43.31	43.31	43	43
17.05	20.46	:	:	:	:	:	:	:	:	15.73	16.17	17.38	18.37	:	:	:	19.14	20.24	81	:	9.41	15-95	:	÷

- 14 - 8

- 20

- 11 - 11 8

- 20 - 12 - 15 - 22 - 14

- 20

- 14

- 20

12

- 14

21 21

- 18

		Prince Phil	Count Her
		•	
	Locality.	Germany	s Do
	Weight of Stag.	:	257 kilos
ontinued.	Spread.	:	368
EER-	Widest inside.	:	293
REDI	Tip to Tip.	:	471°
CONTINENTAL RED DEER—continues	Circum- ference between Bez and Trez.	:	:
	Circumfer- ence below Crown.		86
OTHER	Circumfer- ence above Middle Point.	7.48	7.t ₆ 9\$
	Circum- ference above Burr.	8.66	8 8
	Circum- ference of Burr.	10-04	† 6
	Length along Curve.	42.91	4218
	Number of Weight in lbs. Points. Avoirdupois.	18.92	:
	Number of Points.	- 16	- 20

9.0 7.28

6 6 9 7.68

8.27

9.84 9.64

42.91 42.91

90.91 20.06

91 -- 20 91 – 99.8

	Locality.
	Weight of Stag.
ontinued.	Spread.
DEER—con	Widest inside.
RED I	Tip to Tip.
CONTINENTAL	Circum- ference between Bez and Trez.
CONTE	Circumfer- ence below Crown.
OTHER	Circumfer- ence above Middle Point.
	Circum- ference above Burr.
	پ. پ

Owner.

	:
	,
	Weight of
red.	
ntin	
OTHER CONTINENTAL RED DEER—continued	Circum-
ED I	į
H	į
ENTA	Circum- ference
CONTIN	Circumfer-
OTHER	Circumfer- Circumfer-
	Circum-

- Philip of Saxe-Coburg Jotha. Ienry Fünfkirchen.

 - Prince Philip of Saxe-Coburg and Gotha. Count Jose Hoyos. Count Fery Nádasdy. Count B. Keglevich. Count Bombelles. Count Szechenyi. G. Jankovich.

Szilvás ņ.

ņ. å

265 kilos

:

2.68 7.87

7:09†

10-24 9.05

19:11

42.13 42.13

17.6

19.1

8.5

42.3 42.7

- 20 - 18 91 -11 - 20

- 14

. 8

9.64

18.85

94

42·I

18.8

18

:

Ö.

Munkacs

- Viscount Powerscourt (k).

Austria

- - E. N. Buxton.

Germany

311 35%

264

Galicia

Styria Gotha

ಜ್ಞ

- H.R.H. the Duke of Saxe-Coburg and Gotha. British Museum. Sir Clement Hill, Bart.
 - Viscount Powerscourt. Count Fery Nádasdy.

Germany Hungary

25

8.07

7:09

9.25

10.24

41.73

18.26

- 18

91

. 12

0

- 15

12 :

- Ö, <u>ي</u>
- ő.

Germany

Ď.

36

26

26

Prince Philip of Saxe-Coburg and Gotha,

Hungary .

:

Do.

230 kilos

Germany Hungary

32‡

275 :

> ₽6 8.27

13

햠

: : 100.7

10.63

12.21

40.04

21.01

8


- 14

Prince Nicolas Ghika.	Prince Philip of Saxe-Coburg	H.R.H. the Duke of Saxe-	Count Tassilo Festetics.	Viscount Powerscourt.	H.R.H. the Duke of Saxe-	Do.	A. von André.	Do.	Prince August Leopold of	H. J. Elwes.	Prince Philip of Saxe-Coburg	H.R.H. le Duc d'Orléans.	H.I.M. the German Emperor.
Roumania.	Hungary .	Tyrol .	Agarév .	Hungary .	Tyrol .	Gotha .	Hungary .	Do.	Do.	Macedonia	Hungary .	Tenuta la	Rominten .
:	:	÷	÷	:	:	:	:	:	:	:	÷	: -	st. 1bs. 24 6 (clean)
:	:	:	:	40	:	49	374	364	÷	÷	:	:	44 4
314	56	278	:	314	36	28	30‡	33	238	30	128	184	:
28¥	:	20 3	:	%	31	:	214	274	:	214	:	12	:
;	:	×	:	53	÷	÷	74	÷	:	5‡	:	44	:
n r S	7.5	÷	:	:	36	:	7.4	54	1 +8	;	7.5	:	:
:	7. 8	:	:	:	:	:	F 9	÷	÷	÷	:	፥	÷
:	1 6	:	8.10(?)	:	:	፥	† 6	88 88	:	:	78	:	: .
:	:	፧	:	:	01	01	01	₹6	13	:	108	:	÷
40 8	401g	40,	40-10	9	04	9	394	38₿	3718	37	36₿	314	%
:	50	:	18.50	:	:	:	:	:	26 }	÷	:	÷	194
- 11	- 14	- 13	- 20	18	- 12	- 18	15	00	- 14	- 14	1 20	01	44

NOTES AND EXPLANATIONS.

on this third tine, forming almost a cup. Mounted on a papier-	i. Tops palmated 6 in. across. j. Right trez tine forked, tops palmated. k. Killed in 1809. Bought at Linz, Austria, in 1898.	A remarkable specimen of a palmated head.o. Trez tine on right antler forked.	 p. From the Hildesheim Collection, near Hanover. f. The circumference of these antlers is taken above middle ints tine. 	Viscount Powerscourt informs me that the 18 stag heads in the entrance hall at Powerscourt, mounted on papier-mache heads, were bought at Munich in 1863, by the advice of the late Count Arco-Zinneberg, the owner of the famous collection in the Wittelsbacher Platz at Munich. The hall where they hang contains upwards of 2500 hads of German Red-deer and Roe-deer of extraordinary weight and size. forming with the one exception of the King of Saxony's collection at Moritabure, near Dresden.
 Many specimens marked thus are uneven pointers. a. Length of brow tine, 22³/₂ in. Bez, 23 in. 	b. " 21½ in. On papier-mache head. very large 8 pointer. c. Palmated tops 6 in. across. Length of brow tine, 17½ in.	d. Length of brow tine, 21 in. Left trez tine, 21 in. c. Shot by a peasant in a drive in Buckowina, in 1896.	f. Bought in Munich. g. Length of left trez tine, 25 in. Bought in Vienna in 1863. h. Right antler is forked at trez tine, and carries three points	Viscount Powerscourt informs me that the 18 stag heads in the entrance advice of the late Count Arco-Zinneberg, the owner of the famous collection ir heads of German Red-deer and Roe-deer of extraordinary weight and size form

were bought at Munich in 1863, by the e they hang contains upwards of 2500 collection at Moritzburg, near Dresden, advice of the late Count Arco-Zinneberg, the owner of the famous collection in the Wittelsbacher Platz at Munich. The hall whe heads of German Red-deer and Roe-deer of extraordinary weight and size, forming with the one exception of the King of Saxony's the finest in the world. Count Arco became so well known as a collector that every fine specimen was brought to him for purchase.



Antlers of Red Deer. From a specimen in the Castle at Moritzburg.

After Dr. A. B. Meyer.

i.—ANCIENT CONTINENTAL RED DEER.

Length (straight).	Circum- ference.	Tip to Tip.	Spread.	Points.	Locality.	Owner.
•••				33 + 29	?	H.M. the King of Saxony, Moritzburg.
1-48	14 of burr	•••	6 ft. 310	12+11	?	Do.
		•••	•••	25+12	}	Do.
•••			•••	28	?	H.R.H. the Duke of Saxe-Coburg and Gotha.
-46½	8% above trez	5 ft. 2‡	•••	22	Alpine Stag?	Count Erbach-Erbach.
•••	93	•••	•••	28	Do.	Do.
	•••		•••	22	Switzerland	Do.
(on curve) 394	54	16	22½ inside	9+9	?	Viscount Powerscourt.
39½	6 1	22	31½. 39 outside	6+6	Germany .	Do.

¹ Weight, 411 lbs.

j.—NEW ZEALAND RED DEER (introduced).

Average height at shoulder, 47 inches.

Length on out- side curve.	ference between bez and trez.	Tip to Tip.	Widest inside.	Spread.	Points.	Weight.	Local	ity.	Owner.
-42	5 1	•••		$37\frac{1}{2}$	9+7	•••	Otago	•	W. Allen.
-411	6	16	•••	$28\frac{1}{2}$	6+5		Do.		Do.
-41	6‡			311/2	6+5		Do.	•	J. S. Handyside.
-41	54			37	7+6		Do.		C. R. Westmacott.
-39	$5\frac{1}{2}$			31	6+6		Do.	•	H. McLean.
-38	5	•••		33	6+6		Do.	•	E. C. Studholm.
37½	54	194	28½		6+6	400 lb. esti- mated	}Do.	•	C. R. Westmacott.
-37	5		•••	34	6+5		Do.	•	W. Telford.
-36½	6 8	••	321/2	•••	9+9	•••	Wairarap	a	J. S. Handyside.
-36	44			331	5+5		Otago		W. Telford.
311	53	101	18	•••	6+7		North Isl	and	Rupert Wilkin.



Skull and Antlers of Caspian Red Deer shot in the Western Caucasus by St. George Littledale.

CASPIAN RED DEER or MARAL (Cervus elaphus maral).

In this variety of the red deer, which probably intergrades with the typical race in the western Carpathians, the height at the shoulder reaches to about $4\frac{1}{2}$ feet, and the build is stouter, the neck thicker, and the head longer and more pointed than in the true red deer. The reddish summer coat of immature animals is very generally marked with numerous yellowish spots; and the colour of the winter coat is dark slaty gray on the back, with the tail-patch of a very bright

yellow, and a large amount of black on the shoulders, thighs, and under parts. The large and massive antlers are generally less complex than those of the typical race, the number of points being seldom more than eight on a side, and frequently only six; while the bez tine, which may be wanting, is often shorter than the long and upwardly curved brow tine, and the fourth tine is generally more distinct from the crown. The average weight is given as about 40 stone.

The typical locality of this race of red deer is the Caspian provinces of Northern Persia, whence it extends into the Crimea, and probably Asia Minor, and so on into Transcaucasia, the Caucasus, probably Circassia, and the Galician Carpathians. The exact limits between the range of this and the typical race are not yet determined; and it is noteworthy that specimens from the Caucasus have shorter faces than those from Northern Persia, and thus approximate to the true red deer. In Asia the term "maral" is applied not only to this animal, but also to Bedford's Deer and the Altai Wapiti.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Spread.	Points.	Esti- mated Weight. st. lbs.	Locality.	Owner.
48½	ó	30	411	•••	5+6		Ichater Dagh, Crimea	H.R.H. the Duke of Saxe- Coburg and Gotha.
481	7‡	(single	antler)		12		Asia Minor .	Lord A. Hay, British Museum.
48	7‡						Caucasus .	O. O. T
478	5 1	38	•••	•••	6+5	•••	Do	Do.
47	7 1	36	374	•••	6+7	•••	Do	Do.
46 2	58	18	33		8+8		Do	Do. British Museum.
45 2	7 2	25 1	35₹		8+8		Do	St. George Littledale.
-45½	8 1	65 1	•••		14	•••	Do	Grand Duke Mikhaelo- vitch.
45½	7 1	32 3	40½	46	8+8		Do .	Prince Demidoff.
45 8	5 \$	42	358		6+7		Ak Dagh, Asia Minor	F. C. Selous.
45 1	7 3			•••	9+6		Asia Minor .	Sir Edmund G. Loder, Bart.
-45	8	36	42		11+10	53 4	Caucasus .	Prince Demidoff.
44 ⁸	6 <u>1</u>	334	35 1		7 + 5		Do. .	St. George Littledale.
44‡	6 7	(single	antler)		9		Asia Minor .	
431	6		40		8+6		Asia Minor .	Museum. M. Le C. Findlay.
43½	6 <u>3</u>	211	35		7+7		Caucasus .	St. George Littledale.
431	5 2	16	28		6+6	•••	Crimea	Earl of Dunmore.

D

CASPIAN RED DEER or MARAL (Cervus elaphus maral)—continued.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Spread.	Points.	Esti- mated Weight,	Locality.		Owner.
$-42\frac{1}{2}$	6 <u>1</u>	25		34	10+11		Crimea (?)		H. J. Elwes.
42½ -42	7	27	311	341	8+7		Caucasus		Prince Demidoff.
-42	6	42		50	10+9		Crimea (?)		H. J. Elwes.
411	5 2	i 5 ½	28 1		6+5		Caucasus		H.R. H. le Duc d'Orléans.
401	S∰	22½	32 1		6+6		Ak Dagh		H. O. Whittall.
411 401 401	5 2 5 2 5 2	28 1	341		8+7		۶		Duke of Bedford.
40	51	21]	27½		6+6		Ak Dagh		H. O. Whittall.
39\$	5½ 5¾	33	36́₹	•••	8+7		Do.	•	F. C. Selous.



Head of Caspian Red Deer shot in Asia Minor by F. C. Selous.

BARBARY RED DEER (Cervus elaphus barbarus).

For characters, see under heading of Cervus elaphus.

Length on out- side curve.	Circum- ference between bez and trez.	Tip to Tip.	Widest inside.	Spread.	Points.	Locality.		Owner.
38 2	58	•••		•••	6+5	North Africa		Sir Edmund G. Loder, Bart.
36 <u>1</u>	44	227	284		4+4	Do.	•	British Museum.
36 8	41/2	171	258		4+4	Do.		Do.

DUKE OF BEDFORD'S DEER (Cervus xanthopygus).

Apparently allied to the red deer, but the antlers probably with not more than seven points each, and the coat rather more wapiti-like. Tail comparatively short, and limbs relatively long, as is the face. summer the head and neck are dark slaty, as are the inner sides of the limbs, while the rest of the upper parts is bright foxy red (occasionally browner), with the tail-patch sometimes totally wanting, or rather indistinct, and little or no black on the under parts and inner surfaces of the thighs; in winter the upper parts brownish gray with a very large and conspicuous bright orange tail-patch and a blackish mane, the tips of the hairs showing a large amount of black.

This deer inhabits Manchuria, and probably some of the other districts of North-Eastern Asia, but the western limits of its range are still undetermined. It is regarded by Monsieur E. de Pousargues as inseparable from the Manchurian wapiti; but this is not borne out by the specimens now living in England. If they survive, the question can be decided in a year or two.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.		Owner.
-20 <u>1</u>	48	141	16 <u>1</u>	4+3	Manchuria		Paris Museum (Type Specimen).



Head of Hangul. From a specimen shot by J. G. Apcar in Kashmir.

HANGUL or KASHMIR STAG (Cervus cashmirianus).

In this very distinct species the first or brow tine arises at a considerable distance above the burr, or coronet, of the antlers, instead of close to it, as in the red deer; the bez tine is usually longer than the brow; the total number of points is generally only five aside, although a third tine may occasionally be added to the normal terminal pair, thus forming an imperfect cup; and the beam of each antler is much curved in towards the middle line of the head. The tail is short, and not included in the light patch on the buttocks, which, at least frequently, is very small; and the tuft on the hind cannon-bone is situated lower down than in the red deer. In winter the general colour of the coat is brown, brownish ash, or liver-colour, with the hairs speckled; the light area on the inner side of the buttocks

being dirty white, with a blackish line on the inner sides of the thighs, the upper side of the tail black, and the lips, chin, and inner surface of the ears white or whitish. In the fawns the spotting is stated to remain much longer than in the red deer. In the pairing season the old stags squeal like a wapiti, instead of roaring in the red deer fashion.

The typical hangul inhabits the forest districts of the north side of the vale of Kashmir and some of the neighbouring valleys, at elevations ranging from 9000 to 12,000 feet in summer, but descending to about 5000 feet in winter. In this race (C. cashmirianus typicus) the terminal or fifth tines of the antlers are so much bent inwards as to be separated by a comparatively small interval. In the forests of the Yarkand river the species is represented by a second local race (C. cashmirianus yarcandensis), in which the antlers are less spreading, with their terminal tines less inclined inwards, and therefore more widely separated. The trez tine in this race is typically larger and longer than either of the lower ones, the brow and bez, in this respect and in the closer proximity of the brow and bez tines more resembling the Caspian than the Kashmir Stag. The height at the shoulder varies from about 4 feet to 4 feet 4 inches; the average weight being about 450 lbs.

This deer, although first discovered by the late Dr. H. Falconer in the Kashmir valley, was named by Dr. G. R. Gray of the British Museum. The finest pair of antlers of which Mr. A. O. Hume has any record were given by Raja Gulab Sing to Colonel King, then commanding the 14th Dragoons. On his death they passed to Captain, afterwards, I think, Colonel, Prettyjohn of the same regiment. What became of these antlers Mr. Hume was never able to ascertain, but he measured them at Meerut in 1852 or 1853, and the record stands, R. 52 L. 53½, measured along the curve inside. Girth 10 inches at burr, and 7 half-way between bez and trez tines. They were a very wide-branching, symmetrical pair.

Length on out- side curve.	Circum- ference between bez and trez.	Tip to Tip.	Widest inside.	Points.	Lo	cali	ty.		Owner.
-48	•••			•••	Kashmir				The late Dr. Leith Adams.
-47	7 8	21	36	7+5	Do.		•	-	Bombay Natural History Society's Museum.
47	6 3	21‡	36 3	5+5	Do.				Sir Edmund G. Loder, Bart.
47	6 <u>‡</u>	30	35½	8 + 8	Do.				Duke of Wellington.
-47		•••				?			Major A. E. Ward.
45 8	8	35	41	6 +6	Kashmir .				Hume Collection, British Museum.
451	6	254	36	8 + 8	Do.				Sir Victor Brooke's Collection.
45	63	19	34	6+6	Do.				Col. R. Pole-Carew, C.B.

HANGUL or KASHMIR STAG (Cervus cashmirianus)—continued.

Length on out- side curve.	Circum- ference between bez and trez.	Tip to Tip.	Widest inside.	Points.	Locality.	Owner.
-44 ⁸	6	20	43	5+5	Lidar Valley	Officers' Mess, Q.O. Corps of Guides.
-44§	61/2	3112	44¥	5 + 5	Kishenganga Valley	Do.
$-44\frac{1}{2}$	6	211	36 1	5+5	Do	The late Capt. E. W. Codrington.
44	7‡	30 3	40 7	5+5	Sind Valley	P. H. G. Powell-Cotton.
44	6 <u>₹</u>	231	36 1	5 + 5	Do	Naval and Military Club.
44	6	27	3 6	5 × 5	Kashmir	Hon. Walter Rothschild.
438	5 7	157	32	5 + 5	Do	Hon. Charles Ellis.
438	6 <u>1</u>	258	36½	•••	Do	Do.
43	6	20	35	6+5	Sind Valley	A. O. Hume, C.B.
43	5å	26 1	37 1	6+5	Kashmir	Martyn Kennard.
42	$4\frac{1}{2}$	134	29 1	5+5	?	Duke of Bedford.
42 .	5₹	13½	26½	5+5	Kashmir	E. L. Phelps.
413	54	23½	35	5+5	Do	Major Kingsley Foster.
418	5 8	232	33	5+5	Do	R. Lydekker, British Museum.
411	6	158	29	7+6	Tral Preserve	P. W. Cobbold.
414	5 1	34	49	6+5	?	C. H. Seely.
-41				13	Tral Preserve	The Maharaja of Travancore,
-41			•••	12	Do. }	G.C.S.I.
-4 I	6	23	37½		Kashmir	The late Major W. D. B. Fenton.
40 1	58	317	39½	6+5	Do	F. W. H. Walshe.
40½	6 <u>‡</u>	151	28	5+5	Do ,	Sir Robert Harvey, Bart.
40 <u>‡</u>	7∰	24½	3	5+5	Do	Capt. H. W. Codrington.
39½	53	2 I ½	33	5+5	Do	H. Maude.
39	54	26	351	5+5	?	Major A. Nugent.
381/2	5	141	29	5+5	Rewa Nalla	The late Major A. Burton.
37½	5	17	28 <u>1</u>	5+5	Kashmir	Reginald Beech.
371	5₩	26 2	34½	6+6	Do	Major C. S. Cumberland.
37	5 7	22 8	314	6+6	Do	W. R. Bindloss.
37	5	20	29	5+5	Do	H. Z. Darrah.
37	54	19½	311	5+5	Do	J. G. Apcar (see Illustration).



Skull and Antlers of Yarkand Stag. From A. O. Hume's specimen.

YARKAND STAG (Cervus cashmirianus yarcandensis).

For characters, see Hangul, p. 37.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Spread.	Points.	Locality	y.		Owner.	
40 1	51/2	234	313	•••	5+5	Near Maralb	ashi		A. O. Hume, C.B. illustration.)	(See
40	5 1	25	31	•••	6+5	Maralbashi		•		
391	6	$24\frac{1}{2}$	25		7+6	Do.			A. O. Hume, C.B.	
394	5₽	16 1	28 3		6+6	Do.			David T. Hanbury.	



Skull and Antlers of Shou. From A. O. Hume's specimen.

THE SHOU (Cervus affinis).

A very large stag with antlers of the general type of those of the hangul, but larger, and with the beam bent suddenly forward at the trez tine, so that the upper half overhangs the face, the number of points being usually five; the brow tine is less constantly longer than the bez. General colour probably rufous brown, with a light tail-patch in winter.

The habitat of this imperfectly known deer appears to be the districts immediately north of Bhutan, and probably the valley eastwards of Chumbi, which drains northwards into the Sangpo. An apparently allied, although smaller, deer is found in Russian Turkestan (Bokhara or Khiva).

Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.	Owner.
55\$	6 1	17‡	40 8	7+6	?	A. O. Hume, C.B. (See illustration.)
548	6 §	21 5	37 1	5+5	?	Dr. Campbell, British Museum.
55‡	6 7 (see ill	26 1 ustratio	44 n, p. 55)	5 + 5	?	The late B. H. Hodgson, British Museum.
534	6 7	30	453	4 ÷ 5	?	Hume Collection, British Museum.
-53	9		40	5+5	?	The late Col. H. C. B. Tanner.
52	8			•••	?	Hon. Walter Rothschild.
49½	71	38	45½	5 + 5	?	Col. J. Biddulph.
48 3	6 <u>1</u>	19 3	38 1	7+5	?	Sir Edmund G. Loder, Bart.
-48	63	303	¹ 39	5+5	Tibetan Frontier	H. J. Elwes.
-48	6 8	17	34½	6+5	?	H.R.H. le Duc d'Orléans.
47 ⁷	5 å	30 3	40]	5+5	?	British Museum.
-43		•••	•••		Chumbi Valley	The late Sir Ashley Eden,
42‡	5 7	235	311	5+5	?	G.C.S.I., Indian Museum. British Museum.
41	6 1			6+4	?	Duke of Bedford.
39 3	5	20	311	5 + 5	?	Edward P. Tennant.
					1 6	

1 Spread.

THOROLD'S DEER (Cervus albirostris).

Thorold's deer is of the same approximate dimensions as the hangul, from which it is readily distinguished by the more flattened antlers, which have no bez tine, and do not curve inwards, but are suddenly bent backwards at the point of origin of the trez; the total number of points being either five or four. Equally distinctive is the pure white muzzle and chin, the white inner surface of the ears, the reversal of the hair on the middle of the back, so as to form a kind of hump on the withers with the points of the hairs directed towards the neck, the low position and large size of the gland-tuft on the hind cannon-bone, and the shortness of the tail, which is included in the very large straw-coloured area of the buttocks. The general colour of the coat is uniformly dark brown, with the hairs, which are remarkable for their coarse and brittle nature, minutely speckled.

The Tibetan plateau, with perhaps some of the neighbouring parts

of Central Asia, is the home of this fine species of deer, which was originally described by the late Colonel Przewalski under the name given above. Subsequently two examples were obtained by Dr. W. G. Thorold, to the north-east of Lhasa, at an elevation of between 13,000 and 14,000 feet, which, under the impression that they indicated a new species, were named *C. thoroldi* by Mr. W. T. Blanford.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.		Owner.
38 1	4	37	27	5+4	Central Tibet .	•	Hon. Walter Rothschild.
38	5‡	•••		5+5	Do		British Museum.
-36	41/2	32	32	5+5	? 1		Indian Museum.

¹ Bought at Darjiling, thither brought by Tibetans.

TRUE WAPITI (Cervus canadensis).

Wapiti are very large deer of the red deer group, easy of recognition by the form of their antlers, which are of great size, carrying more than five tines, curving backwards, and much flattened in the upper half. They always have the bez tine developed, but their most characteristic feature is the great size of the fourth tine, which is larger than either of the others, and with the fifth, which is also long, forms a nearly straight fork; the fourth, fifth, and sixth tines being situated almost in the plane of the portion of the beam immediately below them, so that they more or less completely hide one another when viewed from the front aspect. The brow tine rises close to the burr, and is nearly as long as the bez. The tail is extremely short, the light tail-patch very large; and the neck and under parts are blackish, the general colour of the summer coat being yellowish brown on the upper-parts.

Wapiti (known in America as Elk) range from North America to North-Eastern and Central Asia; the typical form being the East American wapiti (C. canadensis typicus), in which the legs are comparatively short, and the portion of the antlers above the fourth tine is fully developed, the height at the shoulder reaching to about 5 feet 4 inches, and the weight from 700 to 1000 lbs. On the other hand, the West American wapiti (C. canadensis occidentalis) differs by the abortion of the upper part of the antlers, the darker colour, and lighter build.



Head of East American or true Wapiti.
Shot by W. Moncreiffe. For measurements see next page.

TRUE WAPITI (Cervus canadensis)—continued.

					-			
Length on outside curve.	Circum- ference between bez and trez.	Circum- ference of burr.	Tip to Tip.		Widest outside.	Points.	Locality.	Owner.
-70		I41 above burr.		•••	68	6+6	Olympic Mts., Washington	W. F. Sheard.
-66		•••			60	6+6	Wyoming .	J. Darley.
-65	7 1	•••			40	7+7	Laramie Plains, Wyoming	Schoverling, Daly, and Gales.
-643		9₹	411	49	•••	6+7	Wyoming .	James J. Harrison.
641	8		311	45	52	7+6	?	Viscount Powers- court.
-64‡	8	•••		48		7+7	N. W. Wyoming	A. Rogers.
-634	8 1		491	48 <u>1</u>		7+9	North Prong .	Frank Cooper.
62	78	•••	33 1	50 §		7+7	Snake River, Colorado	Ernest Farquhar.
61 3	71/2	•••	•••	45		6+6	Bighorn Mts., Wyoming	., ,,
-6 1	8			52		6+ 6	Do	II. Seton-Karr.
61	8		36 <u>4</u>	46½	55	8 × 8	?	Viscount Powers- court.
¹-603	7 2			52		6+6	?	W. A. Baillie Groh- man.
60½	8‡			55		7+6	Wyoming .	E. Grant.
60§	71		43	46 1	•	6+6	Do	Major C. C. Ellis.
60	6 <u>3</u>		31 1	411	44	9+7	?	Viscount Powers- court.
-593		13	•••		•		Wyoming .	The late Sir Samuel Baker.
59½	81	3	37‡	47	•••	8+8	S.E. Do	Lieut General B. Hankey.
59½	7½			¹ 45		6+6	?	II. Seton-Karr.
58 1	83	•••	39‡	$43\frac{1}{2}$	53½	7+6	?	Viscount Powers- court.
$58\frac{1}{2}$		9 between brow and bez.		46 <u>1</u>	50½	10+7	Wyoming .	Hon. T. A. Brassey.
$-58\frac{1}{2}$	81		•••	44 ½		6+6	Ю	J. D. Cobbold.
-58½	8 1		48		50	6+6	?	F. B. Tolhurst.
58	914	42	47‡	7+6		7+6	Wyoming .	A. II. Straker.
58	78		43½	49		9+8	Do	W. Moncreiffe. (See illustration.)
58	7		46	49	511	6+6	?	Viscount Powers- court.
57½	6 <u>7</u>		24 1	35	•••	6+6	Montana	Capt. Abdy.
57₿	7		47	48§		6+6	Wyoming .	Hon. Charles Ellis.
571		9∄	32	42‡		7 + 7	Do	Sir Humphrey de Trafford, Bart.
			1 M	easured l	by Americ	an Exhibit	tion Committee.	rimoru, Datt.

TRUE WAPITI (Cervus canadensis)—continued.

Length on outside curve.	hattuaan	Circum- ference of burr.	Tip to Tip.	Widest inside.	Widest outside.	Points.	Locality.	Owner.	
57	8 <u>1</u>		$35\frac{1}{2}$	34	43	6+6	Wyoming .	Viscount Powers- court.	
-57	7 1		•••		•••	7 + 7	Do	Count F. Trautt- mansdorff.	
57	7 1		53	$49\frac{1}{2}$	61	8+9	Wyoming .	Sir Edmund G. Loder, Bart.	
57	7 1	•••	35 lb	41 ½	•••	6+6	Yellowstone Park	British Museum.	
-56½	6 7		•••	•••	46 ğ	6+6	Two Ocean Pass	Theodore Roose-	
56	8		394	40 1	44	8+6	?	Viscount Powers-	
-56	7 3					6+6	Wyoming .	Count F. Trautt- mansdorff.	
56	7‡		35	$43\frac{1}{2}$		8+7	Saskatchewan Dis- trict, Canada	Earl of Ava.	
558	7 8		43 1	481		7+7	Wyoming .	Hon. Charles Ellis.	
55₿	7		43½	478	•••	6+5	Saskatchewan Dis- trict, Canada	Capt. G. Dalrymple White.	
-551	8		38	43	••	6+6	Wyoming .	James J. Harrison.	
-55 <u>₹</u>	7 ½		•••	483		8+7	?	Otho Shaw.	
55 1	6 1		44	45⅓		6+6	?	A. H. Pollen.	
55 8	7 l		46 ‡			7+6	?	St. George Little-	
55	8‡	•••		41}	•••	6+6	Wyoming .	Major Maitland Kirwan.	
55	8			45½		7+6	Do	Do.	
55	7 l	•••		47 ½		6+5	Do	E. N. Buxton.	
-55	81		53	54½		17	?	Sir H. B. Meux, Bart.	
-55	8 1			411		6+6	3	A. Pendarves Vivian.	
¹ 55	7 3		21½	35½	43	7+6	?	Viscount Powers- court.	
² 55	6₽					14	America	Duke of Bedford.	
55	6 1 ̄	•••			•	6+6	Colorado	Crawford G. Logan.	
-543	63		39	$43\frac{1}{2}$	•••	6+6	Montana	Count Scheibler.	
54½	8		37	$45\frac{1}{2}$	•••	6+6	Wyoming .	Sylvester Browne.	
54½	• • • • • • • • • • • • • • • • • • • •			•••	•••	14	Colorado	Sir Peter Walker, Bart.	
54½	8		34‡	45½	•••	6+5	Wyoming .	LieutCol. Hon. W. Coke.	
54½	9	•••		palm	nated	11+7	Montana (?) .	G. Wrey.	
54½	6 3		28	37	46	6+6	?	F. J. Mitchell.	
1 Shot by the Earl of Dunmore.							² Shed antlers weighed 20 lbs.		

¹ Shot by the Earl of Dunmore.

² Shed antlers weighed 20 lbs.

TRUE WAPITI (Cervus canadensis)—continued.

Length on outside curve.	Circum- ference between bez and trez.	Circum- ference of burr.	Tip to Tip.	Widest inside.	Widest outside.	Points.	Locality.	Owner.
548	61/2	•••	38 8	44 1	•••	7+6	Wyoming .	E. N. Buxton.
-541	8			43½	•••	10+10	Montana	W. A. Tulloch.
54 1	7 §		498			9+7	Wyoming .	Hon. Walter Roths- child.
-54	81		•••	•••		13	Teton Mountains	H. Lennard.
¹ 54	71			48		8+7	Wyoming .	Moreton Frewen.
54	71/2		32 8	39½	47	11+8	Do	Viscount Powers- court.
54	78	•••	30 2	37 8		6+6	Do	Capt. E. G. Ver- schoyle.
54	74	•••	•••	48		7+8	Do	Moreton Frewen.
54	7		29 <u>1</u>	351	44	7+6	Do	H. Seton-Karr.
54	7 2		48	44		12	?	M. P. Grace.
-53½	8	•••	43	441	•••	8+7	Wyoming .	Count E. Hoyos.
-53½	61		39	42		6+6	Do	Captain G. Dal- rymple White.
53½	7		411	42	•••	7+7	Colorado	E. T. Logan.
53 1	78	•••	47	49 3	•••	8+7 palmated	Wyoming .	Ford G. Barclay.
-53	•••	121/2		•••		···	Do	The late Sir Samuel Baker.
53	8 <u>‡</u>	•••	29½	36	38	7+6	Do	Duke of West- minster.
53	71	•••	41	43½	•••	6+6	Do	Major H. J. Ferguson.
53	71	13	37 1	411	51	7+7	3	Hon. Walter Roth- schild.
-52 3	71	•••	38 1	41#	•••	6+5	Wyoming .	Charles Makin.
52 1	7		32	43	49	7+6	Do	J. B. Gilliat.
52½	$6\frac{1}{2}$		•••	34 2	•••	7+6	Do	Ford G. Barclay.
52½	61		39	42	•••	6+6	Do	Captain G. Dal- rymple White.
52 8	54	•••	49 8	•••	•••	6+5	Colorado	Colonel Ralph Vivian.
52	81	•••	•••	•••	•••	6+6	Wyoming .	Hon. F. Thellusson.
52	6 3	•••	29 1	36 1	•••	6+5	Colorado	T. C. E. Goff.
-52	71/2		46 <u>1</u>	•••	•••	6+6	Do	Dublin Museum.
52	71/2	•••	30	33	•••	6+6	Wyoming .	Captain G. Dal- rymple White.
52	8		48 <u>1</u>	49½ 1 S	 hot by Si	8+7 ir G. Gore, I	Do	Major H. C. Mor- land.
				_				

TRUE WAPITI (Cervus canadensis)—continued.

Length on outside curve.	Circum- ference between bez and trez.	Circum- ference of burr.			Widest outside.	Points.	Locality.		Owner.
52	8	•••	381	44	•••	8+8	Wyoming		Lord Rodney.
52	81	•••	37	40	•••	6+6	Vancouver 1		Barclay Bonthron.
52	6 <u>1</u>	••	36	36 1	41	6+4	Manitoba .	•	Earl of Dunmore.
-52	81		•••	42	•••	14	Do		Prince Henry of Liechtenstein.
-52	7		30½	39		6+5	Washington		P. B. Vander-Byl.
51 1	71		42	46 <u>1</u>		6+6	Territory Colorado .	•	Captain E. G. Vers-
511	$6\frac{1}{2}$		41	441	•••	7+7	Idaho .		choyle. H. C. Nelson.
-518	81				50		Wyoming		G. O. Shields.
51	7‡		30	35		7+7	Do		J. Turner-Turner.
51	61		39	45	•••	6+6	Do		Captain E. G. Vers- choyle.
-51	83		•••	41		7+6	?		Captain Joscelin Bagot.
50 §	7		42	48 1		7+7	Colorado .		Colonel Ralph Vivian.
50 <u>1</u>	7		36	47	•••	7+6	Wyoming	•	Major C. F. Blane.
-50 1	71/2		•••	56 1	58 1	7+6	Do		Hugh Peel.
50 <u>1</u>	64		50	43	•••	6+6	Do	•	Major Maitland Kirwan.
501	7		431	43½	•••	6+5	Do	•	W. Moncreiffe.
50}	61	•••	411	47½		7+6	Do	•	Licutenant - Colonel Hon. W. Coke.
50½	71/2	•••	35	41	•••	6+6	Do		Hon. H. S. Somer-
50	•••		49		•••	8+7	Do		Otho Shaw.
50	6 <u>1</u>	•••	44	44 1	47	6+6	Do		V. Cholmondeley.
-50		11	•••	•••	•••	13	Montana .		T. W. H. Clarke.
50	71/2		391	40		6+7	Wyoming		Prince Demidoff.
50	7 1 8			471		6+6	?		J. M. Hanbury.
50	81		39	40		6+6	3		Duke of Bedford.
50	78		43	40 <u>3</u>		6+6	?		Duke of Portland.
50	7		30]	39		6+5	Washington	•	P. B. Vander-Byl.
50	61		40 <u>1</u>	362	40	7+6	Manitoba	•	Major C. S. Cum-
-49 8	62	••		331	45	6+6	?		berland. H.R.H. the Duke of Saxe-Coburg and Gotha.

1 West American race.

 $\begin{tabular}{lllll} \textbf{TRUE WAPITI (Cervus canadensis)} &-- continued. \\ \end{tabular}$

Length on outside curve.	Circum- ference between bez and trez.	Circum- ference of burr.	Tip to Tip.	Widest inside.	Widest outside.	Points.	Locality.		Owner.
-49½	7활	•••	45‡	45 1		7+7	?		H.R.H. le Duc d'Orléans.
49½	8		•••	43		11+8	Wyoming		Otho Shaw.
$49\frac{1}{2}$	6‡		34½	374	•••	6+6	Do		J. Kenneth Foster.
498	7 ⁵	•••	38 7		•••	15+10	Do		Moreton Frewen.
-49 1	•••	97	28	36 1	•••	7+7	Canada .		Paris Museum.
49 1	84	• • •	381	4112	•••	6+6	Montana .		L. B. Lee.
-49 1	6 <u>1</u>	•••	38	46 1		10+7	Vancouver 1		A. E. Leatham.
49 1	6 1			45		13	Montana .		W. A. Fordham.
49	$7\frac{1}{2}$		26 [‡]	34	41	7+6	Wyoming		J. L. Scarlett.
49	$6\frac{1}{2}$		$48\frac{1}{2}$	51	•••	8+7	Do		Hon. Gathorne Hardy.
49	7 3		33	41		8+5	Do		H. Seton-Karr.
-48 3	•••	11	27 5	32‡		7+7	Canada .		Paris Museum.
-48	•••	I I ½	•••		•••	17	Montana .		T. W. H. Clarke.
48	$7\frac{1}{2}$		41	40 <u>3</u>		8+7	3		Duke of Bedford.
48	6 ‡	•••	41	43		7+7	Wyoming		F. C. Selous.
48	6	•••	40½	39	•••	5+6	Do		Captain G. J. Fitz- gerald.
47 1	5≇		48	42		8+6	Colorado.		E. T. Logan.
47 1	7‡	•••	40	41	•••	7+7	Wyoming		Sir Victor Brooke's Collection.
47	$6\frac{1}{2}$		40	42	•••	6+6	Do		W. W. Ashley.
-45	71	•••	37	37		7+8	Vancouver 1	•	Clive Phillipps- Wolley.

¹ West American race.

ALTAI WAPITI (Cervus canadensis asiaticus).

From the typical wapiti this well-marked local variety differs by its inferior size, relatively longer body and shorter limbs, and absolutely larger antlers; the general colour of the coat being yellowish tawny at all times of year. This wapiti inhabits the forest-clad portions of the Altai and Thian-Shan ranges; it was first described by Dr. Severtzoff under the name of *C. maral*, var. asiatica, and subsequently by Mr. Blanford, on the evidence of detached antlers obtained by the Second Yarkand Mission, as *C. eustephanus*.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Spread.	Points.	Locality.	Owner.
¹ -55	8		•••		6+7	Bought at Kashgar	Earl of Northbrook.
-51	10,9		•••		•••	?	W. T. Blanford.
50 2	6	29½	381	•••	8+7	Altai	Duke of Bedford.
-49 §	8	41	42 l		7 + 7	East Thian Shan	Paris Museum.
481	6 1	36	33		6+6	Altai	Duke of Bedford.
45½	6 <u>‡</u>	344	39½	47	5 + 5	Do	H. J. Elwes.
433	57	201	35	•••	6+5	?	A. O. Hume, C.P.
417	6 <u>1</u>	$37\frac{1}{2}$	40½	44	5+5	Altai	H. J. Elwes.
39	48	32	36	37 ½	6+6	Thian Shan .	H. J. Elwes.

¹ Shed antlers bought by Sir Douglas Forsyth.

MANCHURIAN WAPITI (Cervus canadensis luehdorfi).

Antlers of a much shorter and stouter type than in the Altai wapiti, with the portion above the fourth tine (which is also relatively smaller) generally only slightly developed, and sometimes aborted. Not unfrequently "sports" in the neighbourhood of the fourth and fifth tines; and in one instance the fourth tine itself split into a regular fork. Build and colour very similar to that of the typical wapiti, but the height apparently lower. General colour in winter brownish gray, in summer light brown, with the dark winter mane and under parts of the true wapiti.

Although the antlers of this race (known in this country by the under-mentioned and other specimens in the collection of the Duke of Bedford) are at first sight very unlike those of either the Altai or the true wapiti, yet they present the essential wapiti characters. These are shown in their flatness, the preponderating size of the fourth tine, and the position of the fifth tine in the same plane as the latter. They are much more like the antlers of the West American wapiti (a fine pair of which are exhibited in the British Museum); these being relatively short and stout, with a tendency to the abortion of all the tines above the fifth, and also to the production of "sports."

As mentioned above, this deer is identified by Monsieur E. de Pousargues with C. xanthopygus; but there is no evidence that the coat is red in summer, and the antlers of immature specimens of the latter now living in England do not display decided wapiti characters. Moreover, these animals roar somewhat like a red deer, instead of squealing or whistling like a wapiti.

Distribution.—Northern Manchuria and Amurland.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Spread.	Points.	Locality.	Owner.
33 1	54	18 1	$21\frac{1}{2}$	·	6+6	?	H. J. Elwes.
32 1	5 1	26 7	26 3	29 7	7+6	?	Do.
313	6	17	25%	29	8+7	?	Duke of Bedford.
3112	5	231	24	26	6+6	?	Do.
31	41	26½	257	30½	6+5	?	Do.



Antlers of Japanese Sika.

From a specimen presented to the British Museum by Viscount Powerscourt.

JAPANESE SIKA (Cervus sica).

This species is the typical representative of a small group of deer in which the antlers are shorter and simpler than is usually the case in the red deer group, and have generally four tines, including a trez, but lacking a bez. The coat is spotted, at least in summer, and there is a black-bordered white area in the region of the tail, which is relatively long. In the Japanese sika the white area on the buttocks is large, and extends on to their lateral surfaces; while the coat is chestnut red with numerous white spots in summer, and browner, with no, or only indistinct traces of, spots in winter. These deer are distributed over Northern China, Manchuria, and Japan, and are represented by two closely allied races differing chiefly in size. In the true Japanese sika (C. sica typicus), which inhabits Japan and Northern China, the height at the shoulder varies from about 2 feet 8 inches to 2 feet 10 inches, whereas in the Manchurian sika (C. sica manchuricus) it reaches

3 feet 3 inches. Both races have been acclimatised in English and Irish parks.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Points.	Locality.			Owner.				
311	. 5 1	27 1	4+4	Japan			Sir Edmund G. Loder, Bart.				
25	4, <u>l</u>	201	4+4	Do			Do.				
22	4	16∄	5+4	Bred in Ireland .			Sir Victor Brooke's Collection.				
21 5	38	16 <u>1</u>	10+6	Island of Yezzo.			Do.				
181	3 2	•••	9	Bred in Ireland .			Do.				
$-18\frac{1}{2}$	3 1 8	11	4+4	Kobe, Japan .	•	•	Dr. Percy Rendall.				
16 <u>8</u>	318	•••	4+3	Japan	•		British Museum.				
16 8	3	98	4+3	Bred in Ireland.			Hon. John Ward.				
151	2 3	9 1	4+4	Bred in England			Duke of Bedford.				
1 152	2 3	144	4+4	Bred in Ireland.			Marquis of Hamilton.				
2-15½	•••	•••	6+5	Do	•		Viscount Powerscourt.				
154	38	121	4+4	Do			Hon. R. A. Ward.				

¹ Weight 10 stones 3 lbs. as it fell.

This stag when killed was estimated to be fifteen or sixteen years old, and had no teeth left.

The following specimens belong to the Manchurian race.

Length on out- side curve.	Circum- ference.	Tip to Points.			Locality.					Owner.		
-26	41	138	4+4	Manchuria					•	Paris Museum.		
151	3		4 ÷ 4	Woburn .						Duke of Bedford.		

HYBRID JAPANESE and RED DEER.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Weight.	Locality.	Owner.
2 9‡	$4\frac{1}{2}$	172	21	5+4	14 st.	?	Viscount Powerscourt.
281	41	141	18	5+4	14 st. clean	?	Do.
26½	4	$21\frac{1}{2}$	21	4+4	···	?	Do.

FORMOSAN SIKA (Cervus taëvanus).

Nearly allied to the Japanese and Manchurian sikas, but distinctly spotted in winter, when the coat retains more or less of the rufous summer tinge. The dark line down the middle of the back is very strongly marked, there is a more distinct black bar above the white tail-patch, and the limbs are shorter, and the body proportionately longer. The height at the shoulder is about 2 feet 11 inches. This species is confined to the mountains of the island of Formosa.

Length. ference. Tip.		Tip to Tip.	Widest inside.	Points.	Locality.	Owner.		
193	3	13	163	4+4	Island of Formosa		British Museum.	
19	31			4+4	Do.		Duke of Bedford.	



Head of Pekin Sika, from a specimen in the Museum at Woburn Abbey.

PEKIN or DYBOWSKI'S SIKA (Cervus hortulorum).

In addition to its larger size (at least 3 feet 7 inches at the shoulder), this species is distinguished from the Manchurian sika by the smaller size of the white tail-patch, which in fully adult individuals does not extend on to the sides of the buttocks, although it does so in younger animals. The head and neck are bluish gray, and in immature animals spots persist in the winter coat, although, except on the hind-quarters, they may disappear more or less completely at this season in fully adult bucks, whose coats become very long and shaggy, especially on the throat and neck. Hinds are more brightly coloured in winter than the stags, and retain more distinct spotting. This deer was first named by the late Consul Swinhoe from an immature buck and doe taken at the sack of the Summer Palace, Pekin, and was afterwards obtained in

the wild state in the Ussuri district of North-Eastern Manchuria, when it received the name of *C. dybowskii*.

Length on out- side curve.	Circum- ference.		Widest inside.	Points.	Locality.	Owner.
27	41	23½	20	4+4	Manchuria	Hon. Walter Rothschild.
26	34	18	151	4+4	Do	Edward P. Tennant.
24	4	$22\frac{1}{2}$	19	4+4	Do	Hon. Walter Rothschild.
23½	$3\frac{1}{2}$	18‡	16 <u>1</u>	4+4	Do	Duke of Bedford.
1 5 g	3	20	•••	4+4	Ussuri, South Man- churia	British Museum.



Skull and Antlers of Shou. From a specimen in the British Museum.



Head of Common Fallow Deer.

FALLOW DEER (Cervus dama).

Antlers normally without a bez, but with a trez tine, above which the beam is palmated, with numerous snags on the hinder edge. Coat spotted with white in summer (except in the black breed), with a black-bordered white area in the neighbourhood of the long tail. Height at shoulder about 3 feet; weight about 140 lbs. clean. The original distribution includes Greece, Spain, Portugal, Anatolia, Rhodes, Sardinia, Asia Minor, Northern Palestine, and North-Western Africa, but the species has been introduced into Great Britain and some other countries.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Spread inside.	Points.	Width of Palm.	Locality.	Owner.
1-31	5	•••		30	7	?	Sir Victor Brooke's Collec- tion.
² -30	44	23½	26 <u>1</u>	10+9	41/2	Drummond Castle, Perth	

¹ Recorded by J. G. Millais (British Deer and their Horns).

² Weight of antlers, 8 lbs. 1 oz. on skull, no lower jaw (Millais, British Deer).

FALLOW DEER (Cervus dama)—continued.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Spread inside.	Points.	Width of Palm.	Locality.	Owner.
- 30	$4\frac{1}{2}$	22	37 outside	16 + 10	7	Drummond Castle, Perth	J. G. Millais.
-29½	5	17	28½	14+13	71	Petworth Park, Sussex	Do.
1-281	4	141	26	10+11	6	Woburn	Duke of Bedford.
2 28 $^{\frac{1}{2}}$	4	(Spar	n 34) side	19	6	Colebrooke	Sir Victor Brooke's Collec- tion.
³ 28	4	(26)		18	5	Do	Do.
4-27\$	5	20½	$21\frac{1}{2}$	13+7	5#	Do.	A. Basil Brooke.
27 3	41	32½	25	11+10	•••	Woburn	Duke of Bedford.
271	4	23		10+8	•••	England	J. Carr Saunders.
-27					•••	Nr. Blair Castle .	Dowager Duchess of Atholl.
26 7	34	12	17½	10+7		?	British Museum.
-26 1	41	$23\frac{1}{2}$	20½	9+7	57	Perthshire	A. Basil Brooke.
-26	5	20 1	234	10+10	$4\frac{1}{2}$	Tasmania	T. W. H. Clarke.
-26	5‡	20	22½	11+13	7	Ashton Park, Lan- cashire	J. Whitaker.
25 3	41	26		8+8	5À	Ireland	Sir Victor Brooke's Collec- tion.
-25	4	24 ³	25 1	11+10	5≹	Tasmania	T. W. H. Clarke.
-25	41	211	24	11+10	7	England	Sir Edmund G. Loder, Bart.
$23\frac{1}{2}$	3 ‡	14	191	10+9	41/2	?	G. O. M. Herron.
-23	41/2	15	$18\frac{1}{2}$	10+9	5	?	Dublin Museum.
21 8	48			11+10	5	Colebrooke, Ireland	Sir Victor Brooke's Collection.

¹ Weight dressed, 165 lbs.

Seven years old, as he fell 237½ lbs., 198 lbs. cleaned.
 (Millais, British Deer).
 Seven years old. 3 Seven years old, as he fell 224 lbs. (Millais, British Deer).




Head of Mesopotamian Fallow Deer, from a specimen in the British Museum.

MESOPOTAMIAN FALLOW DEER (Cervus mesopotamicus).

Larger and brighter coloured than the common fallow deer, with the spots near the middle of the back tending to form longitudinal stripes, and less black on the tail. Antlers of a totally different type, being somewhat expanded at the origin of the trez tine (which is large, and situated some distance above the short brow tine), but at the summit only moderately flattened, and breaking up on the hinder border into several snags.

Distribution.—The mountains of Luristan in Mesopotamian Persia.

	Circum- ference.	Tip to Tip.	Spread inside.	Points.	Width of Palm.	Locality.	Owner.
curve. -29 -21 4	4 1 8 3 8	30½ 14½	24 1 	10+11 9+7	5½ 	Asia Minor Do	
201	5	14		6+5		Luristan Mountains	(Père A. David). Sir Edmund G. Loder, Bart.



Skull and Antlers of extinct Giant Fallow Deer (Irish Elk).

EXTINCT GIANT FALLOW DEER (Cervus giganteus).

(Commonly called "Irish Elk.")

A huge deer, probably standing at least 6 feet at the shoulder, with the antlers enormously expanded, and carrying several large tines on the front border, of which the one above the trez is the longest; the brow tine being often flattened and forked. In its typical form this magnificent deer occurs in the prehistoric deposits of Ireland, England, and probably some of the western districts of the Continent.

Sprea to 7	ıd Tip Fip.	Length insid anti	le of	both a	ntlers	Circum- ference above burr.	Width of Palm.	Points.	Owner.
ft. -1 I	in. 3	ft. 7	in. 5½	ft. 		121	194	17	Mrs. Donaldson-Hudson.
-10	$4\frac{1}{2}$	6	9	13	10	12	20	•••	Earl of Bessborough.
10	2	5	81			97	19‡	19	British Museum.
-10	2						•••	•••	Thomas Bate.
9	8	•				•••		•••	Hon. Walter Rothschild.
9	5	6	2	12	5	11	211	11+11	Viscount Powerscourt.
9	5	6	o	12	10	131	211	15+13	Sir Edmund G. Loder, Bart.

EXTINCT GIANT FALLOW DEER (Cervus giganteus)—continued.

Spre to	ad Tip Tip.	Length round inside of antler.	Length of both antiers across skull.	Circum- ference above burr.	Width of Palm.	Points.	Owner.
9	3	6 2	13 5	10	24	12+10	Duke of Westminster.
9	2	•••	13 6	•••	151		Mrs. Graham Lloyd.
-9	2	•••	11 10	10∤	221	9+9	Dublin Museum.
-9	2		•••	•••		20	Thomas Bate.
8	114	6 г		83	173	10+13	Hon. Charles Ellis.
8	II	5 10	•••	111	17	9+9	Viscount Powerscourt.
8	10	5 9	11 9½	9‡	17	12+11	Duke of Westminster.
-8	10		12 10	•••	131		Mrs. Graham Lloyd.
-8	9	•••			183	9+8	H. J. Elwes.
8	7	5 9½		10	20	12+12	Viscount Powerscourt.
7	6	5 3½		85		10+11	Sir Victor Brooke's Collection.



Antlers of Indian Sambar. From a specimen in the British Museum.

INDIAN SAMBAR (Cervus unicolor).

The typical representative of the Rusine group of deer, in which the antlers are rounded and three-tined, both the bez and trez being wanting, and the summit of the beam simply forked. Height reaching to 5 feet 4 inches at the shoulder. Antlers large and rough, with the brow tine given off at an acute angle to the beam, and the two terminal tines of nearly equal length. Hair coarse and shaggy, uniformly dark umber-brown, with some chestnut on the buttocks, at all ages. Face-glands very large, and capable of being turned inside out. Ears large, and the tail relatively long and bushy. Young uniformly coloured. Weight about 600 lbs., when cleaned about

Lancah Circum-

410 lbs. The wooded hilly districts of India and Ceylon form the habitat of the true sambar, which probably extends into Assam.

The largest, or rather longest, pair of antlers Mr. A. O. Hume ever met with were from the Central Provinces, and measured 48 inches along the curve inside (must have been nearly 50 inches on outside curve). Both brow tines were broken off. They were sent him by Mr. R. Blewitt.

Length on outside curve.	Circum- ference above brow tine.	Tip to Tip.	Widest inside.	Points.	Locality.	Owner.
48	7	Single dr		3	Khandesh	R. H. Madan.
-46 7	•••	49		3+3	Cent. Provinces	R. Wordsworth.
46 <u>1</u>	6₿	24 1	301	3+3	Do	A. O. Hume, C.B.
-46½	9 below brow tine	45			Gurhwal	R. M. Nash.
458	6 §	174.	32 lg	3+3	?	The late Dr. H. Falconer, British Museum.
45	7 3	$22\frac{1}{2}$	33 ‡	3+3	Mayoghur, Cent. Pro-	Sir John Morris, K.C.S.I.
45	7‡	44 8		3+3	Western Ghats	Major-Gen. Sir Arthur Ellis, K.C.V.O.
¹ –45		•••		•••	Cent. Provinces	Bombay Natural History Society's Museum.
-45	9				Orissa	H.H. The Maharaja of Travancore, G.C.S.I.
-45	8	Picked	l up by Jones		Cent. Provinces (?) .	Heighway Jones.
44 1 8	7 2	448	45#	3+3	Rangeer, Do	Col. W. J. Morris.
-44	•••	34		3+3	Central Do	J. D. Inverarity.
² -44	9				Rewa	Major A. E. Ward.
44	6	9‡	248	3+3	Do	Capt. C. F. Pinney.
44	61	19‡	314	3+3	Do	H. E. M. Davies.
43₿	5∯	Single l	horn	3+3	Do	Hume Collection, British
$43\frac{1}{2}$	51	17	28	3+3	Do	Museum. Sir E. P. Bates.
431	6 3	25 1	34	3+3	Cent. Provinces	Viscount Powerscourt.
43 8	6‡	20 8	29 1	3+3	Do	Sir Robert Harvey, Bart.
-431	10			3+3	Do	Capt. C. Hutton Dowson.
-43	101				Cuttack	Indian Museum.

¹ This is only the measurement of a portion of a Sambar antler, and was recorded in the fournal of the Bombay Natural History Society, iii. p. 228. The animal was shot by Mr. R. Gilbert in the Central Provinces, but got away minus this piece of his antler.
² Height at shoulder, 35½ inches.

INDIAN SAMBAR (Cervus unicolor)—continued.

Length on outside curve.	Circum- ference above brow tine.	Tip to Tip.	Widest inside.	Points.	Locality.	Owner.
43	$7\frac{1}{2}$	351	38	3+3	Gwalior	Sir Greville Smyth, Bart.
43	6	24½	30	3+3	Khandesh	A. Cumine.
-43	91/2	26	23	3+3	Cent. Provinces	Capt. J. H. Gwynne.
42 8	58	26	32 1	3+3	Do	A. H. Pollen.
$42\frac{1}{2}$	6 1	15	27 <u>3</u>	4+4	Ghats of Simrol	Col. J. Evans, British Museum.
-42	6 <u>‡</u>	Single	antler	•	?	Sir Edmund G. Loder, Bart.
42	6	26 <u>‡</u>	$35\frac{1}{2}$	3+3	Khandesh	A. Cumine.
41 8	6 1	33 1	35 1	3+3	Western Ghats	Major-Gen. Sir Arthur Ellis, K.C.V.O.
41 3	7	28 1	344	3+3	?	Hon. Walter Rothschild.
418	7 1	314	36 <u>3</u>	4+3	Nepal	The late B. H. Hodgson, British Museum.
411	61	30½	36½	4+4	Cent. Provinces	P. Jay.
41	6 8	181	29	3+3	Do	Duke of Bedford.
¹ -4I	81/2	•••		3+3	N. Gujerat	S. C. Law.
40 8	6	22	27½	3+3	,	Sir Edmund G. Loder, Bart.
40½	$7\frac{1}{2}$	24½		3+3	?	Major James Grant.
40½	5 8	33	37 1	3+3	,	Hume Collection, British Museum.
40 <u>1</u>	6 1	281g	32	3+3	Nilgiris	Sir Victor Brooke's Collection.
-40		20½	•••	3+3	Asirghur	J. D. Inverarity.
40	6	311	33 3	4+4	Cent. Provinces	H. P. Whitney.
40	58	32 1	36	3+3	Ghats of Simrol	Col. J. Evans, British Museum.
-39½	8	26 <u>1</u>	•••	•••	Nimar	Captain J. N. MacLeod.
39 1	6	16 <u>1</u>	201	2+2	Nepal	H.R.H. the Duke of Saxe- Coburg and Gotha.
-39	6	18	28		Satpura Range	Col. J. Biddulph.
-39		28 1	30 1	3+3	Asirghur	J. D. Inverarity.
38 <u>1</u>	6	26	311	3+3	?	Viscount Powerscourt.
-38½	51	38	29½	3+3	Girnar Hill, Kathiawar	LieutCol. L. L. Fenton.
38 1	63	23 1	28 7	3+3	?	Sir Robert Harvey, Bart.
38 1	6 7	30 1	35	3+3	Nilgiris	Sir Victor Brooke's Collection.
					1 Malformed	:= ==:

1 Malformed.

INDIAN SAMBAR (Cervus unicolor)—continued.

Length on outside curve.	ference above	Tip to Tip.	Widest inside.	Points.	Locality.	Owner.
¹ -38		328			Mandla District, Cent.	Capt. B. H. Boucher.
-38	61/2			3+3	Provinces Cent. Provinces	H. Lennard.
38	6 <u>1</u>	293	418	3+3	Chanda District, Cent.	Sir John Morris, K.C.S.I.
38	5₹	28	28	3+3	Provinces ?	Capt. E. H. R. Hibbert.
374	6	29‡	314	3+3	Cent. Provinces	LieutCol. M. Cust.
37½	6 3	218	258	4+3	Do	Major C. S. Cumberland.
37 ½	51/2	43	31	3+3	Chanda District, Cent.	L. Gisborne Smith.
$37\frac{1}{2}$	5 7	29	311	3+3	Provinces Do	C. D. Twopeny.
371	$7\frac{1}{2}$	23	29	3+3	Do	Col. M. M. Bowie.
371	5\$	$23\frac{1}{2}$	31	3+3	Do	C. F. Egerton.
378	51/2	19	232	3+3	Nimar, Cent. Provinces	LieutCol. H. Wade-Dalton.
37 1	5₿	19§	30₽	3+3	Central Provinces .	Major C. S. Cumberland.
37	51/2	201	28 <u>1</u>	3+3	?	M. Loam.
37	61	27 1	28 <u>3</u>	3+3	?	H. C. V. Hunter.
-37	6 _۲ ت	211	24	3+3	Mount Aboo	Viscount Edmond de Poncins.
37	9	13	31	3+3	Mounar Valley	A. H. Sharp.
36₹	6 1	23½	22	3+3	Rajputana	Col. J. Biddulph.
36½	54	243	27	3+3	Benares	Sir Comer Petheram.
36½	5	234	24½	3+3	?	A. M. Caccia.
36½	5₽	29	34	3+3	Nimar, Cent. Provinces	LieutCol. H. Wade-Dalton.
-36	7	19	•••	3+3	?	Count J. Potocki.
				Ce	ylon Specimens.	
321	6 1	24	26]	3+3	Ceylon	A. R. Hay.
311	5	20	17	4+3	Do	Surgeon-Major G. E. Hale,
-30	8?	21½	19§	3+3	Do	D.S.O. J. Ryan.
29	$4\frac{1}{2}$	25	221	3+3	Do	Earl Cairns.
27‡	41	16‡	20	3+3	Do	A. M. Naylor.
-201	5₹	15	15	3+3	Do	Dr. Percy Rendall.
				1 Heig	ght at shoulder, 52 inches.	



Frontlet and Antlers of Malayan Sambar.

Drawn from a Burmese specimen in the British Museum.

MALAYAN SAMBAR or EQUINE DEER (Cervus unicolor equinus).

This local race is nearly as large as the Indian sambar, but the antlers are generally shorter and thicker, with the hinder or inner tine of the terminal fork much shorter than the front one, and arising as a spur from the inner hind margin of the beam, of which the front tine forms the direct continuation; the brow tine also generally longer. General colour of coat of adult darker, usually a light ring round the eyes, the ears smaller, often with a white margin, and the tail very bushy. Young spotted. The distributional area extends from Assam and Cachar through Burma and the Malay Peninsula to Siam, Hainan, Borneo, and perhaps Sumatra.

MALAYAN SAMBAR or EQUINE DEER (Cervus unicolor equinus)—
continued.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.	Owner.
30 2	6 1	174	19	7+7	Borneo	Sir Edmund G. Loder, Bart.
-30½	5 1	274	1 30	3+3	Burma	VetCapt. G. H. Evans.
30 1	48	20 3	211	3+3	Do	British Museum.
29 8	6	16 §	20 1	3+3	Garro Hills, Assam.	Hume Collection, British Museum.
-29 1	5 1	$22\frac{1}{2}$	1 23 $\frac{1}{2}$	3+3	Burma	VetCapt. G. H. Evans.
-28½	5	26 1	1 29½	3+3	Do	Do.
26 3	6	12	13.3	3+3	Garro Hills, Assam.	Hume Collection, British
26½	6 3	11		3+3	Borneo	Sir Edmund G. Loder, Bart.
-26½	6 3	24½		3+3	Perak	Perak Museum.
23 1	7	15 8		3+3	Borneo	H. B. Low, British Museum.
19 §	41	16 7	16 7	4+3	Assam	Hume Collection, British
16 ‡	41	5₽		3+2	Borneo	Museum. W. B. Pryer, British Museum.

FORMOSAN SAMBAR (Cervus unicolor swinhoei).

This race is very closely related to the preceding, from which it is distinguished by its shorter head, concave profile, longer limbs, and certain differences in colour; the lower part of the legs being brownish or whitish yellow, and the bushy tail black all round. It is confined to the island of Formosa.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Points.	Locality.					Owner.		
194		9	3+3	Island of Formosa					British Museum.		
16 1	3ቔ	16	3+3	Do.					Do.		
13 §	4 1	13 1	4+4	Do.					Do.		
5 7	2 1	6 <u>‡</u>	3+2	Do.					Do.		

LUZON SAMBAR (Cervus unicolor philippinus).

Nearly allied to the two last, the height at the shoulder being about 28 inches, the build stout and massive, with the hind-quarters specially elevated, and the form that of a small Malayan sambar. On the head is a blackish streak starting from over each eye to form a line down the middle of the face separated by a band of pale fawn from a moustache-like dark mark in the muzzle.

Distribution.—The island of Luzon, in the Philippines; introduced into the Marianne islands, and described as a separate species under the name of *C. mariannus*.

Length on outside curve.	Circum- ference.	Tip to Tip.	Points.	Localit	y.		Owner.
21	48	51/2	3+3	Island of Luzon		•	Sir Edmund G. Loder, Bart.
19 §	5₺	71/2	4+4	Do.			Capt. Belcher, British Museum.
18 <u>7</u>	51	93	3+5	Do.			Do.
188	5 1	143	4+3	Do.			Do.
18	45	13	4+4	Do.			Do.
16	4	101	3+3	Do.			Sir Edmund G. Loder, Bart.
151	5 1	111	3+3	Do.			Capt. Belcher, British Museum.

BASILAN SAMBAR (Cervus unicolor nigricans).

Length on outside curve.	Circum- ference.	Tip to Tip.	Points.	Local	ity.		Owner.
131	48	12	3+3	Basilan Island		•	A. H. Everett, British Museum.
134	41/2	115	3+3	Do.			Do.

SZECHUAN SAMBAR (Cervus unicolor dejeani).

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.	Owner.
30 8	5 1	15 8	181	3+3	Ta-tsien-lou Szechuan	Paris Museum (type specimen).



Skull and Antlers of Javan Rusa. From a specimen in the British Museum.

JAVAN RUSA (Cervus hippelaphus).

General form, coat, and colour sambar-like; but the ears smaller the tail thin, the hairs on the back banded with coloured rings, and the under parts, chin, and inner sides of buttocks whitish. Antlers comparatively slender and only moderately rough, with the brow tine medium or short, and making a large acute angle with the beam; the hinder or inner tine of the terminal fork much longer than the front or outer one, and forming the continuation of the beam, from the

front or front outer surface of which the front tine arises as an off-shoot; the two antlers enclosing a lyrate space. Young, uniformly coloured. There are two races of this species—one the Javan rusa (C. hippelaphus typicus) of the approximate size of a red deer, and the other the Moluccan rusa (C. hippelaphus moluccensis), from Celebes and the Moluccas, said to be smaller, and without a distinct mane on the neck or tuft to the tail.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.			Owner.			
-36		111		3+3	Mauri	tius (int	trodu	ced)	VAdmiral Sir William Kennedy.		
35½	48	224		3+3	Java	•			Sir Victor Brooke's Collection.		
341	47	17‡	221	3+3	Do.	•			Commander C. Keppel, C.B., R.N.		
¹ 34					Rodrig	guez (in	trodu	ced)	VAdmiral Sir William Kennedy.		
33 1	47	191	25	3+3	Java				Commander C. Keppel, C.B., R.N.		
33	41/2	25½		3+3	Do.	•			Duke of Bedford.		
3 2	$4\frac{1}{2}$	19	16 1	3+3	Do.				The late H. J. H. Platt.		
28]	41/2	1112	$12\frac{1}{2}$	3+3	Do.				Sir Edmund G. Loder, Bart.		
	1 Weight 19 stone 3 lbs.										

MOLUCCAN RUSA (Cervus hippelaphus moluccensis).

See Javan rusa above.

Length on outside curve.	Circum- ference.	Tip to Tip.	Points.	Locality.	Owner.
36 1	4	181	3+3	?	Sir Edmund G. Loder, Bart.
27]	4‡	141	3+3		Duke of Bedford.
14 3 8 1	3 1	•••	3+3	3	Do.
8 3	4	5 1	•••	Batchian .	A. R. Wallace, British Museum.
6₫	3∄	58	•••	?	British Museum.

BAVIAN DEER (Cervus kuhli).

A small deer allied to the Javan rusa (C. hippelaphus), standing about 27 inches at the shoulder, of light build, and of a uniform brown colour, without a dark stripe down the back.

Distribution.—The Bavian Islands, between Borneo and Java.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.	Owner.
93	27	10 1	107	3+3	Bavian Islands	British Museum.



Head of Male Hog-Deer.

HOG-DEER or PARA (Cervus porcinus).

Allied to the Bavian deer, but the antlers larger, the build longer and lower, and the summer coat of the adult, as well as that of the young, spotted with yellowish white. General colour in winter rufous or yellowish brown, somewhat speckled above, and much darker beneath; in summer, upper parts paler and more or less spotted. Antlers on long pedicles, with the hinder time of the terminal fork the shorter. Height at shoulder from about 25 to 29 inches. Weight about 90 to 100 lbs. The largest pair ever seen by Mr. A. O. Hume belonged to a specimen he shot in the Ganges Khadir, near Meerut. It measured 20 along the beam inside, and had a mid-beam girth of 3.5. It was destroyed in the Mutiny, the house in which it was hung, with several hundred others, having been burnt down.

Distribution.—India, throughout the Indo-Gangetic plain from Sind and the Punjab to Assam, thence through Sylhet to Burma and Tenasserim.

Length on outside curve.	Circum- ference above brow tine.	Tip to Tip.	Tip to Tip.		Locality.			Owner.
-23 1	37	12 7	Burma					VetCapt. G. H. Evans.
-213	$3\frac{1}{2}$	7‡	Do.					Do.
-2I	31/2	183	Do.					Do.

HOG-DEER or PARA (Cervus porcinus)—continued.

Length on outside curve.	Circum- ference above brow tine.	Tip to Tip.	Locality.	Owner.		
-2 I	31	154	Pegu	Major-Gen. E. M. Norie.		
-20½	31	121/2	Burma	VetCapt. G. H. Evans.		
-20	31	•••	Nepal	Bombay Natural History Society's		
-201	41/2	174	?	Museum. J. Whitaker.		
-20 1	3	91	N.W. Provinces .	J. Nugent.		
-20	•••	•••	Burma	The late Capt. R. C. Beavan, Indian Museum.		
-198	31	16	Meerut, N.W.P	W. Q. Winwood.		
191	31	135	Dudla Swamp, N.W. Provinces	Capt. W. E. Stobart.		
-191	38 €	•••	Nepal	Bombay Natural History Society's Museum.		
-198			Ganges Khadir .	H. S. King.		
19‡	31	91	Do	A. O. Hume, C.B.		
191	3 1	161	Burma	Sir Victor Brooke's Collection.		
198	38	183	3	H. C. V. Hunter.		
-19	41	518	Upper Burma .	C. W. A. Bruce.		
-18 8	410	15	Do	Do.		
1-18	6 <u>1</u>	19	Ganges Khadir .	Col. R. Pole-Carew, C.B.		
-18	•••	•••	3	W. Gillman.		
173	31/2	9‡	Nepal	The late B. H. Hodgson, British Museum.		
-171		•••	Ganges Khadir .	Major F. D. V. Wing.		
178		91	India	J. Carr Saunders.		
-17	318	•••	Naini Tal Terai .	Major-General Alexander A. A. Kinloch.		
16 7	3	108	?	British Museum.		
-16 1	38 €	148	Nepal	A. E. Leatham.		
16 1	27	8§	India	The late Gen. Hardwicke, British Museum.		
151	2ģ	7 1	Do	Sir Edmund G. Loder, Bart.		
15	$2\frac{1}{2}$	•••	Laos States, Cam- bodia	A. Waley.		

¹ Ten points.



Head of Chital, shot by Lieut.-Col. F. H. Whitby.

CHITAL or AXIS (Cervus axis).

This beautiful species is distinguished from all the other members of the Rusine group, except the Philippine spotted deer (C. alfredi), by the body being profusely spotted with white at all seasons and all ages; the general colour of the upper parts being light rufous fawn, with a dark stripe from the nape to the tip of the tail and a black band on the muzzle. Height at shoulder from 36 to 38 inches; live-weight estimated at about 250 lbs. Antlers supported on short pedicles, long, slender, and moderately rough; the brow tine making nearly a right angle with the beam, and the front tine of the terminal fork, which forms the continuation of the beam, much the longer.

Distribution.—India and Ceylon.

CHITAL or AXIS (Cervus axis)—continued.

Length on outside curve.	Circumfer- ence above T the first point.	ip to Tip.	Points.	Locality.	Owner.
-38 1	4 3	191	3+3	Asirgarh Jungle, Central Provinces	LieutCol. M. Cust.
38	4 3	18	3+3	East Berar	LieutCol. F. H. Whitby. See Illustration.
-38	•••		•••	Narbada Valley .	The late Capt. J. Forsyth.
37½	3 ³	19	3+3	Siwalik Hills	B. R. M. Glossop.
371	41	24 §	4+4	Bassim, C.P	A. O. Hume, C.B.
378	4	$24\frac{1}{2}$	3+3	Déhra Dún	Do.
371	41	16 7	4+5	?	Sir Victor Brooke's Collection.
36 2	4	198	4+4	,	The late W. C. Oswell.
36 1	41/2	25	4+3	Berar	C. H. Seely.
-36 1	41/2	15 §	4+3	Nepal	A. E. Leatham.
-36 1	5₹	194	3+3	Central Provinces .	Capt. M. McNeill.
-36 1	•••	164	4+4	Do	Capt. J. H. Gwynne.
36	41	251	3+4	3	British Museum.
-36	5	15	4+3	Mysore	VetCapt. G. H. Evans.
-36	48	21	4+3	?	Major James Grant.
$35\frac{1}{2}$	4‡	181	4+4	?	Sir Edmund G. Loder, Bart.
351	4	121	5+5	7	G. W. Hatch.
35 1	41	22 7	3+3	Western Ghats .	Major-Gen. Sir Arthur Ellis, K.C.V.O.
-351	4	21	4+4	?	James J. Harrison.
-35	41/2	•••		N. Kanara	Bombay Natural History Society's Museum.
35	41/2	201	4+3	Kota, Rajputana .	Major H. C. Morland.
-35	•••	16 1		Mandla, Central Provinces	Capt. B. H. Boucher.
35	4	244	3+3	Nepal	Capt. G. Roos Keppel.
-35	31/2	18	•••	?	W. S. Murray.
-347	5₹	I 2 8	4+4	N.W. Provinces .	Surgeon-Capt. E. M'K. Williams.
347	41/2	172		?	H.R.H. the Duke of Saxe- Coburg and Gotha.
$-34\frac{1}{2}$				Mirzapore	Indian Museum.
-341		221	3+3	Tapti Valley	J. D. Inverarity.
-341	3 7			North Kanara .	LieutCol. L. L. Fenton.

CHITAL or AXIS (Cervus axis)—continued.

Length on outside curve.	Circumfer- ence above T the first point.	ip to Tip.	Points.	Locality.	Owner.
34	44 .	16 <u>1</u>	3+3	?	Col. R. Pole Carew, C.B.
-34	•••	•••		?	Otho Shaw.
-34	•••	29 .	3+3	Narbada Valley .	J. D. Inverarity.
34	$3\frac{1}{2}$	24	4+3	?	W. H. Cobb.
331/2	48	16 <u>1</u>	3+3	Central Provinces .	H. Douglas Taylor.
33½	48	238	4+5	Do.	Major C. S. Cumberland.
33½	•••		•••	Ahree District, Central Provinces	Colonel F. C. Lister-Kay.
33½	48	195	3+3	?	J. Carr Saunders.
-33½	•••	18	3+3	Central Provinces .	H. St. Lennard.
33 1	3₹	18 1	3+3	?	Col. Scott Chisholme.
33 1	3 2	13	2+3	Yeddacurra	Sir Victor Brooke's Collection.
331	37	148	3+4	3	Sir James Anderson, Bart.
-33	41	16 1	5+4	?	Dublin Museum.
-33 33	6 1 3 2	22 19 3	4+4 3+3	Déhra Dún N.W. Provinces .	Major-General Alexander A. A. Kinloch. Capt. Chambers Didham.
32 7	48	211	3+3	?	Hon. Walter Rothschild.
32 8	37	14	3+3	?	Sir James Anderson, Bart.
32½	41	238	5+4	South India	Sir Victor Brooke's Collection.
-321	41	141	5+4	North Oudh	Major J. W. M. Cotton.
32	4	16 <u>1</u>	4+4	Jabalpur	A. Leslie Renton.
-31½	34	171	3+3	Ceylon	Count Scheibler.
311	3	101	3+3	Do	A. R. Hay.
30 2	3 1	201	3+3	Do	Washington Singer.
30	31/2	15	4+3	Do	LieutCol. H. Wade-Dalton.
29 7	418	201	3+3	Do	British Museum.
28 2	3 ‡	111	3+3	Do	Capt. G. O. Bigge.
28 1	41/2	171	3+3	Do	Surgeon-Major G. E. Hale,
27	31/2	81	3+3	Do	D.S.O. Capt. Lewis Jones.
-25½	3	18	3+3	Do	Dr. Percy Rendall.



Side view of Antlers of Swamp-Deer. From a specimen in the British Museum.


SWAMP-DEER (Cervus duvauceli).

This species belongs to the Rucervine group, in which the antlers resemble those of the Rusine section in the absence of the bez and trez tines, but have the beam regularly forked, and each branch again dividing, so that there are at least four tines. In the swamp-deer the antlers are smooth and flattened, with a long brow tine rising almost at right angles to the beam, which is undivided for about half the total length of the antler, and then splits into a fork, each branch of which is usually simply forked, but sometimes divided in a more complicated manner. General colour bright rufous brown, often speckled near the back. Height at shoulder from 3 feet 8 inches to 3 feet 10 inches; weight about 51 stone 3 lbs.

Distribution.—India, exclusive of Ceylon.

SWAMP-DEER (Cervus duvauceli)—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.	Owner.
41	6 <u>1</u>	351	38	8+5	Central Provinces	Capt. W. W. Hancock.
41	5½		antler) oken	6+6	Do	Do.
39 1	5	351	37½	5+6	Do	Major C. S. Cumberland.
38‡	5½	(shed	antlers)	6+5	Do	C. F. Egerton.
-38		43		6+6	Do	J. D. Inverarity.
-38				6+6	Do	Capt. B. H. Boucher.
378	54	235	$28\frac{1}{2}$	7+6	5	Sir Edmund G. Loder, Bart.
36½	5		20 (outside)	23	Mavella District	Measured by Major A. E. Ward.
36	5	29 1	338	6+5	Nepal	The late B. H. Hodgson, British Museum.
-36		29		6+6	Do	J. D. Inverarity.
$-35\frac{1}{2}$					Chutia Nagpur	Indian Museum.
35월	48	27	29 3		3	Sir Victor Brooke's Collection.
35 1	5 3	221		7+6	Nepal	H. C. V. Hunter.
-34 3		33	•••	12+8	Mymensing	F. B. Simson.
-341	51/2	24 8	$28\frac{7}{8}$	6+5	Nepal	A. E. Leatham.
341	478	104	28	9+6	Mandla, Cent. Provinces	Major A. G. W. Malet.
$33\frac{1}{2}$	48	28 <u>1</u>	32 1	5+5	N.W. Provinces	Capt. Chambers Didham.
32 8	5 1	228	26 8	6+5	Gowhatti, Assam	A. O. Hume, C.B.
328	5	33 1	35	6+6	3	Sir Robert Harvey, Bart.
323	5	27½	•••	7+6	Kuch Behar	Major Henry Streatfeild.
32 <u>1</u>	41	32	$33\frac{1}{2}$	5+5	Central Provinces	Capt. E. D. White.
32 1	5 1	198	$24\frac{1}{2}$	7+7	Nepal	H.R.H. the Duke of Saxe- Coburg and Gotha.
32	4	29	19	4+4	;	Sir H. D. Tichborne, Bart.
31	4 1	•••		•••	Herkapur	Capt. J. H. Purvis.
-30 §	5	315		7+6	Raipur, Central Provinces	Capt. M. M'Neill.
30½	54	28 1	32 3	8+8	Central Provinces	H. Douglas Taylor.
30 <u>₹</u>	41	24‡	27	5+5	?	A. M. Caccia.



Antlers of Schomburgk's Deer. From a specimen in the British Museum.

SCHOMBURGK'S DEER (Cervus schomburgki).

Allied to the preceding, but the antlers smooth, rounded, and more complex; the brow tine very long, frequently forked, and arising nearly at right angles to the beam, which is very short, compressed, and regularly forked, with each of the main branches about equally developed and again forking in a similar manner to terminate in long cylindrical tines. General colour uniform dark brown. Height at shoulder about 3 feet 5 inches.

Distribution.—The northern districts of Siam.

Length on out- side curve.		Tip to Tip.	Widest inside.	Points.		Localit	y.	Owner.
-32 1	5 1	•••		12 + 11	Siam			Sir Edmund G. Loder, Bart.
30 1 29 2 29 2 29 2 28 1 28 1	5	15 §	33	10+10	Do.			British Museum.
29 7	51		26 7	9+8	Do.			Do.
294	51 41	23 5 28 8	318	10+11	Do.			Do.
291	6	191	27	6+7	Do.			J. Carr Saunders.
28 1	5 1	T I	298	11+9		?		Sir Victor Brooke's Collection.
-28.5	5.25	9.5	28.5	10+9	Siam			Indian Museum.
27 A	58	14	24½	9+8	Do.			British Museum.
27 { 27 {	48	18 1	24 7	8+7	Do.			Do.



Skull and Antlers of Thameng. From a Burmese specimen.

THAMENG or ELD'S DEER (Cervus eldi).

Although belonging to the same group as the two preceding species, this deer is readily distinguished by the peculiar form of the antlers. These are rounded and rough, with a long curved brow tine, forming a continuation of the curve of the beam, which is set at right angles to the pedicle; the beam unbranched for some distance, much curved, and finally forked, with the outer prong more subdivided than the inner. Height at shoulder about 3 feet 9 inches; weight from 210 lbs. to 245 lbs. There are two races of this species. First, the Burmese thameng (C. eldi typicus), ranging from Manipur through Burma to the Malay Peninsula, in which the antlers are rounded throughout, and the coat is uniformly umber-brown. And, secondly, the Siamese thameng (C. eldi platyceros), from Siam and Hainan, in which the tips of the antlers are flattened with a number of small snags, and the coat is redder, with yellowish spots.

Length on outside curve, not including brow tine.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Length of brow tine.	Locality.		Owner.
42	5	29	24	3+2		Burma .		A. H. Collins.
41	5 1	27 §	36	5+5		Upper Burma		Surgeon-Major O. E. P. Llovd.
¹ 39€	5	31 8	36∰ (sī	20 nall poir	15 3	Siam .	•	Sir Edmund G. Loder, Bart.
38 7	6 1	25½	30 8	16+19		3		British Museum.

¹ Measured on front of antler from highest tip to tip of frontal tine 56% inches.

THAMENG or ELD'S DEER (Cervus eldi)—continued.

Length on outside curve, not including brow tine.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Length of brow tine.	Locali	ty.	Owner.
381	6 <u>‡</u>	24	30 2	6+5	•••	Manipur		A. O. Hume, C.B.
38	48	29 1	36 3	5+5	•••	Pegu		G. R. Radmore.
371	58	20]	30½	5+5	•••	?		British Museum.
37	41	24 ⁸	32 1	5+5		Burma		A. H. Collins.
36 1	5 1	28 3	314	4+3		Manipur		A. O. Hume, C.B.
354	48	192	26]	4+3		Do.		Viscount Powerscourt.
351	41/2	221	314	6+6		Kyaikto, Burma	Lower	J. W. Clough.
351	6 <u>‡</u>	24	30\$	6+3	18 <u>1</u>	Manipur		A. O. Hume, C.B.
35	41/2	30 1	36	5+5	•••	Lower Bu	rma .	G. R. Radmore.
34 8	51/2	27½	32	6+6	•••	Burma		A. O. Hume, C.B., British Museum.
34½	4	211	27	6+6		Do.		Viscount Powerscourt.
34 1	5	24	32 1	6+6		Do.		Hon. Walter Rothschild.
34	5 7	23½	29 2	10+10		Do.		J. Carr Saunders.
33 2	42	178	248	5+4		Do.		Capt. G. H. Mockler.
33½			40	•••	15	Pegu		Col. R. Pole-Carew, C.B.
33 1	5 1	33 1	37 1	5+5	14	Burma		Sir Edmund G. Loder, Bart.
33	48	28 7	32 1	3+3		Do.		Sir Victor Brooke's Collec- tion.
32 1	58	201	20½	4+3		Do.		Gen. Sir Henry Collett, K.C.B.
32	41/2	18 1	24 1	6+5		Do.		Col. J. Biddulph.
-32	•••	•••			•••	5		Maharaja of Travancore, G.C.S.I.
311	5	25 8	29 1	7+6		5		The late B. H. Hodgson, British Museum.
24 <u>3</u>	3 1	19½	21 8	5+5		Burma		Dr. W. P. Y. Bainbrigge.
-24	41/2	18	181	5+5	••.	Do.		Dublin Museum.
181	3ቔ	121/2	131	6+5		?		The late R. Swinhoe, British Museum.
13 §	3 1		128	6+5		Siam		British Museum.

THAMENG or ELD'S DEER (Cervus eldi)—continued.

Extreme length of right antier round the out- side curve, high- est point, to tip of brow tine.	ference.	Tip to Tip.	Widest span.	Number of points.	Loc	ality.		Owner.
-561	5€	28	37	13	Burma	•		W. F. Lostus-Tottenham.
-55 1	41/2	34½	42 1	12	Do.			Do.
-55	7	28½	38½	12	Do.	•	•	VetCapt. G. H. Evans.
-55	53	30	37	15	Do.			Do.
-55	5		42	13	Do.			Do.
-53½	51/2		38	16	Do.			W. F. Loftus-Tottenham.
-52 3	41/2	24 5	36 2	12	Do.	•		Do.
-51 1	5		331	7	Do.			Do.
-51	5 1	27	37½	10	Do.			VetCapt. G. H. Evans.
−50]	7 1	36	39 1	14	Do.			Do.
-50½	51/2	•••	291	14	Do.	•		Do.
-50 1	5	30½	37	8	Do.			Do.
-50	51	25½	33	12	Do.	•		W. F. Loftus-Tottenham.
-49½	6 <u>1</u>	27	371	9	Do.	•		VetCapt. G. H. Evans.
-49½	5₺	27½	36	10	Do.			W. F. Loftus-Tottenham.
-48½	5 1	23 1	36	13	Do.			Do.
-47½	5	•••	348	13	Do.	•		VetCapt. G. H. Evans.
-46½	6	16	•••	9+6	Do.			S. E. F. Jenkins.
-46	54	20	30	10	Do.			VetCapt. G. H. Evans.
-44½	5	•••	23	10	Do.			Do.
-44	5	22	36	14	Do.			Do.
-43½	$5\frac{1}{2}$	30	33	10	Do.			W. F. Lostus-Tottenham.
$-43\frac{1}{2}$	51/2	16 1	$27\frac{1}{2}$	14	Do.			VetCapt. G. H. Evans.
-43	43	•••	34½	12	Do.	•		Do.
-40½	43	28	33½	9	Do.			Do.
-40	5½		40 <u>‡</u>	10	Do			W. F. Loftus-Tottenham.



Frontlet and Antlers of Indian Muntjac.

INDIAN MUNTJAC or BARKING DEER (Cervulus muntjac).

This species is the typical representative of a genus of small Oriental deer differing widely from all those included in Cervus. antlers, which do not exceed half the length of the head, have a short brow tine and an unbranched beam, and are supported on long skincovered pedicles, continued downwards as convergent ridges on the forehead, whence the name of rib-faced deer. Tufts of bristly hair occupy the position of the antlers in the females. The muzzle has a large naked portion, and although there is generally a pair of glands on the face, there are none either on the hock or the cannon-bone. are spotted, but the adults uniformly coloured. In the Indian muntjac, which is one of the reddish-coloured species, and whose range extends from Ceylon and India through Burma to China, the Malay Peninsula, Sumatra, and Java, the height at the shoulder varies from 20 to 22 The Chinese muntjac (C. reevesi), from Southern China and Formosa, is a much smaller species, also reddish in colour; but in the rare Tenasserim muntjac (C. fea), and the larger but equally scarce hairy-fronted muntjac (C. crinifrons) of Eastern China, the general hue of the body is dark purplish sepia-brown, with white on the buttocks and under surface of the tail. The average height at the shoulder is about 26 inches, and weight about 28 lbs.; a female stands about 23 inches and weighs about 32 lbs. In 1852 Mr. Wilson ("Mountaineer") had a specimen of which Mr. A. O. Hume recorded the measurements as follows: antlers round the curve outside, 9.5 and 9.0 long. days he did not collect horns, and only noted them.

INDIAN MUNTJAC or BARKING DEER (Cervulus muntjac)—continued.

Length on outside curve of antler from burr to tip.	Circumference above burr.	Tip to Tip.	Locality.	Owner.
-108	•••	•••	Java	H. Van Son.
7₩	41/2	$3\frac{1}{2}$	Lombok	Hon. Walter Rothschild.
$^{1}-7\frac{1}{2}$			Near Mussuri	Major A. E. Ward.
71	31	37	Lombok	Hon. Walter Rothschild.
-6 3	$2\frac{3}{4}$	$2\frac{1}{2}$	Perak	Perak Museum.
6 3	21	3	Ranikhet	Surgeon - Major B. W. C. Deeble.
-6 3	•••	•••	Dugshai, Punjab .	J. Johnston-Stewart.
6₽	21	31	Java	J. C. Van Son.
$-6\frac{1}{2}$	•••		Do	H. Van Son.
61/2	2 🖁	$3\frac{1}{2}$	Nepal	The late B. H. Hodgson, British Museum.
6 <u>1</u>	3	38	3	A. O. Hume, C.B.
$-6\frac{1}{2}$	$2\frac{1}{2}$	318	Namba Forest, Assam	Surgeon-Capt. H. S. Wood.
-6 <u>1</u>	2 3	•••	North of Mussuri .	Capt. Harry V. Brooke.
6	25	4 3	Taroy, Burma	VetCapt. G. H. Evans.
-5½		•••	?	Indian Museum.
5½	$3\frac{1}{2}$	$2\frac{1}{2}$	Assam	A. H. Straker.
$5\frac{1}{2}$	2	2	Laos States, Cam- bodia	A. Waley.
5 8	3		, ,	Col. R. J. Heber-Percy.
51	2	31	S. India	Sir Victor Brooke's Collection.
51	31	31	Assam	P. Russel.
-54			Mandla Dist., C.P	Capt. B. H. Boucher.
5	3	3	Assam	Major C. S. Cumberland.
-5	$3\frac{1}{2}$	3ቔ	North Kanara	LieutCol. L. L. Fenton.
			¹ Weight 37 lbs.	

TIBETAN MUNTJAC (Cervulus lachrymans).

Nearly allied to the last, but smaller.

Length on out- side curve of longest antler.	Circumference.	Tip to Tip.	Locality.				Owner.
21/3	18	3	Ningpo				The late R. Swinhoe, British Museum.
$-2\frac{1}{2}$	I ½	3 1	Do.				Dublin Museum.



Head of Hairy-fronted Muntjac.

THE HAIRY-FRONTED MUNTJAC (Cervulus crinifrons).

Length on out- side curve of longest antler.	Circumference.	Tip to Tip.		Loca	lity.		Owner.
I ½	•••	41/2	Ningpo				British Museum.

TENASSERIM MUNTJAC (Cervulus feæ).

Antlers about two inches long.

EUROPEAN ROE (Capreolus vulgaris).

Roe may be recognised by the rudimentary tail, and the medium-sized antlers rising close together and almost vertically from the head, without a brow tine, and regularly forking at a point about two-thirds the total length, with the posterior prong again subdividing, so that the number of points is usually three. There is no gland and tuft on the hock, but one on the upper part of the hinder cannon-bone. In the European roe the height at the shoulder is about 26 inches. In winter the coat is dark speckly brown with a large white rump-patch, but in summer foxy red, with little or no white behind. The range embraces the greater part of Europe as far as the Caucasus, and probably Asia Minor.

EUROPEAN ROE (Capreolus vulgaris)—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
13	4	$14\frac{1}{2}$	Germany		Viscount Powerscourt.
13	4	14½	Do		Do.
13	$3\frac{1}{2}$	10‡	Do		Do.
-13	$6\frac{1}{2}$	81	Do		H.R.H. the Duke of Saxe-Coburg and Gotha.
-13	7	10	Do	•	Do.
-13	61	8	Do		Do.
I 2 4	3	81/2	Do	•	Viscount Powerscourt.
121	4	$9\frac{1}{2}$	Do		Do.
121	44	9	Servia		Do.
$-12\frac{1}{8}$	34	44	Austria		Capt. John Marriott.
$^{1}-12\frac{1}{8}$			Perthshire		R. Moncrieff.
115		6	Monymusk, N.B	•	Sir Arthur Grant, Bart.
1 I I ½			Orton, Speyside .		Sir G. Macpherson Grant, Bart.
111	3	83	Germany		Viscount Powerscourt.
-11‡	6	6	Perth		J. G. Millais.
114	41/2	8‡	Servia		Viscount Powerscourt.
-II I	4 3	•••	Scotland	•	Duke of Bedford.
-11 8	41/2		(Single antler dug up)		J. G. Millais.
11	71	•••	Ross-shire	•	H. M. Warrand.
-11		71	Aberdeenshire .		Col. Gordon Cumming.
11	518	71/2	Ross-shire	•	H. M. Warrand.
11	38	83	Germany	•	Viscount Powerscourt.
11	7 1 8	6 7	Sligo, Ireland		Sir Henry Gore Booth, Bart.
101	•••	•••	Beaufort, Ross-shire	•	J. G. Millais.
101	6 <u>1</u>	6	Ballindalloch	•	Sir G. Macpherson Grant, Bart.
101	$3\frac{1}{2}$	54	Inverness	•	C. Macpherson Grant.
10‡	31/2	8‡	,		Major James Grant.
-104		•••	,		J. G. Millais.

¹ and 2 Recorded by J. G. Millais (British Deer and their Horns).

³ Abnormal (Peraque).

EUROPEAN ROE (Capreolus vulgaris)—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.		Loca	lity.		Owner.
-101	31	Single antler	Dorset				A. Du Cane.
10	21	41/2	Ross				Sir Edmund G. Loder, Bart.
9 3	24	31/2	Austria				H.R.H. le Duc d'Orléans.
9 3	81	12 points	Sligo, Ir	elano	ı.		Sir Henry Gore Booth, Bart.
9 7	5	5	Dorset				J. E. Harting.
9 7	3 3	5 3	Moray	•			Sir Victor Brooke's Collection.
91/2	33	41/2		?			Col. Ralph Vivian.
9 <u>1</u>	3 1	7 4	New Fo	r e st			Hon. Gerald Lascelles.
-9‡	51/2	5	Beaufort	, Ros	s-shi	re	J. G. Millais.
-9‡	4	51 ¹ 8	Ross-shi	re			Col. A. J. Warrand.
-91	$3\frac{1}{2}$	7 1 6	Do.				Do.
9	3	61/2	Banff				H. Brinsley Brooke.
83	23	51/2	Glenrinn	ies			J. G. Williams.
85	3	4 3 antiers	Scotland	۱.			Col. Ralph Vivian.
81	6‡	48	Lissadell	l, Ire	land		Sir Henry Gore Booth, Bart.
81	3	41	Scotland	١.			C. C. Branch.
$8\frac{1}{2}$	3	$4\frac{1}{2}$	France				British Museum.
812	243	•••	Islay				Capt. M. M'Neill.
$-8\frac{1}{2}$	41	3	Scotland	l			C. V. A. Peel.
8‡	35	271	Lissadel	1			J. Kenneth Foster.



Head of Siberian Roe.

SIBERIAN ROE (Capreolus pygargus).

Larger than the last, the height at the shoulder varying from 28 to 34 inches; the ears shorter and more hairy, the white rump-patch larger, and the antlers longer and more rugose with numerous nodose snags.

Mr. St. George Littledale says the specimens he observed barked like a "barking deer."

Distribution.—From the Altai and mountains of Turkestan to Siberia, and probably the Caspian provinces of Persia.


SIBERIAN ROE (Capreolus pygargus)—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-18 18	1 63 4	•••	?	Carl Hagenbeck.
16 .	41/2	12	?	Viscount Powerscourt.
151	41	8 <u>1</u>	?	Do.
15	43	•••	Siberia	Sir Edmund G. Loder, Bart.
-141 ³ 6	3 1 5	•••	Do	Paris Museum.
14	31/2	134	Altai, Mongolia .	St. George Littledale.
13 7	27 .	135	Semirechensk Altai	(See illustration.) H. J. Elwes.
132	3	6	Asia?	British Museum.
132	3 3	1112	?	H.R.H. le Duc d'Orléans.
135	4	9	Upper Yenisei Valley)
131	3 1	83	Semirechensk Altai	
131/2	3 \$	9§	Upper Yenisei Valley	H. J. Elwes.
131	48	$12\frac{3}{4}$	Do	
131	$3\frac{1}{2}$	111	S. Siberia	W. A. L. Fletcher.
138	3 18	91	Siberia	Paris Museum.
134	4	91/2	Do	Duke of Bedford.
131	4	87	S. Manchuria ² .	H. E. M. James, British Museum.
134	37	91	Upper Yenisei Valley	H. J. Elwes.
134	4 7	14	Do	Do.
123	3	7 \$	Semirechensk Altai	Do.
I 2 ½	3	63	?	H.R.H. le Duc d'Orléans.
123	31/2	74	Caucasus	H. J. Elwes.
121	31/2	3	?	Duke of Bedford.
118	4	8 1	Manchuria ²	British Museum.

 $^{^{1}}$ Circumference of burr. 2 These probably belong to the somewhat smaller Manchurian Roe.



Skull and Antlers of Siberian Roe shot by St. George Littledale.



Antlers of Père David's Milou Deer at Different Ages. From specimens at Woburn Abbey.

PÈRE DAVID'S MILOU DEER (Elaphurus davidianus).

This remarkable deer differs from all the preceding, except the roes, by the absence of a brow tine to the antlers, which are large and branching, the beam forking at a comparatively short distance above the burr, and the front prong of the fork again dividing, while the hind prong is long and straight. The bushy tail is longer than in any other deer, and the neck of the male is maned. There is a gland-tuft on the upper half of the hind cannon-bone, but none on the hock. In the adult the colour is uniformly tawny, but spotted in the young. Height at shoulder about 3 feet 9 inches.

Distribution.—Northern China; probably unknown in the wild state.

PÈRE DAVID'S MILOU DEER (Elaphurus davidianus)—continued.

	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.	Owner.
32 7	6월	13§	187	8+8	Near Pekin	Sir Edmund G. Loder, Bart.
-30 1	5	351 ¹ 6	35 1 g	11+10	Imperial Park, Pekin	Paris Museum (Type Specimen), Père A. David.
28 1	58	$26\frac{1}{2}$	•••	3 ÷ 3	?	Hon. Walter Rothschild.
27	5 8	20}		6+5	?	British Museum.
¹ 25	5	•••		6+4	Bred in England .	Duke of Bedford.
22	4 Ē			4+3	Do	Do.

¹ Back point of this head measured 28 inches from beam of antler.



Antlers of Virginian Deer. From a specimen in the British Museum.

COMMON AMERICAN or VIRGINIAN DEER (Mazama americana).

With the exception of the wapiti and elk, all the deer of America are distinguished from those of the Old World, save the roe and milou deer, by the absence of a brow tine to the antlers, which are either regularly forked or spike-like, and quite different from those of either the roe or milou deer. In the Virginian deer they are large and complex, with a long sub-basal snag, and the front prong of the main fork developed at the expense of the hinder, and carrying a number of snags on its upper surface. Tail long. A gland-tuft on the hock, and a small cylindrical white one with a black centre near the lower end of the hind cannon-bone. Colour of upper parts chestnut in summer and bluish gray in winter, with the under surface of the tail and the buttocks pure Typically from Eastern North America, where the height at the shoulder reaches to 3 feet 1 inch, but represented by numerous races in other parts of the Continent, which gradually decrease in size and complexity of antlers towards the south, where they extend to Peru, Bolivia, and Guiana. Weight, 12 st. 7 lbs. (F. C. Selous). Commonly called white-tailed deer.

COMMON AMERICAN or VIRGINIAN DEER (Mazama americana)— continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.		Owner.
27 \$	58	Single antler		16	N. America .	•	British Museum.
27 1	48	14 1	19	6+6	Do		Do.
26‡	4 3	9 1	20	20	Do		Major James Grant.
25 ‡	41/2	9	19	11	Maine		H. S. Wellcome.
25 8	48	108	19	15+13	Texas		Capt. F. Cookson.
25 1	4‡	12 7	171	11+9	N. America .		British Museum.
24½	41/2	123	181	6+6	Do		J. Carr Saunders.
24½	41	11 1	19½	6+6	Do		British Museum.
-24	48	1 191		18	Nebraska .		G. B. Grinnell.
-23 3	6	12	16 <u>3</u>	5+4	Wyoming .		James J. Harrison.
23	5	5	16	6+6	B. Columbia .		J. Turner-Turner.
-221	4	1 15 3	•••	12	Medora, N.D.		Theodore Roosevelt.
$21\frac{1}{2}$	33	31	131/2	4+4	New York State		Sir Edmund G. Loder,
21	5	183	19	4+4	?		Bart. L. C. R. Messel.
191	41	12	15	5+5	Wyoming .		F. C. Selous.
19½	31/2	121	151/2	5+5	B. Columbia .		T. P. Kempson.
173	44			27+25	Do		Moreton Frewen.
1 0 12‡	4	8		3+3	Do		J. Turner-Turner.
					1 6		

¹ Spread.

MEXICAN DEER (Mazama americana mexicana).

One of the smaller races of the preceding, the height at the shoulder being about 2 feet 9 inches, and the antlers smaller and simpler.

Distribution.—South Mexico.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.		Owner.		
13 A	38	64	113	3+3	Mexico			British Museum.	
8‡	31/2	6	$6\frac{1}{2}$	3+3	Do.	•		Sir Edmund G. Loder, Bart.	
8 <u>‡</u>	$2\frac{1}{8}$	44	6§	3+4	Do.			Sir Victor Brooke's Collection.	



Head of Mule-Deer. From a specimen in the possession of E. S. Cameron.

MULE-DEER (Mazama hemionus).

Antlers with a much shorter sub-basal snag than in the Virginian deer, beyond which the beam is directed outwards for a short distance, and then curves upwards to form a regular fork, both prongs of which are usually equal, and generally subdivide so as to form five points on each side. Ears very large and heavy; tail moderate, terminating in a bush-like tuft. Gland-tufts on hock and cannon-bone coloured like the leg; the latter of these elongated and situated in the upper half of the cannon-bone. General colour of upper parts reddish tawny in summer, brownish or rufous speckled gray in winter, with a brown

horse-shoe mark on the forehead. Height at the shoulder, 3 feet 3 or 4 inches in the typical form. Weight, 17 stone 2 lbs. (F. C. Selous). Distribution.—The greater part of North America westward of the Missouri, extending from British Columbia to California. There are several local races, among which the South Californian (M. hemionus peninsulæ) is one of the smallest.

	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.	Owner.
30	54		41	17	White River, Colorado	H. A. James.
28 §	$4\frac{1}{2}$	134	173	5+5	Wyoming	Ford G. Barclay.
28 <u>1</u>	5	183	21	6+4	British Columbia .	J. McI. M'Iver Campbell.
28 <u>1</u>			24½		White River, Colorado	Major Maitland Kirwan.
27	5‡	19½	21 §	6 ÷ 5	North America .	Sir Edmund G. Loder, Bart.
27	5‡	19½	$22\frac{3}{4}$	5+5	British Columbia .	D. H. Crake.
-26 7	5			35	North Dakota	Theodore Roosevelt.
-26 3	•••	•••	•••	26	3	W. A. Baillie-Grohman.
26 3	43	19§	201	5+5	Wyoming	Ernest Farquhar.
26 <u>1</u>	5	151	181	6+5	British Columbia .	T. P. Kempson.
26 <u>1</u>	5\$		171	12	Wyoming	Hon. F. Thellusson.
26 g	4 1 8	15 7	22 7	6+6	Do	Capt. F. Cookson.
26	$4\frac{1}{2}$	20	21 3	7+5	Do	The Maclaine of Lochbuie.
26	5	15‡	191	4+4	British Columbia .	J. V. Colby.
-25 §	$5\frac{1}{2}$	143	28	11+8	Frazer River, B.C	A. E. Leatham.
$25\frac{1}{2}$	5½		$21\frac{1}{2}$	8	Do	Sir Peter Walker, Bart.
$-25\frac{1}{2}$	43	•••		28	Montana	P. Liebinger.
$25\frac{1}{2}$	43	18	24 1	5+5	Colorado	E. T. Logan.
$25\frac{1}{2}$	41	27	2 9	5+5	Wyoming	Capt. G. J. Fitzgerald.
25 8	41	19	20	5+5	Do	A. H. Pollen.
25	51	27 8	27 🖁	16+13	Montana	Moreton Frewen.
25	5	$20\frac{3}{4}$	24	5+5	Wyoming	J. L. Scarlett.
25	5	10	163	5+4	Do	F. C. Selous.
-25	54	$17\frac{1}{2}$	$21\frac{1}{2}$	7+5	Do	James J. Harrison.
24½	54	•••	$23\frac{1}{2}$	10	Maine	H. S. Wellcome.

MULE-DEER (Mazama hemionus)—continued.

Length on out- side curve.	Circum- ference.		Widest inside.	•	Locality.		Owner.
$24\frac{1}{2}$	51/2		25	10	Wyoming		T. W. H. Clarke.
24½	5	19	24	5+5	Do		F. C. Selous.
24	5	13\$	17	10	British Colum	nbia	T. P. Kempson.
24	$4\frac{1}{2}$	201	20	6+5	Do.		R. Rankin.
24	44	19‡	$25\frac{1}{2}$	5+5	Wyoming		Capt. J. M'Call Maxwell.
-24	41/2	153	191	5+5	Do		Count E. Hoyos.
$23\frac{1}{2}$	7	18	21	26	British Colum	nbia	J. Turner-Turner.



Head of Mule-Deer. From a specimen shot by J. McI. M'Iver Campbell.


BLACK-TAILED DEER (Mazama columbiana).

Nearly allied to the mule-deer, but of inferior size, with relatively smaller ears and finer hair; but specially characterised by the shorter gland and tuft on the hind cannon-bone, and the larger amount of black on the tail, of which only the basal third of the lower surface is white.

Distribution.—Western North America, from British Columbia to California.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.		Owner.			
$27\frac{3}{4}$	6	r 5 ‡	19‡	9+6	British Colum	bia		G. Wrey.		
-22	41		1 22		Vancouver			Clive Phillipps-Wolley.		
-201	5	17	17	5+7	Do.			A. E. Leatham.		
19‡	41	14	16½	3+3	S. California			Sir Victor Brooke's Collection.		

¹ Outside.



Malformed Marsh-Deer Antlers.

MARSH-DEER (Mazama dichotoma).

Antlers without a sub-basal snag, forking regularly, with both prongs again dividing, and the upper one usually more complex than the lower. Ears large, with white hair internally. Hair long and coarse, reversed on the withers for a short distance. General colour of upper parts bright rufous chestnut in summer, browner in winter; legs black from the knees and hocks downward. No tuft on cannon-bone. Size, approximately that of a red deer. Although ten is the usual number of points, sports are common.

Distribution.—From Brazil to the inner wooded districts of Argentina.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.	Owner.		
24½	5	16	18	5 + 5	Chaco of Paraguay .	Vice - Admiral Sir William Kennedy.		
23½	6 7	26	25	5+5	?	Sir Edmund G. Loder, Bart.		
23 8	5 1 8	152	16 1	5+4	S. America	British Museum.		
231	6		191	12	Do	G. R. Stuart.		
22 §	6 1	20	$20\frac{1}{2}$	5+5	Brazil	Sir Victor Brooke's Collection.		
$-22\frac{1}{2}$	$7\frac{1}{2}$		154	6+6	Argentina	Kenyon Slaney.		

Н

MARSH-DEER (Mazama dichotoma)—continued.

Length on out- side curve.	Circum- ference.	Tip to Tip.	Widest inside.	Points.	Locality.			Owner.		
228	51/2	25		28	Argentina	•	•	A. Vans-Agnew.		
211	51/2	22	16 <u>1</u>	6+6	Paraguay			S. Pulley.		
$21\frac{1}{2}$	5	124	16	5+5	Do	•		Vice - Admiral Sir William Kennedy.		
-21 		193		6+5	Entre Rios			′		
20	51/2	20		4+4	Paraguay			Count Henry Coudenhove.		
-20	68	19‡	183	4+4	Uruguay.	•		Staff-Surgeon J. Dowson, R.N.		
171	41	11		10	Paraguay			Hon. Walter Rothschild.		



Side view of Antlers of Marsh-Deer. From a specimen in the British Museum.

PAMPAS DEER (Mazama bezoartica).

A small deer nearly allied to the last, but with the front prong of the antlers simple, and the hinder one divided. A whorl in the hair on the middle of the back and another at the base of the neck, so that the fur of the withers is directed forwards for a considerable distance. Colour of upper parts light reddish brown, under parts and lower surface of tail white; upper surface of latter black.

Vice-Admiral Sir William Kennedy, in his Sporting Sketches of South America, thus describes it:—"The gama is of a light buff colour, carries a pretty symmetrical pair of horns, generally six points, but emits so strong an odour as to be called the stinking deer. Height at shoulder, 2 ft. 6 in."

Distribution.—Brazil to Northern Patagonia, in open districts.

Length on outside curve.	Circum- ference.	Tip to Tip.	Points.	Locality.	Owner.
148	28	13‡	3+3	Argentina .	British Museum.
141	3	11	3+3	South America.	Sir Edmund G. Loder, Bart.
14	41	111	3+3	Paraguay	Vice-Admiral Sir William Kennedy.
13	$2\frac{1}{2}$	128	3+3	?	W. Livingstone Learmonth.
-12½	$3\frac{1}{2}$	7	12+11	Uruguay	J. Burnett.
121	21	81/2	3+3	?	Sir Victor Brooke's Collection.
1118	3 7 8	131	3+3	3	The late Charles Darwin, British
108	$2\frac{1}{2}$	131	3+3	North Patagonia	Museum. British Museum.
10	21	128	3+3	Argentina .	Count Henry Coudenhove.
71/2	21/4	51/2	3+3	3 .	Duke of Bedford.

PERUVIAN GUEMAL (Mazama antisiensis).

Together with the closely allied Chilian guemal (M. chilensis), this species constitutes a group characterised by the antlers forming a single fork, of which the front prong is the smaller, and curves upwards towards the hinder one. There is no gland-tuft on the hind cannon-bone, the short tail is rather bushy, and the hair coarse and brittle.

Distribution.—The high Andes, from Peru to Northern Chili.

Length on outside curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
9½ 9Ř	7 § 2½	4 8 81	Tinta, South Peru Ceuchepate, Peru (11,000 ft.)	H. Whitely, British Museum. Do.

WOOD BROCKET (Mazama nemorivaga).

The brockets are some of the smallest deer included in the genus *Mazama*, of which they are the typical representatives. They are recognisable by their simple spike-like antlers, the tufted crown of the head, and the radiation of the hair of the face from two whorls, which causes that on the nose to be directed downwards. The present species is distinguished by its small size (height at shoulder about 19 inches), its pale pepper-and-salt brownish or gray colour, the streak on the forehead, and the absence of a gland and tuft on the hock.

Distribution.—Guiana, Colombia, Bolivia, Brazil, and Trinidad.

Length on front of horn.	Girth.	Girth. Tip to		I	ocalit	y.		Owner.	
-48	34	4	Trinidad					Dr. Percy Rendall.	

MUSK-DEER (Moschus moschiferus).

From all living deer except the Chinese water-deer this species is distinguished by the absence of antlers, whose function as weapons is discharged in the male by long upper tusks. The tail is rudimentary, the fur coarse and brittle, and the lateral hoofs are very large. The males have a glandular pouch which secretes the musk from which the species takes its name. Height at shoulder about 20 inches, at rump 22 inches.

Distribution.—The forest districts of the Himalaya as far west as Gilgit, at elevations of 8000 feet or more in summer, to Tibet, Siberia, and Western China.

Length of Tusk. Locality			ity.	у.			Owner.	
-34	Himalaya							J. Johnston-Stewart.
3	Gurhwal		•			•		A. O. Hume, C.B.
3	Amurland					•		British Museum.
23	Himalaya							Rowland Ward.
211	Do.							Dr. Percy Rendall.
21	Kashmir							Sir Edmund G. Loder, Bart.
21	Do.							Major A. Nugent.



A. H. Neumann's Northern Giraffe Skull. From De Winton, P.Z.S. February 1897.

NORTHERN GIRAFFE (Giraffa camelopardalis).

Ihuhla of the Swazis.

Indhlulamiti of the Zulus.

Tuthla of the Basutos.

Ngabe of the Masawaras.

Giri or Halgiri of the Somalis.

Kameel of the Boers.

The long and stilted limbs, the peculiarly-formed head, and the dappled hide, render giraffes distinguishable at a glance from all other living ruminants; with none of which they have, indeed, any very close affinity, although their nearest relatives are the deer. So great are their distinctions, that they constitute a family by themselves—the Giraffida. For many years it was supposed that this family was represented only by a single species; but it is now ascertained that the North and South African giraffes are markedly distinct from one another; the difference being apparently sufficient to admit of their being regarded as species. As a family characteristic, one of the most marked peculiarities of the giraffes is to be found in the horns. These consist of a pair of bony processes arising from the head between the ears, and covered during life with skin. They are never shed; and in the adult are immovably united to the bones of the skull, although separate in young animals. In addition to these, there is a more or less distinct third horn, or boss, situated on the forehead between the eyes. Giraffes have a long, extensile tongue, hairy lips,

and broad, low-crowned molar teeth. There are no tusks in the upper jaw; and the false hoofs are likewise wanting.

In the present species the third horn is well developed, measuring from three to five inches in height in old bulls. The ground-colour of the coat varies from white to fawn; the orange-red or reddish chocolate dark blotches having sharply defined edges, with the spaces between them generally narrow and distinctly demarcated, even in old individuals. Beneath the knees and hocks the legs are white in typical examples.

Distribution.—From the Tana river northwards through the Galla country and Somaliland to Abyssinia, Kordofan, and thence probably across Africa, in suitable districts to Senegambia from about Lat. 15° N. to the Equator. An unusual tall female giraffe from Nigeria has been regarded as indicating a western race of this species, under the name of G. camelopardalis peralta. The giraffes of Nyasaland and the Kilimanjaro districts have been described by Dr. P. Matschie as distinct species, but their right to separation stands in need of confirmation.

SOUTHERN GIRAFFE (Giraffa capensis).

In the southern giraffe the third horn, even in old bulls, is so reduced in size as scarcely to merit that name at all. The ground-colour of the coat varies from white to dull fawn; and the dark blotches, which may be either dun or deep coffee colour, always have the centre darker than the edges, and the latter irregular and not sharply defined, so that in the adult the intervening light spaces are broad. In young individuals, however, the skin shows very clearly defined white lines between the dark blotches, forming a conspicuous network pattern; the margin of the blotches receding with age. In size, the species appears very similar to the last; and in both kinds the old bulls become much darker than the cows or immature males.

Distribution.—Within the last half century this species has ranged from the Orange to the Zambesi rivers. Northward of this latter river on the eastern half of the continent, at least, no giraffe is found for about 12 degrees; but north of the Rufizi river they again appear and continue through German East Africa, reaching westward to Lake Tanganyika, and occurring east of the Mari escarpment and south of the Tana river in British East Africa (De Winton).

SOUTHERN GIRAFFE (Giraffa capensis) -continued.

Total height.	At shoulder.	Locality.	Owner.					
ft. ins. -18 7	ft. ins. 	South-East Africa .	. F. Vaughan Kirby.					
1-18 o	I2 O	South Africa	. The late Sir W. Cornwallis Harris.					
-17 6	Average height	South-East Africa .	. F. Vaughan Kirby.					
& 17 O		W. Matabeleland .	. F. C. Selous.					
§ 16 6	•••	N. Kalahari	. Do.					
-915 6 to 16 0	Average height	South-East Africa .	. F. Vaughan Kirby.					

NORTHERN GIRAFFE (Giraffa camelopardalis).

Total height.	At shoulder.	Locality.	Owner.	
ft. ins. -16 0	ft. ins.	Central East Africa		A. H. Neumann.

¹ Portraits of the Game and Wild Animals of Southern Africa.



Head of Male Prong-horn.

PRONG-HORN (Antilocapra americana).

Although commonly termed an antelope, this animal differs from all the members of the *Bovidæ* by the forking of the horns. These are annually shed from their bony sheaths and replaced by a new pair, which commence to grow up beneath the old ones before they are cast off. In consequence of this difference the species represents a family (*Antilocapridæ*) by itself. Horns generally absent in the female. Ears long and pointed, tail short, and neck maned. General colour chestnut, with a white rump-patch, and white bars on the neck. Height at shoulder, 36 inches; weight, 70 to 80 lbs. clean.

Distribution.—Western United States, from British Columbia to Mexico.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Locality.				Owner.		
-171	$6\frac{1}{2}$		9	N.W. Canada					J. Whitaker.	
-17			20 outside		?				Otho Shaw.	
-16	$6\frac{1}{2}$			N. Dakota					Theodore Roosevelt.	
154	6]	5 7		Wyoming .					St. George Littledale.	

PRONG-HORN (Antilocapra americana)—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Locality.	Owner.
-15 3	51	•••	20 outside	?	Otho Shaw.
-15 §	6	2 7		Teton Mountains	H. Lennard.
15 <u>1</u>	$4\frac{1}{2}$	9 1		Wyoming	W. R. Cookson.
-15‡	57	6]		Do	Count E. Hoyos.
15 1	6 1	7 5		?	Col. Ralph Vivian.
15	5₹	54	104	Wyoming	St. George Littledale.
148	5	5‡		New Mexico	Capt. F. Cookson.
141	5 3	I 2	•••	?	C. F. Bengough.
$-14\frac{1}{2}$			121	Wyoming	T. W. H. Clarke.
14½	6		91/2	Laramie Plains, Wyoming.	Ford G. Barclay.
141	6≸	5‡	•••	Wyoming	LieutCol. Hon. W. Coke.
14‡	5 8	6‡	10	Do	Count Scheibler.
141	6	28		?	Sir Victor Brooke's Collection.
141	7	41		?	J. McI. M'Iver Campbell.
14	5½	101	•••	Wyoming	F. C. Selous.
14	5‡	133		Colorado	E. T. Logan.
14	6	6∯		Wyoming	Sir Peter Walker, Bart.
14	6	3 1		Do	J. B. Gilliat.
14	5½	9‡	•••	?	J. Carr Saunders.
138	51	48		Wyoming	W. W. Ashley.
13½	5½	7‡		Do	Capt. Hugh Fraser.
13½	$5\frac{1}{2}$	7		Do	Capt. G. J. Fitzgerald.
13	$5\frac{1}{2}$	47	•••	Do	Capt. J. M'Call Maxwell.
13	$5\frac{1}{2}$		$5\frac{1}{2}$	Do	Crawford G. Logan.
13	61/2	$5\frac{1}{2}$	$5\frac{1}{2}$	Colorado	Sir Edmund G. Loder, Bart.
13	$6\frac{1}{2}$	$8\frac{1}{2}$	9	Wyoming	Dublin Museum.



Horns of Bubaline Hartebeest. From Julius Jeppe's specimen.

BUBALINE HARTEBEEST (Bubalis boselaphus).

This species brings us to the great family of hollow-horned ruminants, or *Bovidæ*, in which the horns are in the form of unbranched hollow sheaths, which are never shed, and are supported on bony cores. The hartebeests are large antelopes with naked muzzles, abnormally long faces, doubly curved horns, small face-glands, large valvular nostrils (of which the lower lids are covered with short hairs), long, tufted tails, and large lateral hoofs. The present species is the smallest of the group, standing only 43 inches at the shoulder. It has a short pedicle supporting the horns, which are in the form of the letter W, and the colour is uniform tawny, with the tail-tuft black.

Distribution.—Northern Africa (interior of Morocco, Algeria, and Tunisia) and Arabia.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			p to Tip. Locality. Ov			Owner.
141	8§	98	North Africa				British Museum.		
141	63	9 1		;			Col. Ralph Vivian.		
134	$9\frac{3}{3}$. 9 1	Tunisia .				Julius Jeppe.		
13 1	7	5‡	North Africa				British Museum.		



Horns of West African Hartebeest. Shot by J. W. Carroll.

WEST AFRICAN HARTEBEEST (Bubalis major).

Closely related to the preceding animal, of which, when fully known, it may turn out to be merely a local race with more massive horns. Body said to be of a uniform grayish brown, face deep brown, the fore-legs streaked with dark brown or blackish from the knees downwards, and the tail-tuft black.

Distribution.—Gambia, Lower Nigeria, and the interior of the Cameroons.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
25½	113	122	Nigeria	P. A. Clive.
25 8	124	111	Yauri, Hausa States	J. W. Carroll. (See illustration.)
25 1	12‡	138	Near Borgu	Capt. N. C. Welch.
25	113	10‡	Benue River	Julius Jeppe.
231	1112	22½	Yauri, Hausa States	J. W. Carroll.
-23.5	12.5	malformed 6	Togoland	Berlin Museum.
23	121	9‡	Niger Sudan	Hon. Walter Rothschild.
22	11	19\$	Ibi, Nigeria	Capt. A. H. Festing.
213	121	malformed 8‡	Near Borgu	Capt. N. C. Welch.

WEST AFRICAN HARTEBEEST (Bubalis major)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality	·.	Owner.
1 21 $\frac{1}{2}$	121		Lokoja .		Capt. C. A. Wilding.
21½	101/2	134	Do		Capt. A. H. Festing.
21 1	12	121	Gambia .		H. L. Stephen.
-21 1	11	9 1	Do		Paris Museum.
21	11	9‡	Lokoja .		Col. F. D. Lugard, C.B., D.S.O.
21	12	11	Do		F. H. Barber.
21	I I ½	103	Do		A. Ohlsson.
20 3	10‡	7	Nigeria .		British Museum.
20	11	10‡	Do		Mr. Justice Hopley.
♀ 20	. 9₫	6	Lokoja .		A. W. M. Brodie.
-19 8	I I ½	103	Gambia .		Dr. Percy Rendall.
§ 19 3	9	$6\frac{1}{2}$	Lokoja .		F. H. Barber.
9 19	8‡	13	Nigeria .		Major A. J. Arnold, D.S.O.
Ş 18	8‡	14	Do		Hon. Walter Rothschild.
\$ 18	9	7	Lokoja .		The late Dr. Higgs.
♀ 17½	83	$7\frac{1}{2}$	Benue River		Julius Jeppe.
♀ 17‡	9	5 ½	Lokoja .		Col. F. D. Lugard, C.B., D.S.O.

¹ Height at shoulder, 54 inches.



Frontlet and Horns of Tora Hartebeest. Shot by Lieut.-Col. Hon. W. Coke.

TORA or TETEL HARTEBEEST (Bubalis tora).

Horn-pedicle of medium length, the horns themselves in the form of an inverted bracket (—). Height at shoulder about 48 inches. Colour uniformly pale tawny fulvous, with the exception of the tail-tuft and chin, which are black. Although of the same uniform colour, this species is easily distinguished from *B. boselophus* by its superior size, higher gait, and differently shaped horns.

Distribution.—Upper Nubia, Northern Abyssinia and Kordofan.


Length on front curve.	Circum- ference.	Tip to Tip	Locality.		Owner.
21	9	19 1	Sudan		LieutCol. Hon. W. Coke. (See illustration.)
20 7	9 1	141/2	Do		77 337 1. 35 .9 1 10 1
20	98	15§	Do		British Museum.
20	$9\frac{1}{2}$	143	Lake Zuay, Abyssinia		Prince de Lucinge.
193	9	16 1	Dombelas, Abyssinia		British Museum.
194	9	134	Do		Sir Victor Brooke's Collection.
181	83	151	Sudan		Col. Ralph Vivian.
18 1	•••	16,3	Upper Basalam Rive Abyssinia	er,	J. Menges.
18‡	81	111	Nubia		Julius Jeppe.
18	81	141	Settite River		W. D. James.
171	9 3	18	Bogos-land, Abyssinia		British Museum.

TORA or TETEL HARTEBEEST (Bubalis tora)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Local	i ty.		Owner.
17	$9\frac{1}{2}$	131	Settite River			W. D. James.
♀ 14½	6 <u>1</u>	5 #	Sudan .			Hon. Walter Rothschild.
♀ 141	6‡	18	Upper Nubia			Julius Jeppe.
♀13₹	$6\frac{3}{4}$	14	Do.			Sir Edmund G. Loder, Bart.
♀ 13 §	••	148	Abyssinia .			J. Menges.



Head of Tora Hartebeest.



Head of Female and Skull of Male Neumann's Hartebeest.

NEUMANN'S HARTEBEEST (Bubalis tora neumanni).

Nearly allied to the typical tora, but distinguished by the stouter horns, on which the rings are less distinct and do not extend so far round, and the generally richer coloration. The horns are also less divergent. Mr. Walter Rothschild, the describer of this form, gives the following particulars:—

Colour of hair fulvous fawn, much richer on the back, where there are also some darker spots, which may be stains or natural; below very much paler. Chin blackish, tip of tail black. The male is brighter and darker in colour than the female. There are also on the back some patches with longer, thicker, almost whitish-buff hair, perhaps remains of the winter fur.

Description.—East Africa, in the neighbourhood of Lake Rudolph. Discovered by A. H. Neumann.

Length on outside curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
16 1	103	81	N.E. of Lake Rudolph .	•	•	A. H. Neumann.
♀ 13½	7‡	91	E. shore of Lake Rudolph	•		Do.
♀138	7 1	10	?			Lord Delamere.



Head of Swayne's Hartebeest. Shot by Major H. G. C. Swayne.

SIG or SWAYNE'S HARTEBEEST (Bubalis swaynei).

Sig of the Somalis.

Korkei of the Gallas.

Horn-pedicle and horns of the same general type as in the last species. Height at shoulder about 47 inches; weight about 300 lbs. General colour pale chocolate-brown, with white tips to the hairs; face black, except the muzzle and a line between the eyes; shoulders and upper part of fore-legs, a well as a patch on upper part of hindlegs, also black.

Distribution.—Interior of Northern Somaliland and Shoa; in Somaliland it is found on the dry plateau known as the haud.

Length on front curve.	Circumference.	Tip to Tip.	Locality.	Owner.
201	87	26 <u>‡</u>	Somaliland	Major H. G. C. Swayne.
191	$9\frac{1}{2}$	22½	Do.	G. H. Cheetham.
-19‡	•••	241	N. Somaliland	J. Menges.
-19	9 1	18‡	Do	Capt. M. M'Neill.

SIG or SWAYNE'S HARTEBEEST (Bubalis swaynei)—continued.

Length on front curve.	Circumference.	Tip to Tip.	Locality.	Owner.
183	9‡	16	N. Somaliland.	Ford G. Barclay.
181	9‡	19‡	Do	T. W. Greenfield.
183	83	18	Do	Major H. G. C. Swayne.
-18	8‡	103	Do	Count J. Potocki.
18	9	22	Do	B. R. M. Glossop.
174	11	17	Galla Country.	Viscount Edmond de Poncins.
173	83	21	Somaliland .	Count J. de Bylands.
172	10	211	Do	E. Lee Townshend.
172	10	20	Do	Digby Davies.
171	81	19½	Do.	T. W. H. Clarke.
171	9‡	19	Do	Sir Edmund G. Loder, Bart.
171	9	213	Do	J. J. Richardson.
171	10	213	Do	Capt. J. M'Call Maxwell.
171	9 ‡	21	Do	H. A. Bryden.
$-17\frac{1}{2}$	9	19½	Do	C. V. A. Peel.
17‡	81	21	Do	Major C. F. Blane.
-17‡	81	19‡	Do	J. Johnston-Stewart.
171			Do	Major George Douglas.
-17g	1010	18 <u>1</u>	Abyssinia .	Prince A. de Lucinge.
17	9	10	Somaliland .	W. W. Ashley.
17	9	20 1	Do	J. Byng Paget.
17	91/2	19	Ю	Sir H. D. Tichborne, Bart.
17	83	15½	Do	Capt. George Campbell.
17	91/2	187	Do	R. M'D. Hawker.
-17	9	17	Do	A. E. Pease.
17	83	171	Do	Capt. F. C. Quicke.
161	9	121	Do	W. R. Bindloss.
-16½	91/2	15‡	Do	Paris Museum.
161	10	16	Do	J. Kenneth Foster.
16½	9‡	123	Do I	Hon. Walter Rothschild.

SIG or SWAYNE'S HARTEBEEST (Bubalis swaynei)—continued.

Length on front curve.	Circumference. Tip to Tip.		Locality.		Owner
161	101	16	Somaliland		E. T. Marshall.
16 8	9	171	Do		W. D. James.
161	81/2	201	Do		Lord Delamere.
16	8 1	18	Do		H.R.H. le Duc d'Orléans.
16	93	171	Do		Prince Nicolas Ghika.
16	81	15	Do		Capt. C. S. Timins.
16	9	181	Do		R. Wahrmann.
16	10	16 3	Do		J. Benett-Stanford.
9 15 1	6 3	19	Do		Count J. de Bylands.
♀ 15 <u>1</u>	7	11 1	Galla Country	•	Viscount Edmond de Poncins.
- 9 158		•••	N. Somaliland	•	J. Menges.
- 2 13 1	7	19	Somaliland		C. V. A. Peel.



KONGONI or COKE'S HARTEBEEST (Bubalis cokei).

Horn-pedicle moderate, horns bracket-shaped, very short and thick. Height at shoulder about 45 inches. General colour uniform bright fawn, with the lower lip somewhat browner, and the lower part of the rump paler; tail long, with the black tuft ascending some way up the hinder surface.

Distribution.—Eastern Africa, from Usagara northwards to Kilimanjaro and Masailand. This hartebeest is named after Lieut.-Col. Hon. W. Coke, by whom it was first killed. A bull shot by Capt. R. A. J. Montgomerie, C.B., R.N., in Masailand weighed 312 lbs.

Mr. F. J. Jackson says: "This hartebeest ranges as far north as L. Naivasha, south into German territory and east to within a few miles of the coast at the back of the Shimba hills."

Length on front curve.	Circum- ference.	Tip to Tip.		Localit	y.		Owner.
-20 3		•••	East Africa				Julius Jeppe.
194	10	184	Do.				J. Gardiner Muir.
19	9 1	16	Do.		٠	•	Sir Robert Harvey, Bart.
-19	103	14	Do.				Berlin Museum.
185	87	164	Do.	•			F. J. Jackson, C.B.
$-18\frac{1}{2}$	10]	18	West of Mor	mbasa			C. W. Hobley.

KONGONI or COKE'S HARTEBEEST (Bubalis cokei)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
181	91/2	134	East Africa		H. C. V. Hunter.
-18	$7\frac{1}{2}$	$12\frac{1}{2}$	Masailand		Count Scheibler.
$-17\frac{1}{2}$	81	152	East Africa		Do.
178	91/2	11	Do	•	J. Carr Saunders.
17	$9\frac{1}{2}$	12	Do		Lord Delamere.
17	98	13‡	Machakos		Col. F. D. Lugard, C.B., D.S.O.
17	83	141	East Africa	•	S. L. Hinde.
168	8	I 2 3	Do		British Museum.
16 1	9	17‡	Do		Sir Victor Brooke's Collection.
16 1	8§	131	Kilimanjaro		Sir John Kirk, K.C.B.
16 1	101	133	3		C. Steuart Betton.
16 1	10	15	3		Julius Jeppe.
16	9 3	12	East Africa	•	E. Gedge.
15\$	88	10	Taita	•	J. Wray, British Musuem.
15½	104	I I ½	?		Sir Edmund G. Loder, Bart.
9 15₺	7‡	121	East Africa		S. L. Hinde.
₽ 13	63	13	5		Lord Delamere.



Horns of Cape Hartebeest. F. H. Barber's specimen.

CAPE HARTEBEEST (Bubalis cama).


Khama of the Bechuanas. Ingama of the Makalakas.

Horn-pedicle greatly elongated, horns forming a letter V when viewed from the front. Height at shoulder about 48 inches. General colour brownish fulvous, darker than in any of the preceding; face (except between the eyes), back of neck, chin, shoulders, thighs, and tail, black or blackish; lower portion of buttocks with a conspicuous whitish or yellowish blaze, forming a marked contrast to the other colours.

Distribution.—South Africa southwards of the Limpopo, but extending farther northwards along the confines of the Kalahari desert. This fleet and handsome species (the roi hartebeest) is now nearly exterminated in the Cape, although still found in the Transvaal. A few still linger in the old Bushman country in the north-west of Cape Colony. Although practically exterminated in the Orange Free State and in most of the Transvaal (except to the northwest), numbers of hartebeest are to be found in the pleasant country—partly plains, partly open forest—of British Bechuanaland and the Bechuanaland Protectorate. In the North Kalahari and the desert regions about the Botletli River big troops are to be met with. This hartebeest is an extremely good sporting animal, remarkably fleet and enduring, and not by any means easy to bring to bag. Its flesh is very palatable, and its brilliant coat is much sought after by the native tribes.

CAPE HARTEBEEST (Bubalis cama)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-25	•••		Orange Free State .	F. H. Barber. '(See illustration.)
25	11	10	3	C. Rube.
-24 7	11	7	Nata River	H. M. Barber.
$24\frac{1}{2}$	12	$9\frac{1}{2}$	South Africa	C. D. Rudd.
-24 8	10‡	918	Damaraland	Th. Rehbock.
-241	11	7 7	South Africa	Julius Jeppe.
23 1	10	88	Do	The late Sir Andrew Smith, British Museum.
$-23\frac{1}{2}$	1112	11 7	ро	Julius Jeppe.
$23\frac{1}{2}$	I I ½	9 1	Do	Hon. Walter Rothschild.
$23\frac{1}{2}$	1112	101	Khama's Country .	F. C. Selous.
231	108	8 <u>1</u>	Makari Kari Salt Pan	The late J. S. Jameson.
23	12	5½	South Africa	G. Richards.
23	111	6	Do	F. C. Selous.
23	11	11	Do.	A. Beit.
$-22\frac{1}{2}$	11	81	Do	A. Ohlsson.
$22\frac{1}{2}$	103	10 <mark>1</mark>	Do	Sir Edmund G. Loder, Bart.
221	1112	8	Do	A. Moseley.
221	121	6 1	Do	Major R. Hayes Sadler.
-221 ¹ 6	13	11	Do	Paris Museum.
-22	111/2	107	Do	Dublin Museum.
22	ю	8 1	Do.	Sir Victor Brooke's Collection.
-♀20			Orange Free State .	Cape Town Museum.
- 9 19 8	7 11	1111	Damaraland	Th. Rehbock.
\$ 19 3	71/2	9	South Africa	The late J. S. Jameson.
₽ 19 1	8	81	Do	G. Richards.
₽ 19	81	61/2	Kalahari	H. A. Bryden.



Skull and Head of Jackson's Hartebeest.

JACKSON'S HARTEBEEST (Bubalis jacksoni).

Generally similar to the preceding, but without the black blaze on the face, and with the horns less abruptly bent backwards from above. Other differences may be found to exist when the coloration of the body becomes fully known.

Distribution.—The interior of British Central Africa, north of Lake Baringo, Uganda, and probably northwards to the White Nile and westwards to Congoland.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
24 ¹ / ₂	$10\frac{1}{2}$	114	East Central Africa	F. J. Jackson, C.B.
24	10 3	6 7	Do	Do.
23½	12	9	Do	E. Gedge.
$23\frac{1}{2}$	I I ½	6 <u>1</u>	North End of Lake Albert	Col. Trevor Ternan.
231	ΙΙξ	101	Rangata Nyuki	Major A. E. Smith.
23	113	81	E. C. Africa	Sir Edmund G. Loder, Bart.
-22 7	101	112	E. of Victoria Nyanza .	Paris Museum.

JACKSON'S HARTEBEEST (Bubalis jacksoni)—continued.

Length on front curve.	Circum- ference.	Tip to Fip.	Locality.				Owner.
$22\frac{1}{2}$	103	103	E. C. Africa				Capt. E. J. Tickell.
22	103	11	E. of Victoria	Nyar	ıza		Col. F. D. Lugard, C.B., D.S.O.
-22	11	11	E. C. Africa				T. E. Buckley.
21	101	$9^{\frac{1}{2}}$	Do.				Julius Jeppe.
20‡	I I ½	7 ³	Do.				Capt. E. J. Tickell.
19½	1112	$7\frac{1}{2}$	Do.				Lord Delamere.
9 18∤	8	5 7	Uganda .	٠.			Col. Trevor Ternan.
♀ 16 ‡	8	93	E. C. Africa				F. J. Jackson, C.B.
9 16∄	81	$9\frac{1}{2}$	Do.				Julius Jeppe.
ծ 16 <u>1</u>	7월	41/2	Do.			. •	Mr. Justice Hopley.



Skull and Horns of Hybrid Hartebeest. Shot by Major C. II. Villiers.

HYBRID HARTEBEEST (presumed between Bubalis cokei and jacksoni).

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
191	10 1	9 1	Lake Naraku	. J. Ponsonby.
19] 18]	10	9	Mau Plateau	. Major C. H. Villiers.
171	9 1	91/2	West of Eldoma Ravine	. G. E. Smith.



Head of Lichtenstein's Hartebeest.

LICHTENSTEIN'S HARTEBEEST (Bubalis lichtensteini).

Horn-pedicle very short and broad, and the horns themselves much curved inwards towards one another below the terminal backward inclination. Height at shoulder about 50 inches, weight about 300 lbs. General colour tawny fulvous, becoming more rufous along the back, with the chin, tail-tuft, and front of cannon-bones black.

Distribution.—East Africa north of the Sabi River, throughout Nyasaland and Mozambique to Usagara.

Godonko of Zambesia.

Inkulando of the Mashonas.

Kokotombwi in Barotseland.

Konze in the Chilala and Chibisa countries.

Ngondo in the Chinyanja.

LICHTENSTEIN'S HARTEBEEST (Bubalis lichtensteini)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-22 ⁷ 8	11	7 7	?	Mr. Justice Hopley.
-22 <mark>3</mark>			Nyasaland	F. Watkinson.
-21 4	103	9	?	Mr. Justice Hopley.
21½	121/2	9	Batoka Plateau	F. Smitheman.
$-21\frac{1}{2}$			Batokaland	R. T. Coryndon.
21	121	83	Barotseland	H. Timmins.
-2I			?	O. R. Dunell.
-20 4	I 2 🖁	7 1	Portuguese East Africa .	F. Vaughan Kirby.
-20 3			?	Julius Jeppe.
20∯	13	6	Fort Johnstone, B.C.A	C. C. Bowring.
-201 ⁰ 6	1118	7 1 8	British Central Africa	A. S. Hamilton.
20	121	10‡	Congo Free State	Julius Jeppe.
20	121	8	Wami River, East Africa	Sir John Kirk, K.C.B.
20	131	8	Fort Hill, B.C.A	John Yule.
19½	$II\frac{1}{2}$	8 1	S.E. Africa	Hon. Walter Rothschild.
-19 1	113	81/2	British Central Africa	F. Vaughan Kirby.
19‡	10	6 §	East Africa	Sir John Kirk, K.C.B.
19	12	41	Liwondes, B.C.A	C. C. Bowring.
19	113	5‡	Pungwe	J. W. Allen.
19	1112	48	East Africa	British Museum.
-18 7	I I ½	7 1	Barotseland	F. Aitkens.
18‡	12	5 4	Batoka Plateau, Northern Rhodesia	F. Smitheman.
183	12	8‡	Barotseland	Major H. J. Goold-Adams, C.B., C.M.G.
188	118	3 1	Manica Plateau	F. C. Selous.
$18\frac{1}{2}$	12 3	4 1	Pungwe	Hon. T. Thynne.
181	10‡	5	S.E. Africa	F. C. Selous, British Museum.
181	131/2	6	E. Mashonaland	A. Neilson.
181	13	$2\frac{1}{2}$	Songwe, Nyasaland	James Yule.
18‡	12	8	Pangani, East Africa	General Sir Lloyd W. Matthews.

LICHTENSTEIN'S HARTEBEEST (Bubalis lichtensteini)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
18 1	111	51	Wami River, East Africa	•	F. J. Jackson.
181	12	6 1	Pungwe		R. K. Micklethwait.
-18‡	12	6 <u>1</u>	Beira	•	Dr. Percy Rendall.
18	12	34	British Central Africa.	•	J. E. Gough.
-18	123	83	Nyasaland		Major P. W. Forbes.
-18			Beira		Cape Town Museum.
-18	121	6	East Central Africa .		James J. Harrison.
171	113	5‡	Pungwe		C. C. Gouldsmith.
171	12	6 <u>1</u>	Do		H. R. Holden.
-17½	11	34	Do		A. M. Naylor.
-17 8	113	$91^{\frac{1}{6}}$	East Africa		Paris Museum.
17‡	124	3 3	Maskanala		Hon. K. Campbell.
17	123	5₩	Pungwe		Capt. F. H. Lehmann.
17	11	4 1	Lower Shiré		Staff-Surgeon J. Dowson, R.N.
17	10	4 3	East Africa		Dublin Museum.
-17	111/2	51/2	Pungwe		Julius Jeppe.
-17	81	$2\frac{1}{2}$	Do		A. Ohlsson.
16 §	118	6 <u>1</u>	River Sabi, Mashonaland	•	F. C. Selous, British Museum.
♀ 15	87	47	Do		Do.
- 9 15	78	5₿	Portuguese East Africa	٠	F. Vaughan Kirby.
Ŷ 12 1	7	5½	?		Julius Jeppe.



Head of Hunter's Hartebeest. From a specimen shot by H. C. V. Hunter.

HUNTER'S HARTEBEEST (Damaliscus hunteri).

With this species we come to a group of antelopes closely allied to the true hartebeests, but in which the frontal region is not elevated into a horn-pedicle, while the horns themselves form in most cases a lyrate or simple curve, and the face is of medium length. In the present species the slender horns are indeed doubly curved, although without the sudden angulation of the true hartebeests. They slant upwards and outwards, then bend downwards, while their long points are directed upwards. Colour uniform rufous, with a chevron on the face, the inner surface of the ears, and the tail-tuft white. Height at shoulders about 4 feet, and the build light and graceful.

Discovered by H. C. V. Hunter in 1888.

Len	gth.	Circum-	Tip to	Locality.	Owner.
On front curve.	Straight.	ference.	Tîp.	Locality.	Owner.
26‡	22	81	I 2 ½	Bank of the Tana River, East Africa	Sir Robert Harvey, Bart.
24 ⁷ 8	•••	8 <u>1</u>	15½	Do.	H. C. V. Hunter.
245	211	88	9 1	Do.	Sir Robert Harvey, Bart.
♀2 I		54	7	Do.	F. H. Barber.
201		5∰	8 7	Do.	Hon. Walter Rothschild.



Head of Korrigum.

KORRIGUM or SENEGAL HARTEBEEST (Damaliscus corrigum).

Horns with a single slightly lyrate curve. Size medium. General colour reddish, with patches on the face, shoulders, upper part of forelegs, hips, and thighs, and the tail-tuft black; no dark markings on back and feet.

Distribution.—Senegambia and the interior of West Africa. Discovered by Col. Denham and Capt. Clapperton in 1822-24.

Length on front curve.	Circum- ference.	Tip to Tip.	Lo	cali	ty.	Owner.
22½	8₹	113	Gambia		•	Julius Jeppe.
-21 §	98	15 §	Do.		•	Paris Museum.
-21g	9 1	114	Do.			Dr. P. Rendall, British Museum.
19	62	10	Do.			Late Earl of Derby, British
						Museum.
18 3	91	11	Do.			Do.
-15½	7	8	Do.			Dublin Museum.

TIANG (Damaliscus corrigum tiang).

Apparently only a local race of the korrigum, from which it is distinguished, so far as at present known, by its slightly inferior dimensions, and certain differences in the black markings on the face and limbs.

Distribution.—Sennar, Kordofan, and Bahr-el-Ghazal.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
♀ 20 <u>1</u>	6≇	6 7	Bahr-el-Ghazal	The late Consul Petherick, British Museum.
107	6#	7 	Do	Do.

TOPI 127



Head of Topi. From a specimen shot by A. H. Neumann.

TOPI (Damaliscus corrigum jimela).

The southern representation of the last, distinguished by its still smaller size (height at shoulder, 43-44 inches), shorter and more slender horns, and the somewhat brindled appearance of the coat, due to patches of longer hairs on the otherwise short fur. General colour a peculiar purple-brown, with the dark markings less defined and less black than in *D. corrigum typicus*. Weight about 250 lbs.

Distribution.—British East Africa from the Juba to the Sabuki River, and thence to Uganda and Uniamwezi.

Length ou front curve.	Circum- ference.	Tip to Tip.		Owner.			
21	8	61/2	North End of	Lake	Rudo	lph	H. Andrew.
194	6§	38	East Africa				H. C. V. Hunter.
19	71/2	8	Do.			•	Rowland Ward.
18 <u>3</u>	81	6‡	North End of	Lake	Rudo	lph	A. H. Neumann.

TOPI (Damaliscus corrigum jimela)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
$18\frac{1}{2}$	8 <u>a</u>	81	East Africa	Sir Edmund G. Loder, Bart.
181	· 8 ⁸	5‡	Do	J. Carr Saunders.
18½	78	6	North End of Lake Rudolph	Julius Jeppe.
18 1	6 8	5 7	East Africa	Sir Robert Harvey, Bart.
18	8 1	5#	Do	F. J. Jackson, C.B.
171	8	71	Uganda	Hon. Walter Rothschild.
171	7 1	5	East Africa	H. C. V. Hunter.
17	8	7	Do	Sir Robert Harvey, Bart.
₽ 17	52	41/2	North End of Lake Rudolph	A. H. Neumann.
16 2	7	51/2	Do.	F. J. Jackson, C.B.
16 <mark>7</mark>	7 3	$4\frac{1}{2}$	East Africa	Capt. R. A. J. Montgomerie, C.B., R.N.
16 3	6 1	5½	Coast Region between Lanu	Sir John Kirk, K.C.B.
16 3	57	5	and River Juba River Juba	Do.
15 §	7 8	53	Coast Region between Lamu	Hon. Walter Rothschild.
151	81/2	7	and River Juba Do.	E. Gedge.
151	78	5₩	East Africa	F. H. Barber.
$-14\frac{1}{2}$	6 <u>3</u>	63	Coast Region between Lamu	Mr. Justice Hopley.
-144	5	9‡	and River Juba	Berlin Museum.



Head of Bontebok.

BONTEBOK (Damaliscus pygargus).

Distinguished from the preceding species of the genus by the white blaze on the face, which is continuous from the horns to the nose. Height at shoulder about 40 inches. Weight about 200 lbs. Colour of fore part of back rufous fawn darkening into blackish on the rump, flanks, shoulders, front of limbs, and tail-tuft; rump, upper half of tail, under parts, and much of hinder surface of limbs white.

The curious pied marking of the bontebok, its snow-white face, belly, legs, and rump, and the glaze-like, purplish-lilac gloss of the upper part of the coat—which may be described as chestnut upon the neck and warm purplish-brown upon the body—are very remarkable. The horns very nearly resemble those of the blesbok in shape, but the colour is much darker. The blesbok and bontebok formerly gave great sport to the Boers and European hunters. Both of these blaze-faced antelopes run steadily up-wind, "carrying their broad white noses close to the ground like a pack of harriers in full cry."

Distribution.—Cape Colony, south of the Vaal River; now nearly exterminated. Although formerly abounding in tens of thousands on the Karoos of Cape Colony and the plains of the Orange Free State, bontebok are now reduced to a single herd carefully preserved on some flats on the estate of Mr. Vander-Byl, near Swellendam, in the south of Cape Colony.

BONTEBOK (Damaliscus pygargus)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.		
168	6 <u>3</u>	91	?			British Museum.		
16 1	6₽	83	Bredasdorp .			A. C. Campbell.		
1 157	6 7	8 .	Do			F. C. Selous, British Museum		
15½	61/3	81	Do			Hon. Walter Rothschild.		
-15½	6 <u>1</u>	81/2	Do			Dr. W. P. Y. Bainbrigge.		
-15½	61/2	5	Do		•	A. Ohlsson.		
151	61/2	9 8	Do			A. C. Humbert.		
$-15\frac{1}{2}$	6}	9 3	Do			Mr. Justice Hopley.		
15 8	6 1	81	Cape Colony .			Sir Victor Brooke's Collection.		
-15 18	6 1	7‡	Bredasdorp .			Dr. Percy Rendall.		
15	6 3	81	Cape Colony .			F. C. Selous.		
15	6 8	7€	Do. .			Capt. F. Cookson.		
15	6 1	6월	Bredasdorp .			W. S. Curtis.		
-15	6 <u>‡</u>	78	3			Julius Jeppe.		
-15			?			C. T. Jones.		
$-14\frac{1}{2}$			Bredasdorp .		•	Cape Town Museum.		
-141	61/2	88	Do			Julius Jeppe.		
15	6 2	81/2	Cape Colony .		•	F. C. Selous.		
$14\frac{1}{2}$	6 <u>‡</u>	81	Bredasdorp .		•	Hon. Walter Rothschild.		
- ♀ 14 ‡	5 8	78	Do			Julius Jeppe.		
141	5 8	78	Cape Colony .	•	•	F. C. Selous.		
♀ 13½	5‡	78	Bredasdorp .	•	•	Julius Jeppe.		
2 9 13½	5‡	71/2	Do	•	•	F. C. Selous, British Museum.		

¹ Weight, 200 lbs.; height, 412 at shoulder.

² Height at shoulder, 362.

BLESBOK 131

BLESBOK (Damaliscus albifrons).

Nunni of the Bechuanas.

Closely allied to the last (of which it may be only a local race), but with less black on the body and limbs, the blaze divided by a white line between the eyes, and the rings on the horns yellowish.

Formerly to be numbered by hundreds of thousands, the beautiful blesbok has in the last twenty years grown very scarce indeed. can now scarcely be reckoned as a South African beast of chase, being only met with in small numbers on a few Boer farms in the Transvaal and the Orange Free State. Yet, thirty or forty years ago, blesboks often literally darkened the face of the land with their innumerable legions. The north of the Cape Colony, Griqualand West, the Free State, and the plains of the Western and Southern Transvaal may be described as the true home of this charming antelope in the old days. In 1848 Gordon Cumming speaks of a sight he beheld in the blesbok country. "The plains," he says, "exhibited one purple mass of graceful blesboks, which extended without a break as far as my eyes could strain; the depth of their vast legions covered a breadth of about six hundred yards." What a contrast with the scarcity of the present day! Distribution.—Northern plains of Cape Colony, Orange Free State,

Transvaal, and Bechuanaland: now nearly exterminated.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
181	54	12‡	South Africa		Sir Edmund G. Loder, Bart.
-18	$6\frac{1}{2}$	10 §	Orange Free State.	•	F. R. N. Findlay.
17‡	$6\frac{3}{4}$	7	Transvaal		Abe Bailey.
-17 1	7	71/2	Do		Dr. W. P. Y. Bainbrigge.
-16 1	7	71/2	Orange Free State .		Count E. Hoyos.
-163	6	$9\frac{1}{2}$	Do		Julius Jeppe.
- 8 16 1	61	83	Do	•	Do.
-16 1			?		O. R. Dunell.
16 1	7	8	Orange Free State.		Capt. H. D. Livingstone.
16 1	6 8	7 1	?		Capt. G. F. Henry.
161	61	81	South Africa		Sir Edmund G. Loder, Bart.
16	6 <u>8</u>	8	?		Capt. Lord Douglas Compton.

BLESBOK (Damaliscus albifrons)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner
-16	••		?		Cape Town Museum.
-15%	61/2	5½	?		Mr. Justice Hopley.
-153		•••	?		C. T. Jones.
15	6₹	5½	?		A. Beit.
151	63	8‡	?		Capt. F. H. Lehmann.
151	5 1 8	$7\frac{1}{2}$?		C. H. Akroyd.
15‡	$6\frac{1}{2}$	1 10	Driesontein, O.F.S.		F. C. Selous, British Museum.
15‡	6 1	53	South Africa		Hon. Walter Rothschild.
15‡	61	6 <u>1</u>	Zululand		Major-Gen. Sir Arthur Ellis, K.C.V.O.
15 1	$6\frac{1}{2}$	73	South Africa		Hon. Walter Rothschild.
-15	6₹	81	?		Julius Jeppe.
9 15	5 1		Orange Free State .		Capt. H. D. Livingstone.
14‡	61	83	Cape Colony		H.R.H. the Duke of Saxe-Coburg and Gotha.
-14 1	518	7	Transvaal		F. Vaughan Kirby.
14	63	$7\frac{1}{2}$	Do		Sir Victor Brooke's Collection.
14	$6\frac{1}{2}$	$7\frac{1}{2}$	Heidelberg	•	R. H. Sawyer.
-14	8‡	6∤	Transvaal	•	T. E. Buckley.
-14	6	$7\frac{1}{2}$?		A. Ohlsson.
-14	7 3	128	?		Mr. Justice Hopley.
14	4 1 8	6 <u>년</u>	?		Dr. W. P. Y. Bainbrigge.
131	5	5₹	?		F. C. Selous.
124	45	7 ½	Driefontein, O.F.S.		F. C. Selous, British Museum.

¹ Weight, 180 lbs. Height at shoulder, 39½ inches.



Head of Sassaby. From a specimen shot in Mashonaland by F. C. Selous.

SASSABY or BASTARD HARTEBEEST (Damaliscus lunatus).

Incolomo of the Matabele. Ingalowana of the Basutos. *Inkweko* of the Masubias. Inyundo of the Makalakas. Kaboli in the Barotse country and Mzanzi of all Zulu tribes Lake Ngami country.

Luchu of the Masaras. M'tengo in the Chilala and Chibisa countries. Mzanci of the Swazis.

Unchuru of the Makubas.

Horns starting obliquely outwards, with a single upward and backward lunate curve. Height at shoulder nearly 4 feet. General colour dark chestnut rufous, with the face, shoulders, hips, upper portions of limbs, and tail-tuft black, and the region of the groin and margin of the ears white.

The sassaby has the reputation—in the opinion of all hunters who have tested its speed—of being the fleetest and most enduring animal in South Africa. Were it not that this handsome antelope, in common with its near relative the Cape hartebeest, happens to be lacking in presence of mind, it would very seldom fall to the sportsman's rifle. The sassaby is not difficult to circumvent. A troop can be often turned from its course, or brought to a halt, by firing a bullet or two over the heads of the fleeing animals. Or if the leader of the troop can be wounded and turned out, the rest of the herd become confused and now and again offer easy shots. Like the Cape hartebeest the sassaby has extremely drooping quarters. The skin of one of these antelopes, freshly killed, is very beautiful, the wonderful smoothness and the brilliant purplish bloom of the coat being specially noticeable. Distribution.—South-East Africa, from north of the Orange River to the Zambesi, and westward to Lake Ngami, and northwards to British Central Africa (see below).

Length on front Tip to Tip. Circum-Owner. Locality. ference. curve. Mashonaland . Sir John Willoughby, Bart. 15% 71 153 155 71 113 Sir Edmund G. Loder, Bart. 151 8# 121 Daka, S. of Victoria Falls F. C. Selous. 121 S.E. Africa The late Sir Andrew Smith, British 151 Do. . -154 71 I 23 Julius Jeppe. 15# 71 Do. . Sir Victor Brooke's Collection. 12 83 Do. . G. Richards. 15 ΙIĘ 15 7 11 Do. . A. Moseley. 144 8 10 Lebombo Mountains F. Vaughan Kirby. 142 7 115 Pungwe . J. W. Allen. North of Delagoa Bay 142 71 12 A. Cameron. 8 Matabeleland . J. Brander Dunbar. -I4½ 151 S.E. Africa James J. Harrison. -14½ 15 F. C. Selous, British Museum. 141 75 11 Mashonaland . 141 7 1 Do. . Do. 111 Pungwe . Capt. G. F. Henry. 141 7 10 141 Matabeleland . Rev. Dr. R. J. Nevin. 7 10 - 9 14t 6 Do. . Dr. W. P. Y. Bainbrigge. 9 W. Van Ness. 141 Do. . 73 13 1 141 71 121 Bangweolo Flats, British F. Smitheman. Central Africa -141 81 11 Pungwe Count E. Hoyos.

¹ The locality noted is of great interest, as indicating the existence of this species north of the Zambesi.

SASSABY or BASTARD HARTEBEEST (Damaliscus lunatus)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.		
141	71/2	101	Mashonaland .			Hon. Walter Rothschild.		
-141	71/2	13	S.E. Africa .			Dr. W. P. Y. Bainbrigge.		
142	8	14	${\bf Matabeleland}$.			Hon. R. A. Ward.		
-141			Do			C. T. Jones.		
141	64	98	Pungwe			C. M. Swire.		
14	71	10	Do	•		Marquis of Hamilton.		
-14	6 <u>å</u>	12 7	Do	•		A. Beit.		
-14	8	121	Do	٠		T. E. Buckley.		
14	7 1	103	Do			Hon. T. Thynne.		
14	7	1112	Do			R. K. Micklethwait.		
♀ 13 2	6 §	1112	${\bf Mashonal}$ and .		•	F. C. Selous, British Museum.		
9 13 7	6‡	12	Do			H. and C. Beddington.		
₽ 13½			Lebombo Mts.			F. Vaughan Kirby.		
₽ 13½	•••	117	Mabonga River			H. M. Barber.		
\$ 13	6	134	Matabeleland.			Lord Brackley.		



Horns of Brindled Gnu. From Julius Jeppe's specimen.

BRINDLED GNU (Connochætes taurinus).

Ee-vumba of the Makalakas.

Ikokoni of the Basutos.

Inkone-kone of the Amandebele.

Inkongone of the Swazis and Zulus.

Minyumbwe of the Batongas.

Numbo of the Masubias.

Unzozo of the Makubas.

From their near relatives the hartebeests the gnus, or wildebeests, are distinguishable at a glance by their grotesque shape and smooth horns, as they also are by their habits. The short, broad, and massive head has a blunt and bristly muzzle, and tufts of coarse hair on the forehead and chin; the chin-tuft also extending on to the throat. The horns, which are placed on the crown of the head, are approximated at their bases, especially in old bulls, and are nearly smooth, more or less flattened at the bases, but almost cylindrical at the tips; the curvature being at first outwards, or outwards and downwards, and then bending upwards at the tips. An abundant mane of stiff, upright hair clothes the back of the neck; and the tail is covered with longer and softer hairs, reaching considerably below the hocks. It is from the equine form of the tail that these animals were long popularly known by the name of "horned horse." The hoofs are characterised by their narrow form.

The blue wildebeest, as this species is called at the Cape, is a large animal, standing about 4 feet 3 inches at the shoulder. Its most characteristic features are the outward direction of the horns, which are but little expanded at the base, and not unlike those of a buffalo, the uniformly black tail, and the absence of long hair on the lower part of the chest and belly. The general colour varies from grizzled roan to blackish slaty brown, with more or less distinct vertical dark stripes on the sides of the neck and fore-quarters; these stripes being most conspicuous in the lighter-coloured specimens. Typically, the fringe of hair on the throat, like the mane and tuft on the forehead, is black. Distribution.—Formerly ranging from the north of the Orange River for a long distance up East Africa, the brindled gnu is now practically

exterminated in the Orange Free State and the adjacent districts south of the Limpopo. It is, however, still to be met with in parts of Griqualand West and of the Kalahari, as well as in British Bechuanaland; while in Khama's country and Rhodesia, and thence north-



Head of Bull Nyasaland Gnu. From Sir H. H. Johnston's *British Central Africa*, published by Messrs. Methuen.

wards through Central and East Africa, it is comparatively abundant, or, at all events, was so before the ravages of the rinderpest, and it also occurs in Mozambique. In spite of its clumsy and ungainly appearance, the brindled gnu is a rapid mover; and even when severely wounded will not unfrequently succeed in making good its escape from the mounted hunter.

It is generally distributed in South-East Central Africa, and north of the Zambesi is represented by the Nyasaland gnu (C. taurinus johnstoni), which Mr. F. Vaughan Kirby found also in the Mozambique province.

Widest outside.	Widest inside.	Length on front curve.	Breadth of palm.	Tip to Tip.	Locality.		Owner.
	29	31	¹ 13‡	15‡	Sabi Flats		Dr. R. P. Mitchell.
$-33\frac{1}{2}$	$28\frac{3}{4}$		61	24	Do.		B. Secretan.
$32\frac{1}{2}$	29	20	4	26 <u>1</u>	P. E. Africa		F. Vaughan Kirby.
	28½	218	51	178	Matabeleland		G. H. M. Banks.
•••	28½	20 ³	5	16	Pungwe .		C. C. Gouldsmith.
² 31 ½	28	•••	8	18 1	P. E. Africa		F. Vaughan Kirby.
31	•••	23	1 I I 1	191	S.E. Africa		The late J. A. Nicolls.
30½			•••	•••	Do.		F. C. Selous.
301	•••	•••		17	Do.		F. H. Barber.
•••	$27\frac{1}{2}$	$23\frac{1}{2}$	4‡	16 3	?		Julius Jeppe.
	-27	21	41/2	20½	?		Mr. Justice Hopley.
	-26 3	29	5	163	?		Julius Jeppe.
	26½	211	4§	20½	Matabeleland		Major R. Hayes-Sadler.
	26½	19½	413	17	Do.		F. G. Shaw.
	-26½	311	•••	17	Delagoa Bay		H. T. and A. H. Glynn.
301	26 <u>1</u>	22	41	201	Beira .		Rowland Ward.
•••	-26	31	43	21	Lydenburg		F. H. Barber.
•••	26	221	6	18 1	Mababe Plain	•	F. C. Selous.
	26	$21\frac{1}{2}$	7	171	Pungwe .		Frank Harris.
	25½	22	41	17	S. Africa .	•	G. Richards.
•••	♀ 25	221	41/2	10	P. E. Africa		F. Vaughan Kirby.
29	251	21	41/2	183	Zululand .		LieutCol. Hon. W. Coke.
-28 1	25		11	13	?		F. J. Newnham.
•••	24 3	203	43	163	Pungwe .		Capt. F. H. Lehmann.
	-24½	313	¹ I4	16½	Sabi Flats		James J. Harrison.
	24½	20½	5	173	Matabeleland		Rev. Dr. R. J. Nevin.
		¹ Circ	cumference.		² Height at shou	ldei	r, 53½.

BRINDLED GNU (Connochetes taurinus)—continued.

Widest outside.	Widest inside.	Length on front curve.	Breadth of Palm.	Tip to Tip.	Locality.	Owner.
•••	241	20	4	17§	Pungwe .	P. B. Vander-Byl.
	241	21	41/2	151	Do	Marquis of Hamilton.
28§	241	22	5 1	19‡	S. Africa .	Sir Edmund G. Loder, Bart.
	241	152	41	191	Pungwe .	Ford G. Barclay.



Head of Brindled Gnu.

 241	20 ³	41/2	171	Barotseland	H. Timmins.
 $23\frac{3}{4}$	191	41/2	161	Pungwe .	H. R. Holden.
 234	20	51/2	134	S. Africa .	C. D. Rudd.
 231	19‡	$^{1}11\frac{1}{8}$	178	?	H. Atkinson.
 $23\frac{1}{2}$	21½		181	S. Africa .	T. E. Buckley.
 23½	24§	7	154	Beira .	A. M. Naylor.
 $23\frac{1}{2}$	20 1	4§	16]	Mashonaland	J. A. Jameson.

¹ Circumference.

BRINDLED GNU (Connochætes taurinus)—continued.

Widest outside.	Widest inside.	Length on outside curve.	Breadth of Palm.	Tip to lip.	Locality.	Owner.
	$23\frac{1}{2}$	$20\frac{1}{2}$	¹ I 2	145	Botletli River .	H. A. Bryden.
	-23	22 7	411	17 7 7	North of Great Namaqualand	Th. Rehbock.
	-23		1 1 3 ½	14	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	A. Ohlsson.
	-23	20	1 14 $^{1}_{2}$	161	S. Africa	Dr. W. P. Y. Bainbrigge.
	-22 5	2 I $\frac{7}{8}$	6‡		Pungwe	Count E. Hoyos.
	20	18‡	4	13 1	Benguela, Angola	G. W. Penrice.
• • • •		21			Damaraland .	Cape Town Museum.
	♀ 17½	181	18	8	?	Dr. W. P. Y. Bainbrigge.
	27	19½	41/2	15 1	Tushila Plains, B.C.A.	G. N. Barclay.
	24 1/2	19	4	163	B.C.A	J. E. Gough.
	224	18½	41/2	15	Tushila Plains, B.C.A.	C. C. Bowring.
	16½	121/2	1 10	12	E.C.A	James J. Harrison.

¹ Circumference.



Head of White-Bearded Gnu.

WHITE-BEARDED GNU (Connochætes taurinus albojubatus).

Swahili name Nyumbu.

Apparently a local race of the brindled gnu, distinguished by its slightly paler coloration, and the yellowish white throat-fringe, a few whitish hairs being also mingled with the mane.

The widest part of the front of the horns is somewhat different from the corresponding portion of the typical variety, as will be seen by reference to the illustrations.

Distribution.—East Africa, Athi plains, Ukambani, north of Kilimanjaro.

Widest inside.	Length on front curve.	Breadth of Palm.	Tip to Tip.	Locality.			Owner.
2 27 1	22	6	15½	East Africa			Lord Delamere.
² 28			16	Do			Sir John Willoughby, Bart.
² 27	20	1 12	17	Do			R. P. Carroll.
26 1	22∯	•••	183	Do			H. C. V. Hunter.
25 8	22 ⁷ / ₈		161	Do			F. J. Jackson, C.B.
25	20½	1121	$12\frac{1}{2}$	Do			Rowland Ward.
24	20 ह	•••	151	Do			Sir Robert Harvey, Bart.
231/2	18‡	4 1	12	?			S. L. Hinde.
		1	Circumfe	rence.		2 O	utside.

WHITE-BEARDED GNU (Connochestes taurinus albojubatus)—continued.

Widest inside.	Length on front curve.	Breadth of Palm.	Tip to Tip.	Locality	7.		Owner.
² 24	17		15	East Africa			W. Astor Chanler.
22	20	51	16 <u>‡</u>	Athi Plains			E. Gedge.
21 2	19\$	5	131	East Africa		•	Capt. R. A. J. Montgomerie, C.B., R.N.
218	18 1	4	17‡	Athi Plains	•		Col. F. D. Lugard, C.B., D.S.O.
$21\frac{1}{2}$	21	5‡	151	East Africa	•		Capt. J. W. Pringle.
211	2111	1 1 2 1	173	Do			Lionel Decle, Paris Museum.
$21\frac{1}{2}$	19	43	15	Do			E. J. L. Berkeley.
22	19	4	151	Do			Henry Charrington.
₽ 19 <u>1</u>	15	3	131	Do			Lord Delamere.
211	19§	5 ₹	$12\frac{1}{2}$?			Hon. Walter Rothschild.
20½	16 1	4	151	East Africa			Do.

¹ Circumference.

² Outside.



Skull and Horns of White-Bearded Gnu.



Horns of White-Tailed Gnu. From F. H. Barber's specimen.

WHITE-TAILED GNU (Connochætes gnu).

This southern species is the true gnu, being formerly known to the Hottentots by that name, while, by the colonists, it is termed the black wildebeest. Its inferior size (height at shoulder about 3 ft. 10 ins.), the downward curvature of the horns at starting and their great expansion at the base, the pure white tail, and the abundant fringe of long hair on the chest and fore part of the belly, serve at once to distinguish it from the brindled gnu. The general colour is uniform deep amber-brown, passing into black. Females are much smaller than males; and have the horns more slender and less expanded at the base.

Distribution.—The northern range of this species was approximately limited by the Vaal, or northern branch of the Orange River. Like most of the large animals of the Cape it is now rapidly approaching extermination, being apparently at present represented by herds of a few hundred head preserved by the Boers of the Orange Free State. On the plains of the latter country, as well as on the Karoos of Cape Colony, it was formerly found in vast herds, generally in company with quaggas. Fierce and treacherous in disposition, it was especially characterised by its habit of indulging in grotesque capers and frolics on the approach of strangers; a practice totally unknown to its cousin on the farther side of the Vaal River.

WHITE-TAILED GNU (Connochates gnu)—continued.



Head of White-Tailed Gnu.

Length on front curve.	Breadth of Palm.	Tip to Tip.	Locality.	Owner.
-30 8	1 22 1	14	Kalahari	Dr. F. H. H. Guillemard.
-30	107	171	?	Mr. Justice Hopley.
-28	•••	•••	Colesburg	F. H. Barber. (See illustration.)
-27 7	7	11	Orange Free State .	Count E. Hoyos.
$-27\frac{1}{2}$		•••	Wynburg	F. H. Barber. (See illustration.)
$-27\frac{1}{2}$				Cape Town Museum.
-27½	81	13	Colony ?	Mr. Justice Hopley.
26½	8	14½	South Africa	Sir Edmund G. Loder, Part.
$-26\frac{1}{2}$	71/2	74	Orange Free State .	Julius Jeppe.
-261	1 201	123	Do	Dr. W. P. Y. Bainbrigge.
-26	118	16 <u>‡</u>	Do	The Maclaine of Lochbuie.
25	63	15‡	Do	A. Payne-Gallwey.
$-24\frac{1}{2}$	81/2	8	Do	Julius Jeppe.

¹ Circumference.

WHITE-TAILED GNU (Connochætes gnu)—continued.

Length on front curve.	Breadth of Palm.	Tip to Tip.	Locality.	Owner.
24½	6 <u>3</u>	$15\frac{1}{2}$	Orange Free State .	Julius Jeppe.
24	81	15	Do	Hon. Walter Rothschild.
24	81	134	Do	C. D. Rudd.
23	7 \$	174	Do	A. H. Neumann.
23	71/2	15	Do	Bloemfontein Museum.
223	71	15	Do	F. C. Selous.
22	81	154	Do	F. C. Selous.
-22 1	91/2	112	Do	Julius Jeppe.
-2218	8	143	Do	Dr. Percy Rendall.
22	9	171	Do	Hon. Walter Rothschild.
-22	9	112	Do	Julius Jeppe.
213	10§	1 8 §	Do	British Museum.
$-21\frac{1}{2}$	1 20	121	Do	A. Ohlsson.
21	10	181	Do	Dublin Museum.
20	51	113	Do	Julius Jeppe.
♀ 19 2	1 1 3 1 2	103	Do	Dr. W. P. Y. Bainbrigge.
ş 16	4 2	1112	Do	Bloemfontein Museum.
- 2 15½	5	143	Do	Julius Jeppe.
♀14	5	11	Do	Rowland Ward.

¹ Circumference.



Horns of Common Duiker. From a specimen shot by F. C. Selous.

COMMON DUIKER (Cephalophus grimmi).

Puti of the Bechuanas. Impunsi of the Matabele.

Impungi of the Swazis and Zulus. Imputi of the Basutos.

Gwapi of the Chinyanjas.

The common duiker, or "diver," is the southern representative of an extensive group of small or medium-sized antelopes, mostly confined to Africa, but also containing one genus and species from India. all of these the muzzle is naked; face-glands of a more or less elongated form are present, as are false hoofs; the tail is of medium length, the knees have no tufts of long hair, and the females are provided with four teats. The horns, which are short and straight, are generally present in both sexes, but are smoother and more slender in the does than in the bucks; and the upper molar teeth have low crowns, with square grinding surfaces. The African duikers have a single pair of horns, generally present in both sexes, and more or less hidden by a tuft of long hairs growing from the crown of the head; and the face-glands are arranged to form a bare line of pores on each side of the muzzle. The present species—the true duikerbok—is the largest member of a group of three duikers, characterised by the horns (usually absent in the females) inclining upwards at a sharp angle above the plane of the profile of the nose. Other features of the subgroup are the long and pointed ears, the general yellowish or grayish colour, devoid of dark markings, save for a brown nose-spot. Although essentially a southern form, the common duiker, which measures about 23 inches at the shoulder, ranges on the west coast as far north as Angola, and on the eastern side of the continent to British East Africa and Somaliland. Weight about 25 lbs. Throughout its habitat the timid duiker is to be met with wherever sufficient covert exists; and its furtive, squatting, dodging habits are most aptly indicated by

its name. Met with either singly or in pairs, it is never found far away from covert, strictly avoiding both open plains and steep, rocky mountains. Absence of water is, however, no bar to its existence, as it thrives in the heart of the Kalahari. It is occasionally hunted with foxhounds; its flesh is but moderately good.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-6	25	3 1	S. Africa	James J. Harrison.
-5≸	$2\frac{1}{2}$	$2\frac{1}{2}$	Do	Julius Jeppe.
-5½	2 1	27	Selinya, Khama's	F. C. Selous.
51/2	21	28	country Cape Colony	Mr. Justice Hopley.
-5 1	•••		S. Africa	Lionel Phillips.
- 9 5 1	11/2	3	Do	Dr. W. P. Y. Bainbrigge.
5 1	21	11	Zululand	Capt. L. O. Williams.
-51		•••	?	O. R. Dunell.
-5 1	21	11/2	?	Julius Jeppe.
-54	2 1 8	2	Inyamonga, P. E. Africa	F. Vaughan Kirby.
-5 1	2 1	178	Bredasdorp, C. Colony	Mr. Justice Hopley.
5 1	2	2 8	Transvaal	H. T. and A. H. Glynn.
5	21/4	21	S. Africa	G. Richards.
5	21/4	21/2	Do	F. C. Selous.
-5	$2\frac{1}{2}$	3	Transvaal	F. Vaughan Kirby.
-5	•••		Natal	T. E. Buckley.
5	21/2	3 1	Northern Rhodesia .	F. Smitheman.
41/2	•••	27	Algoa Bay	F. C. Selous, British Museum.
41/2	21/4	14	South Africa	G. H. M. Banks.
41/2	2 1	$2\frac{1}{2}$	Do	R. A. Cooper.
-4½	21	$2\frac{1}{2}$	Mashonaland	A. Ohlsson.
41/2	2	23	South Africa	Dr. W. P. Y. Bainbrigge.
-4½		•••	?	Cape Town Museum.
-4½		•••	?	C. T. Jones.
48	2	1 8	Matabeleland	Lord Brackley.
48	2 1	21	Chanda	C. C. Bowring.

COMMON DUIKER (Cephalophus grimmi)—continued.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
48	2	25	Zululand	A. Cameron.
41	1 7	23	E. Africa	Sir Robert Harvey, Bart.
-41	21	2 } \$	Barberton	Dr. Percy Rendall.
418	2	11/2	Benguela, Angola .	G. W. Penrice.
418	17	25	Natal	J. Wahlberg, British Museum.
41	$2\frac{5}{16}$. 21	Near Borgu, West	Capt. N. C. Welch.
4	2	23	Barotseland	R. T. Coryndon.
4	2	1 7	East Africa	F. J. Jackson, C.B.
♀4	18	11	North of Great Nama- qualand	Th. Rehbock.
4	178	112		H. and C. Beddington.
4	2	I 👼	Matabeleland	W. W. Ashley.
♀ 3월	1 ½	18	Transvaal	Julius Jeppe.

ABYSSINIAN DUIKER (Cephalophus abyssinicus).

Abyssinian name Midaku.

Belonging to the same sub-group as the common duiker, and agreeing with the crowned duiker (C. coronatus) in size, this species differs from the latter by the general colour of the fur being grayish brown, instead of bright yellow. It was discovered by the traveller Rüppell, and its habits have been well described by Mr. W. T. Blanford in his Geology and Zoology of Abyssinia.

Distribution.—The highlands of Abyssinia.

Length on front.	Circum- ference.	Tip to Tip.	Locality.		Owner.
-3 18	$2\frac{34}{16}$	2	Abyssinia		Prince A. de Lucinge.
$-3\frac{1}{2}$	1 7	13	Do.		Sir Victor Brooke's Collection.
-31 ³ 6	2	1] §	Galla country		Viscount Edmond de Poncins.
23	1 7	11	Abyssinia		British Museum.



Head of Blue Duiker.

BLUE DUIKER (Cephalophus monticola).

Ipiti of all the Zulu tribes.

In the great majority of the duikers, that is to say all except the three included in the group mentioned above, the horns, which are generally present in both sexes, slope backwards either in or just below the line of the profile of the nose. The present species is included in a sub-group characterised by the smoky-brown or blackish coloration; and is especially distinguished by the rufous legs and uniformly coloured rump, the height at the shoulder being 13 inches. Weight about 26 lbs. Distribution.—Southern Africa, from the wooded districts of Cape Colony northwards to Benguela on the west, and Nyasaland on the east.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
$-2\frac{1}{2}$	•••	$I^{\frac{1}{2}}$	Cape Colony	F. Vaughan Kirby.
$2\frac{1}{2}$	រេទ្ធ	$I^{\frac{1}{2}}$	Tushila Plain, B.C.A.	Hon. Walter Rothschild.
-21	•••	•••	Knysna, Cape Colony	Cape Town Museum.
-2 1	112	15	Do	Mr. Justice Hopley.
-210	1 §	1/2	Natal	Dr. Percy Rendall.
2	•••	13	Do	F. C. Selous.
2	1 5	13	Benguela	G. W. Penrice.
-2	•••	•••	?	O. R. Dunell.
13	18	13	Benguela	Hon. Walter Rothschild.
-13	11	13	?	A. Ohlsson.
-13	•••	•••	?	C. T. Jones.
\$ 1 11	111	1 1 5	Natal	Dr. Percy Rendall.
-1 5	I§	17	?	Julius Jeppe.
1,6	I 16	13	?	Major H. J. Goold-Adams, C.B., C.M.G.
-1 3	1 5	11	?	Julius Jeppe.

MAXWELL'S DUIKER (Cephalophus maxwelli).

This species belongs to the sub-group characterised by the smoky-brown or blackish colour. In size it is small (height at shoulder about 14 inches). The face is coloured like the back; the limbs, like the body, are grayish brown; and the rump is not parti-coloured. It was first brought to England by Col. C. Maxwell, and described by Major Hamilton Smith in 1827.

Distribution.—West Africa, from Gambia to the Gold Coast.

Length on front.	Circum- ference.	Tip to Tip.	Locality.			Owner.		
1 5	21	2	Fanti					Hon. Walter Rothschild.

RED or NATAL DUIKER (Cephalophus natalensis).

Rooi-Bosch-bokje of the Boers. Msumbi of the Swazis and Malonga. Mkumbi of the Zulus. Izikupu of the Basutos.

Chisimbi of the Lower Zambesi natives.

This duiker is classed in another sub-group of small or mediumsized species characterised by the fulvous, rufous, or chestnut groundcolour; and is specially distinguished by its small size (height at shoulder, 17 inches) and completely uniform coloration, having no dark markings on either the face or body.

Distribution.—Natal, Transvaal, and Mashonaland (including all the forest and bush country of the East Coast), also seen by F. Vaughan Kirby and James J. Harrison near the Lualwa River, Mozambique Province.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-3 7	23	2	?	Julius Jeppe.
-38	2 7	21	Shupanga Forest	. F. Vaughan Kirby.
38	21/2	2 1 8	Sabi Flats	. James J. Harrison.
-3 8	2 8	2	Makongwa Mountains,	Dr. Percy Rendall.
-31	3	2	Barberton (about 6000 ft.) Foothills of Kahlamba	. F. Vaughan Kirby.
-31	21	2 8	?	Julius Jeppe.
-31	28	21	Spitzkop	. H. T. and A. H. Glynn.

RED or NATAL DUIKER (Cephalophus natalensis)—continued.

Length on front.	Circum- ference.	Tip to Tip.	Locality.		Owner.
31/8		2 1 8	Natal		Dr. A. Kraus, British Museum.
-3 1	$2\frac{1}{2}$	1 7	Do		Mr. Justice Hopley.
-3			?		C. T. Jones.
-3	•••	2	?		F. J. Newnham.
3	28	1 3	Natal		Major H. J. Goold-Adams, C.B., C.M.G.
- 9 2 <u>4</u>	I 7/8	I	Foothills of Kahlamba		F. Vaughan Kirby.
24	2	•••			Sir Edmund G. Loder, Bart.
23	2 5	21/4	South Africa	•	Dr. Oakshott.
-2 <mark>3</mark>			Lydenburg		Cape Town Museum.
25	$2\frac{1}{2}$	1]	3		F. E. Potter.
-25	•••	•••	3		O. R. Dunell.
$-2\frac{1}{2}$	24	2	Natal	•	Dr. W. P. Y. Bainbrigge.
$2\frac{1}{2}$	$2\frac{1}{2}$	13	S.E. Africa		G. Richards.
$-2\frac{1}{2}$	13	•••	?		A. Ohlsson.
$2\frac{6}{18}$	•••	218	Near Durban		F. C. Selous.
- 9 1 7	14	I	Barberton		Dr. Percy Rendall.
₽ I∯		I	Near Durban		F. C. Selous.
Q IĀ	1 7	2	?		Hon. Walter Rothschild.

BAY DUIKER (Cephalophus dorsalis).

From the last species the bay duiker, together with some allied West African forms, differs by the presence of a black stripe running along the back and continued to the tail. As a species, its special characters are the dark colour of the hams, and the evenly haired tail, which shows no sign of a tuft, and is parti-coloured.

Distribution.—West Africa, from Sierra Leone to the Cameroons; there being a northern and a southern race.

Length on front.	Girth.	Tip to Tip.	Locality.	Owner.
311	21	2	West Africa	Hon. Walter Rothschild.

BANDED DUIKER (Cephalophus doriæ).

The tiger-like transverse black bands on the orange ground of the back suffice to distinguish at a glance this pretty little duiker from all its kindred.

Distribution.—The interior of the West Coast of Africa from Liberia to Sierra Leone, where it is commonly known as the mountain deer.

Length on front.	Circum- ference.	Tip to Tip.	L	ocality		Owner.
-1 7		I	Liberia		•	Dr. J. Büttikofer, British Museum.
ı∯	11	11	Do.			Hon. Walter Rothschild.

YELLOW-BACKED DUIKER (Cephalophus sylvicultor).

This species is distinguished from all the other members of the genus by its large size, coupled with its blackish coloration, and the presence of a yellowish crest and similarly coloured longitudinal patch on the rump. Height at shoulder, 34 inches.

Distribution.—The West Coast of Africa from Liberia to Angola.

Length on front.	Circum- ference.	Tip to Tip.	1	ocality		Owner.
6 3	•••	41/2	Fanti			British Museum.
6 §	38	48	Do.			Do.
6‡	4	3₺	Liberia	•	•	Dr. J. Büttikofer, British
51/2	38	4 1 8	Gaboon	•		Museum. Sir Edmund G. Loder, Bart.
48		21/4	Fanti	•		British Museum.



Head of Male Four-horned Antelope.

FOUR-HORNED ANTELOPE or CHOUSINGHA (Tetraceros quadricornis).

This antelope is the Oriental representative of the African duikers, from which it is distinguished by the following features. Typically there are two pairs of horns, and the face-glands form deep slits on the sides of the muzzle instead of a long naked line; the females being hornless. Height at shoulder about 25 inches, and weight about 40 lbs. General colour dull rufous brown, becoming whitish beneath, with the muzzle, the outer surface of the ears, and a line down the front of each leg blackish brown, and some white on the outer side of the pasterns. The front horns are not unfrequently mere knobs, and may even be wanting, as in most of the Kathiawar specimens.

Distribution.—Peninsular India south of the Himalayas.

Length on f	of horns ront.	Circum	ference.	Tip to	Tip.	Docain,			Owner.	
Rear.	Fore.	Rear.	Fore.	Rear.	Fore.					
-5	•••					Jhalawar	•		H.H. Maharaja Rana Bahadur of Jhalawar.	
$-4\frac{1}{2}$	$2\frac{1}{2}$	21	I 7	21	I	Lulitpur .		•	SurgGen. Walker.	
$-4\frac{1}{2}$	$2\frac{1}{2}$	•••	•••	• • •		Gurhwal.			Capt. Ging.	
48	$2\frac{1}{2}$	21	28	1 §	27	India .			Sir Edmund G. Loder, Bart.	
-4 1	11/2		• • •			Do			J. D. Inverarity.	

FOUR-HORNED ANTELOPE or CHOUSINGHA (Tetraceros quadricornis)—

continued.

Length on fi		Circum	ference.	Tip to	o Tip.	Locality.	Owner.
Rear.	Fore.	Rear.	Fore.	Rear.	Fore.	zocamy.	5 ************************************
4	$2\frac{1}{2}$	ΙĮ	1 7	3	I 18	Indore	Col. J. Evans, British Museum.
4	2	3	17	13	$2\frac{1}{2}$	Karkote Jungle, near Mhow	LieutCol.G.D.F. Sulivan.
-3 4	•••			•••	•••	India	Indian Museum.
31	13	1 7	2	1 1 g	$2\frac{1}{2}$	Do	Sir Robert Harvey, Bart.
-3∰	1 7			•••		Do	J. D. Inverarity.
$3\frac{1}{2}$	2	31	1 7	1 7	18	Central Provinces .	C. F. Egerton.
3 }	2	1 5	1 §	14	$2\frac{1}{2}$	India	H.R.H. the Duke of Saxe- Coburg and Gotha.
$-3\frac{1}{2}$	21	•••	•••			Do	Indian Museum.
$-3\frac{1}{2}$	34	•••	•••	•••	•••	Mandla, C.P	Capt. B. H. Boucher.
38	1 §	2	13			Central Provinces .	Hon. Walter Rothschild.
3 1	18	2	13			Jhalawar	Mr. Justice Hopley.
38	21	25	14	2	1	Central Provinces .	C. D. Twopeny.
-38						Kathiawar	LieutCol. L. L. Fenton.
-31	2	21	218	2 3	17	Central Provinces .	Bombay Natural History
31	17.	13	1 ½	2 <u>5</u>	1 1	N.W. Provinces .	Society's Museum. Capt. R. B. Fell.
31	111	2		21	11	India	A. M. Caccia.



Head of Salt's Dik-dik.

SALT'S DIK-DIK (Madoqua saltiana).

This species, the Beni Israel of the Arabs, is the typical representative of a group of tiny, slenderly-built antelopes characterised by the elongated, trunk-like nose, of which the tip is almost entirely hairy, the tuft of hair on the crown of the head, the short and almost rudimentary tail, and the minute size of the lateral hoofs. In the present species the last tooth in the lower jaw lacks the third lobe found in almost all ruminants, the trunk is but moderately developed, and the general colour fulvous or rufous fawn, scarcely more rufous on the sides than on the back. Height at the shoulder, 14 or 15 inches.

Distribution.—The Coast Range of Eastern Abyssinia, and adjacent districts of Somaliland.

Length on front.	Circum- ference.	Tip to Tip.	Locality.				Owner.
3	ı	I	Abyssinia				Sir Edmund G. Loder, Bart.
$2\frac{3}{4}$	I	I	Do.				Rowland Ward.
25		13	Do.			•	Sir Victor Brooke's Collection.
$2\frac{1}{2}$	$1\frac{1}{2}$		North Kass	ala			Col. Ralph Vivian.
23		15		?			British Museum.





Skull of Phillips's Dik-dik.

Head of Swayne's Dik-dik.

SOMALI DIK-DIKS (Madoqua swaynei, M. phillipsi, and M. guentheri).

General native name, Sakáro.

Different districts of Somaliland are inhabited by the three species of dik-diks above named. The first of these is allied to Salt's dik-dik, but is of smaller size; it inhabits the northern half of Somaliland. Phillips's dik-dik, which also occurs in Northern Somaliland, is intermediate in size between Salt's and Swayne's dik-dik, from both of which it is distinguished by having the back gray and the sides and shoulders rich rufous or cinnamon. On the other hand, Günther's dik-dik is widely different from both, being nearly allied to the under-mentioned Kirk's dik-dik, from which it is distinguished by the form of the nasal bones in the skull.

Height at shoulder about 14 inches. Weight, \$\displays 6\$ lbs.; \$\footnote{9}\$ 5 lbs. (T. W. H. Clarke).

Sakáro Guyu (Madoqua swaynei). Distribution,—Berbera District.

Sakáro Gol-Ass (Madoqua phillipsi).

Northern half of Somaliland.
Plateau of Central Somaliland.


Sakáro Gussuli (Madoqua guentheri).

Length on front.	Girth.	Tip to Tip.	Locality.		Owner.
$3\frac{1}{2}$	11	1 §	Somaliland		J. Kenneth Foster.
$-3\frac{1}{2}$	•••		Do.		Capt. F. A. Wilson.
3 1	11	13	Do.		W. W. Ashley.
3 1	11	11/2	Do.		Major C. F. Blane.

SOMALI DIK-DIKS (Madoqua swaynei, M. phillipsi, and M. guentheri)—
continued.

Length on front.	Girth.	Tip to Tip.	Localit	y.		Owner.
1-3 1	1 1/2	13	Somaliland			A. E. Pease.
-3 1			Do.			J. Brander Dunbar.
3	•••	1	Do.			Sir Edmund G. Loder, Bart.
3	1 1/2		Do.			Col. Arthur Paget.
-3	•••	13	Do.			Viscount Edmond de Poncins.
$-2\frac{15}{16}$	11/2	118	Do.			Do.
-2 7	1‡	I 1/8	Do.	•		J. Johnston-Stewart.
27	18	Ιğ	Do.	•		Lord Delamere.
23	1	I	Do.			T. W. H. Clarke.
23	11/2		Do.			C. Liddell.
-2§	•••		Do.			J. Brander Dunbar.
1-21/2	ī		Do.			C. V. A. Peel.
$2\frac{1}{2}$		•••	Do.			Rowland Ward.
21/3	I	18	Do.			Major H. G. C. Swayne.
$2\frac{1}{2}$		18	Do.			Sir Edmund G. Loder, Bart.
28	11/2		Do.			C. Liddell.
-21/2	1 3	11	Do.			Dr. Percy Rendall.
1-218	I 1	14	Do.		•	Julius Jeppe.
-13	1 1	1 7	Do.	•		Do.
² -3 ¹ / ₄		11	Do.			C. V. A. Peel.
$^{2}-2\frac{1}{2}$			Njemps .			F. J. Jackson, C.B.
		1 M. phillip	si.		2	M. guentheri.

The following are the dimensions of a specimen of *M. guentheri*:— Length, $23\frac{1}{4}$. Height, $15\frac{1}{4}$. Horns, $2\frac{1}{2}$. Weight, $8\frac{1}{4}$ lbs. Shot at Njemps, 26:9:96, by F. J. Jackson, C.B.



Kirk's Dik-dik.

KIRK'S DIK-DIK (Madoqua kirki).

This dik-dik belongs to a small group of species differing from the one containing M. saltiana by the presence of three lobes to the last tooth of the lower jaw, and likewise by the more decidedly trunk-like character of the muzzle. Of the other members of the group, M. damarensis differs by its superior size, and M. guentheri by the still greater development of the trunk.

Distribution.—East Africa, from Southern Somaliland to Ugogo, most numerous on the coast.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
3	•••	11/2	East Africa .	F. J. Jackson, C.B.
2 ⁸ / ₄		15	Do	British Museum.
2 §	1 1/2	18	Do	Col. Trevor Ternan.
25		112	Do	Sir Robert Harvey, Bart.
2 §	11/2	1 §	Kilimanjaro .	H. C. V. Hunter, British Museum.
2§		13	East Africa .	Sir Robert Harvey, Bart.
2∰	11/2	18	East African Coast	Col. Trevor Ternan.
$2\frac{1}{2}$	•••	15	East Africa .	Hon. Walter Rothschild.
$2\frac{1}{2}$	•••	18	Do	Rowland Ward.
28		18	Manda Island.	Sir John Kirk, K.C.B.

DAMARALAND DUIKER (Madoqua damarensis).

Character mentioned under heading of the preceding species.

Distribution.—Damaraland.

Length on front.	Circum- ference.	Tip to Tip.	Locality.				Owner.
$-2\frac{1}{2}$		•••	Damaraland				Cape Town Museum
-2 ¹ / ₂	11	11/2	Do.				(type specimen). A. Ohlsson.

HEMPRICH'S DIK-DIK (Madoqua hemprichiana).

Further information is desirable as to the right of this form to specific distinction.

Length on front.	Circum- ference.	Tip to Tip	Locality.	Owner.
2.5	18	1 1/4	Abyssinia	 Hon. Walter Rothschild.



Head of Male Oribi from Pungwe, South-East Africa.

CAPE ORIBI (Oribia scoparia).

Inla of the Swazis and Zulus. Pulukudukamani of the Basutos.

The oribis, grysbuck, klipspringer, and their allies constitute a group of comparatively small African antelopes presenting the following characters in common. The muzzle has a naked tip, the head is devoid of a tuft of hair, large face-glands open beneath the eyes by a small aperture on each side, the tail is short or moderate, and false hoofs may or may not be retained. Horns are present only in the bucks, and are short, almost, or quite, straight, with smoothed tips and ridged bases. upper molar teeth have tall and narrow crowns. The dik-diks are nearly allied, but differ by their tufted heads, and elongated hairy muzzles. The oribis, which are the largest members of the group, have normal hoofs and hair, and are specially distinguished by the presence of a bare glandular spot beneath each ear, and of a large opening in the skull beneath each eye-socket. In the Cape species the horns of the bucks are comparatively smooth and slender, with only their basal two inches slightly ridged; the tail being tufted and moderately bushy, with its terminal two-thirds black. Height at shoulders, 24 inches.

Distribution.—Typically, Africa south of the Zambesi. On grassy plains this graceful little antelope is still plentiful in many districts; and the gunner in search of bustard or francolin will often see one of them start up from its form before his pointer, to scud away at a great rate, occasionally making springs from side to side. At close quarters

a charge of shot will suffice to bowl over this diminutive little buck, and thus add some capital venison to the larder. Formerly oribi afforded excellent sport with greyhounds in the eastern districts of Cape Colony.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-7½	23	2§	Zomba, B.C.A	D. MacAlpine.
6 1	21/4		E. Griqualand	Jff. Darling, British Museum.
6	2	21/4	Spitzkop	H. T. and A. H. Glynn.
-6			Natal	Cape Town Museum.
-5 18	2	$2\frac{1}{2}$?	Mr. Justice Hopley.
-5 7	2	3 1	?	Do.
53	2	3 1	Barotseland	H. Timmins.
58	2	2 ³	Bangueolo	F. Smitheman.
5€	2 1 8	17	Transvaal	F. R. N. Findlay.
51/2	178	3	Pungwe	·G. L. Bonham.
$-5\frac{1}{2}$	1 7	31	?	Julius Jeppe.
~5½	2	$2\frac{1}{2}$	Gorongoza, P. E. Africa .	F. Vaughan Kirby.
-5½	17	31/2	?	A. Ohlsson.
5½	2	23	M'peta Island, Upper Zam- besi	R. T. Coryndon.
-58	2	$2\frac{1}{2}$	Do.	Do.
-54		•••	?	O. R. Dunell.
5 1	I	21	S.E. of Lake Bangueolo .	Poulett-Weatherley.
5	2	17	Pungwe	Col. G. A. Percy.
5	13	17	?	F. Vaughan Kirby.
-5	1 7	$2\frac{1}{2}$	M'peta Island	F. Aitkens.
-5	2	$2\frac{5}{16}$	Barberton	Dr. Percy Rendall.
-4 7		•••	Transvaal	C. T. Jones.
-4 3	$2\frac{1}{8}$	2 8	M'peta Island	F. V. Worthington.
4 18	14	1 7	Pungwe	A. C. Humbert.
4 3	2	$2\frac{1}{2}$	Cape Colony	British Museum.
4 <u>3</u>	2 lg	2∰	British Central Africa .	J. E. Gough.
4 ‡	13	2	Manica Plateau M	F. C. Selous.

CAPE	ORIBI	(Oribia	scoparia)—continued.
------	-------	---------	----------	--------------

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
4 3	I	17	3	Julius Jeppe.
41/2	15	17	Pungwe	R. K. Micklethwait.
41/2	14	1 8	3	G. Richards.
-4½	1 7	$2\frac{1}{2}$?	Count E. Hoyos.
-4 8	$2\frac{7}{16}$	2 5	Upper Shiré Valley .	Dr. Percy Rendall.
-4 8	1 3	21	?	Dr. W. P. Y. Bainbrigge.
41	1 §	$2\frac{1}{2}$	Pungwe	A. Cameron.
41 ⁵ 6	14	21	Tuchila Plain, B.C.A.	Hon. Walter Rothschild.
4	13	2 l	Pungwe	G. Micklethwait.

ABYSSINIAN ORIBI (Oribia montana).

Miwaka of the Abyssinians.

Very similar in most characters to the Cape species, but with a shorter and less bushy tail, the tip of which has only a few sparse black hairs. Height at shoulder, $22\frac{1}{2}$ inches. These antelopes are shy and rarely seen in the open, preferring the thick bush and long grass. If disturbed they go at a great pace with their heads quite close to the ground. Their flesh is very good.

Viscount Edmond de Poncins states that a fine male weighed 30 lbs. 13 oz., and stood $19\frac{1}{2}$ inches at the shoulder.

Distribution.—Abyssinia and Bongoland.

Length on front.	Circum- ference.	Tip to Tip.	Locality.		Owner.
5		21/2	Abyssinia .		Sir Edmund G. Loder, Bart.
-4 7	28	2	Hawash, Abyssi	nia	Prince A. de Lucinge.
$-4\frac{1}{2}$	2	2	Do.		Viscount Edmond de Poncins.
48	13	$2\frac{1}{2}$	Dombelas, Abys	sinia	British Museum.
4. 12	2.2	2.3	Sudan .		Prince Henry of Liechtenstein.
4	2	2.6	Do		Do.
3 1	2	21	Do		LieutCol. Hon. W. Coke.

WEST AFRICAN ORIBI (Oribia nigricaudata).

The present species is very close to the Abyssinian oribi, but of smaller size, grayer, and with a distinct black tip to the tail, as in the Cape species. Height at shoulder, 20 inches.

Distribution.—The open country of Senegal and Gambia.

Length on front.	Circum- ference.	Tip to Tip.	L	ocality	7.	Owner.
3 18	1 7	13	Nigeria			Capt. A. H. Festing.
3 3	11/2		Kiama			A. W. M. Brodie.
3	18	23	Gambia	•		Charles B. Mosse, British Museum.

HAGGARD'S ORIBI (Oribia haggardi).

Swahili name, Taya.

Still imperfectly known, but distinguished from all the other oribis by the stouter horns, which are strongly ridged for rather more than the basal half of their length. Height at shoulder about 24 inches. Distribution.—The coast districts of East Africa in the neighbourhood of Lamu. Discovered in 1887 by Mr. J. G. Haggard.

Length on front.	Circum- ference.	Tip to Tip.	L	ocality.	•		•	Owner.
1 5 1 5	21	2	East Central	Afric	а			Hon. Walter Rothschild.
¹ 5 ¹	118	2 3	North end of	Lake	Albe	ert		Col. Trevor Ternan.
4∯	2	21/4	East Africa					F. J. Jackson, C.B.
1 4 9 4 1 6	2	21	Lake Albert				٠.	Col. Trevor Ternan.
1 4 1 8	2	25	Do					Do.
41	13	$2\frac{1}{2}$	East Africa					F. J. Jackson, C.B.

¹ Determination provisional.

ZANZIBAR ANTELOPE (Nesotragus moschatus).

The two elegant little antelopes included in the genus Nesotragus are near relatives of the oribis, from which they are distinguished by the absence of a naked glandular patch below each ear and the want of lateral hoofs. They are further characterised by the horns being directed backwards nearly or quite in the plane of the face, and extending at least as far as the back of the head, as also by the large size of the empty spaces in the skull below the sockets of the eyes, and by the nasal bones. In the present species the horns are short and slender, and the colour, inclusive of the tail-tip, fawn gray. Height at shoulder about 13 inches.

Distribution.—Islets near Zanzibar and adjacent coast from Kilimanjaro to Mozambique.

Length on front.	Circum- ference.	Tip to Tip.	Lo	cality.			Owner.
3‡	1 1/2	2	Coral Islan	ds, Za	nzibar	•	Sir John Kirk, K.C.B.
3 1	18	112	Zanzibar				The late Capt. Speke, British Museum.
3 ‡	11	11/2	Do	•	•	•	Sir John Kirk, K.C.B., British Museum.
25	11	13	Do				Sir Victor Brooke's Collection.



Skull and Horns of Livingstone's Antelope, from a specimen shot by F. Vaughan Kirby.

LIVINGSTONE'S ANTELOPE (Nesotragus livingstonianus).

Lumswi of the Shupanga. Intilengana of the Amatonga.

Distinguished from the preceding species by its slightly superior dimensions (height at shoulder, 15 inches), the longer and thicker horns, more rufous coloration, and the blackish upper surface of the tail.

Two specimens killed by F. Vaughan Kirby measured-

		_	-	Male.	Female.
Extreme length over all, tip of	nose	to end	of tail	27 la ins.	25 3 ins.
Length of tail			•	$3\frac{1}{2}$ "	3 "
Perpendicular shoulder-height	•	•	•	$14\frac{1}{4}$,	13 1 "
" height at croup	•	•		$14\frac{3}{4}$,	$14\frac{1}{4}$,,
Girth of neck				$6\frac{1}{2}$ "	6 "
" behind the shoulder .			•	$14\frac{1}{8}$ "	13 5 ,,
D		3.5			

Distribution.—South-East Africa, from Mozambique to Zululand; the form from Zululand being distinguished as N. livingstonianus zuluensis.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
41/2	13	13	Shupanga Forest .	F. Vaughan Kirby.
-4½	•••	13	?	F. J. Newnham.
48	2 1		Gungunyana's country	H. T. Glynn, British Museum.
-48			Delagoa Bay	F. H. Barber.

LIVINGSTONE'S ANTELOPE (Nesotragus livingstonianus)—continued.

Length on front.	Circum- ference.	Tip to Tip.	Locality.		Owner.
-41		•••	3		O. R. Dunell.
-4 1 8			?		C. T. Jones.
-4 		1	Delagoa Bay .		F. C. Selous.
-4	18	13	?		Julius Jeppe.
-4		•••	?		C. T. Jones.
-4	14	18	?		Julius Jeppe.
3 3	13	2	?		C. D. Rudd.
-3 4	I ½	13	Zambesi		Sir Edmund Loder, Bart.
3 11	13	$2\frac{\delta}{10}$	Delagoa Bay .		Hon. Walter Rothschild.
38	11/2	2 1	Do		G. Richards.
-3 8	15	1 §	Do		Julius Jeppe.
-3 8	13	15	Do		Mr. Justice Hopley.
3 18	•••		Do	٠	Dr. Percy Rendall.
3 1	15	15	?		Major H. J. Goold-Adams, C.B., C.M.G.
3 18	13	18	?		F. E. Potter.
-3 1	17	13	3		A. Ohlsson.
310	2		Northern Zululand		A. H. Neumann, British Museum.
3	15	2 1 8	Zululand		A. Cameron.



Head of Grysbuck, from a specimen shot by F. C. Selous.

GRYSBUCK (Rhaphiceros melanotis).

Isikupi of the Basutos.

From the other members of the oribi group the grysbuck and its cousin the steinbuck are readily distinguished by the horns rising nearly vertically from the skull, in which the open spaces below the eyesockets are unusually small. Curiously enough, the grysbuck retains the lateral hoofs, which have disappeared in the steinbuck. In addition to this feature, the former is distinguished by the fur showing a large admixture of white, instead of being uniformly coloured. Height at shoulder, 22 inches.

Distribution.--South Africa, extending as far north as the Zambesi and Mozambique.

Length on front.	Circum- ference.	Tip to Tip.	Locality.			Owner.
4 š	13	17	?			R. A. Cooper.
-4			Caledon, C.C.			Cape Town Museum.
318	12	1 🖟	Cape Colony .			F. C. Selous. (See illustration.)
-3 1	•••		Limpopo .			C. T. Jones.
-3 1	1 ½	2 🖁	Knysna, C.C.			Mr. Justice Hopley.
318 -34 -34 34	1 ½	ış	7			Major H. J. Goold-Adams, C.B., C.M.G.
-31	$I^{\frac{1}{2}}$	$1\frac{1}{2}$	Knysna, C.C.	_	.'	Mr. Justice Hopley.
-3 la -3 la	1\$	21/2	Komati Poort			F. Vaughan Kirby.
-3	I 1/2	2	?			A. Ohlsson.
-3 2 8 -2 8 -2 8	1 1/2	11	?			G. Richards.
$-2\frac{3}{4}$	14	11	?			Julius Jeppe.



Head of Steinbuck.

STEINBUCK (Rhaphiceros campestris).

Ishah of the Swahilis. Ingaina of the Swazis.

is. Impulupudi of the Basutos. zis. Phuduhudu of the Bechuanas. Umgwena of the Matabele.


As mentioned before, this species is at once distinguished from the grysbuck by the absence of the lateral hoofs and the uniform colour of the fur. The general tint of the latter is bright sandy rufous, becoming richer on the head, and frequently with the tip of the muzzle and a horseshoe-mark on the crown brown. Height at withers about $10\frac{1}{2}$ inches.

Distribution.—Africa south of the Zambesi on the east, and the Cunene on the west; northwards of the Tana to Nyasaland represented by Neumann's steinbok (R. campestris neumanni), distinguished by the absence of dark markings on the head. Probably owing to its small size, the steinbuck has managed to escape the fate that has befallen so many of the South African antelopes. Wherever the traveller journeys on the veldt, he is almost certain to meet this species, which may be regarded as the most familiar game animal of the plains. Like many of its kindred, it is independent of water, and can thus exist in the heart of the Kalahari, where water may not occur for a distance of fully fifty miles. The excellence of its flesh renders it a welcome addition to the bag of the hungry

hunter; and in addition to affording sport with the shot-gun, or, more rarely, the rifle, steinbuck may either be hunted with foxhounds or coursed with greyhounds.

Length on front.	Circum- ference.	Tip to Tip.	Locality.		Owner.
-6 7			Graffreinett		F. H. Barber.
-5 7			Kikumbulin		W. Russell Bowker.
-5 7	17	3	?		Mr. Justice Hopley.
-53	13	1 7	?		J. Whitaker.
-58	2	1‡	?		Julius Jeppe.
1 5 ½		I 1 6	Kimberley		F. C. Selous.
-5 1			?		O. R. Dunell.
5 8	2	13	E. Africa		E. Gedge.
-51		•••	?		Cape Town Museum.
-51	18	27	Cape Colony		F. R. N. Findlay.
5 1	13	2	?		Julius Jeppe.
5	•••	2 <u>1</u>	E. Africa		Sir John Willoughby, Bart.
5	1 7	2 7	S.E. Africa		Dr. W. P. Y. Bainbrigge.
-4 3	13	ış	Do		James J. Harrison.
48	1 1/2	23	Portuguese E. Africa		F. Vaughan Kirby.
48	11	21/3	S. Africa		G. Richards.
41/2	13	2‡	Mashonaland		Jff. Darling, British Museum.
41/2	11/2	2 l	?		Major H. J. Goold-Adams, C.B., C.M.G.
41/2	13	1 §	S. Africa	•	H. and C. Beddington.
41/2	15	25	Do	•	F. C. Selous.
$-4\frac{1}{2}$		•••	Matabeleland		T. E. Buckley.
-41/2	17	$3\frac{1}{2}$	Barberton		Dr. Percy Rendall.
48	I§	2 3	Tana Valley		Sir Robert Harvey, Bart.
-4 3		•••	Matabeleland		Major A. St. H. Gibbons.
41	1 1/3	2	Do		Rev. Dr. R. J. Nevin.
41	1 1/2	23	E. Africa		Lord Delamere.
41	1 1/2	2	Matabeleland .		Major James Grant.
41	17	21	Sudan		Col. Ralph Vivian.
418	1 👸	28	Kaokoland		Capt. F. Cookson.
-4 1	•••	•••	,		C. T. Jones.
48	15	13	Zululand		LieutCol. Hon. W. Coke.
4	1 8	$2\frac{3}{4}$	Matabeleland .		Duke of Roxburghe.
-3 7	2	2	Komati Valley, S.A.		Count E. Hoyos.

¹ Killed with hounds.



Skull and Head of Male Klipspringer, from specimens shot in Somaliland by Major H. G. C. Swayne.

KLIPSPRINGER (Oreotragus saltator).

Alakud of the Somalis. Chibila in the Chilala and Chibisa Klipbok of the Boers. countries. Ingululu of the Makalakas.

Ikumi of the Basutos. Ligoka of the Zulus and Swazis. Njerere in the Batoka country.

Sass of the Abyssinians.

Not only from the other members of the oribi group, but from antelopes of all kinds, the agile little klipspringer, or "rock-jumper," is distinguished by the peculiar conformation of its hoofs and the structure The former are large, cylindrical, blunt, and so situated in of its hair. regard to the rest of the limb, that the animal walks on what corresponds to their tips in other antelopes, the whole hoof thus rising vertically from the ground. As regards the hair, this may best be compared with that of the musk-deer, having the same brittle, pithy structure. Lateral hoofs are retained; the tail is reduced to a mere rudimentary stump; and the horns of the bucks rise nearly vertically from the head, with a slight forward curvature, and are ringed for their The speckly olive-gray hue of the fur is too well basal third. known to need description. Height at shoulder from about 20 to 22 inches.

Distribution.—Mountainous and rocky districts in South and East Africa, from the Cape northwards to Abyssinia. Klipspringer-shooting is the best mountain sport to be obtained in Africa; and a pair of these active little animals bounding, as if made of indiarubber, from rock to rock is a sight never to be forgotten. To bag these exceedingly active and shy little antelopes, the sportsman, especially in the steep mountain ranges of Cape Colony, must, however, be prepared for a long and difficult stalk under a blazing sun. Except when they require its hair for stuffing saddles, the Boers leave the klipspringer alone; which is doubtless one reason that it is still comparatively abundant. Its venison is of excellent quality.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
58	218	27	Mashonaland	The late H. H. Eyre.
51	2 1 g	2§	Transvaal	F. Vaughan Kirby.
41/2	$2\frac{3}{16}$	2 11	Wittberg, Cape Colony	H. A. Bryden.
41/2	2	3 1	East Africa	E. Gedge.
-4½			Barberton	Dr. Percy Rendall.
48	13	21	Matabeleland	Hon. R. A. Ward.
48	2	3 1	S.E. Africa	F. C. Selous.
44	2	35	Near Lake Rudolph .	H. S. H. Cavendish.
-41	$2\frac{1}{2}$	3	Somaliland	C. V. A. Peel.
-41	3 1	$2\frac{1}{2}$	South Africa	A. Ohlsson.
-44	ış	21	Do	Mr. Justice Hopley.
-41	2	$2\frac{1}{2}$	Somaliland	Sir Edmund G. Loder, Bart.
41	$2\frac{1}{2}$	13	North Nyasaland .	James Yule.
4	11/2	2	East Africa	Lord Delamere.
4	17	2 1	?	Mr. Justice Hopley.
4	2 1 g	11	Matabeleland	W. W. Ashley.
4	$2\frac{1}{2}$	17/8	South Africa	H. and C. Beddington.
3 7	21/8	2	Do	A. Beit.
34	2	21	East Africa	J. Gardiner Muir.
3 3	13	28	Somaliland	Prince Boris Czetwertynski.
3 3	2	2	Do	T. W. H. Clarke.
3 8	•••	1 7	Abyssinia	British Museum.
-3 3	28	3 1	South Africa	Dr. W. P. Y. Bainbrigge.
-34	2	2	Do	Julius Jeppe.

KLIPSPRINGER (Oreotragus saltator)—continued.

Length on front.	Circum- ference.	Tip to Tip.	Locality.			Owner.		
-34	•••		Matabeleland			J. Brander Dunbar.		
31/2	$2\frac{1}{2}$	1 7	Somaliland			The late W. Babington.		
31/2	14	2	Matabeleland		•	Hon. John Ward.		
$3\frac{1}{2}$	2	23	Do			R. A. Cooper.		
- 9 3½		•••	East Africa			F. J. Jackson, C.B.		
38	1 7	2	Abyssinia .			Viscount Edmond de Poncins.		
31	1 7	2	South Africa			G. Richards.		
31			Somaliland		•	Major H. G. C. Swayne.		
-31	12	4	?			Dublin Museum.		
31 ¹ 6	17	2	Abyssinia .			Viscount Edmond de Poncins.		
3	1 7	23	Somaliland			G. D. E. Chapman.		
3	1 5	13	Do		•	W. W. Ashley.		
3			?			Cape Town Museum.		



Horns of Waterbuck, F. H. Barber's specimen.

COMMON WATERBUCK (Cobus ellipsiprymnus).

countries.

Kring-gaat of the Dutch. Li Tumogha of the Matabele.

Kooli in the Chilala and Chibisa M'dongoma or Matutwi in the Barotse country. Swahili name Koru. Tumoga of the Bechuanas.

The waterbucks and their near allies the kobs, together with the reedbucks and vaal rhebok, constitute a well-defined group of large or medium African antelopes presenting the following characteristics. They have the muzzle naked, no face-glands, a moderately long tail, well-developed lateral hoofs, and the horns confined to the bucks. In shape and size the horns are variable, being either long or medium, but never spirally twisted, and always with smooth tips, below which they are ridged; usually they are at first inclined somewhat backwards. after which they are curved upwards and more or less forwards, although they may have a sinuous curvature, and in the vaal rhebok are straight. The upper molar teeth are tall and narrow. In the waterbucks and kobs, which include the largest representatives of the group, there are no naked patches on the head below the ears, the tail is comparatively long, with a slight terminal tuft, and the lateral hoofs are large. A characteristic feature of the skull is the presence of a deep hollow in the forehead. From its allies the true or common waterbuck is recognisable at a glance by the elliptical white ring on the buttocks, which extends downwards to the thighs. Height at shoulder about 43 inches to 53.

Distribution.—Africa north of the Limpopo along the eastern coast region as far as the Shebeyli River in Somaliland; thus including Nyasaland and British and German East Africa. Never, apparently, very abundant, this handsome antelope, whose head and horns form one of the chief prizes of the South African hunter, has had its range much curtailed of late years. Its present strongholds are the unhealthy districts between the Sabi and Zambesi, the affluents of the latter river, and the Chobi, Okavango, and other rivers above Lake Ngami. Although the flesh is uneatable by Europeans, the excellence of the hide for shoe-leather causes the kring-gaat, as it is called by the Boers, to be persistently hunted.

			•			
Length on front.	Circum- ference.	Tip to Tip.	Locality.			Owner.
-36½			Delagoa Bay .			F. H. Barber. (See illustration.)
-36 1			South Africa .			O. R. Dunell.
-351	9	14	3			Mr. Justice Hopley.
-34½			Limpopo Valley			H. T. and A. H. Glynn.
$-33\frac{1}{2}$	10	21 3	S.E. Africa .			F. Vaughan Kirby.
$33\frac{1}{2}$	91	21½	Do			Hon. Walter Rothschild.
33	98	1112	${\bf Mashonal and} \ .$	•		F. C. Selous, British Museum.
33	83	24	Do			J. G. Millais.
-33	83	17‡	Do			A. Ohlsson.
-33			Lake Ngami.			Cape Town Museum.
32 3	9 3	21 3	South Africa .			Sir Victor Brooke's Collection.
313	9 1	24	Zululand .			Capt. L. O. Williams.
-31 8	81	21 7	North of Pungwe		•	Count E. Hoyos.
-311	9	17	?			Julius Jeppe.
318	81	21 7	?			British Museum.
311	101	138	Pungwe			Earl of Dunmore.

COMMON WATERBUCK (Cobus ellipsiprymnus)—continued.

Length on front.	Circum- ference.	Tip to Tip.	Locality.		Owner.
314	9₿	19‡	?		Sir Edmund G. Loder, Bart.
-314			Transvaal		C. T. Jones.
31			Mashonaland		F. C. Selous.
-31	9 \$	17	E. C. Africa	•	James J. Harrison.
30 3	8	231	Pungwe		Capt. Lord Douglas Compton.
30½	83	25	Do		R. Hughes.
-30½	91	161	E. C. Africa		James J. Harrison.
-30½	9	26	Do		Julius Jeppe.
30	9 1	$23\frac{1}{2}$	Zululand		A. J. Brandon.
-30l	•••	•••	Macloutsie		Capt. G. F. T. Leather.
30 1	81	137	Mashonaland		F. C. Selous.
-30	91	114	Komati River, S. Af	rica .	Count E. Hoyos.
298	9 1	127	Zululand		Hon. Charles Ellis.
29 3	9 1	21 7	Mashonaland		F. C. Selous.
-29 ‡	•••	•••	Somaliland	•	Dr. Donaldson Smith.
-2916	918	1911	Danakil		Prince A. de Lucinge.
29 8	98	•••	Zululand	•	A. H. Neumann.
29	91	5½ malformed	Do	•	A. W. Davis.
29	102	16	Sabi River	•	Sir Thomas Fowler, Bart.
29	91/2	26	East Africa		E. Gedge.
29	91	16 1	Do	•	Major W. H. Williams.
-29	81/2	19	Do	•	T. E. Buckley.
29	9	16	South Africa	•	LieutCol. Hon. W. Coke.
29	9	148	Pungwe	•	Capt. F. H. Lehmann.
29	9	123	Do		R. A. Cooper.
29	8 §	19‡	Do	•	Frank Harris.
-29	8	221	Do	•	C. C. Gouldsmith.
283	9 1	21	?		H.R.H. the Duke of Saxe-Coburg and Gotha.
28 3	8	27	Shiré Valley	•	H. H. Williams.
283	8	154	South Africa		G. H. M. Banks.

COMMON WATERBUCK (Cobus ellipsiprymnus)—continued.

Length on front.	Circum- ference.	Tip to Tip.	Locality.			Owner.
28½	8‡	151	Pungwe			Viscount Ennismore.
$-28\frac{1}{2}$	81		Shiré Valley .			Staff-Surgeon J. Dowson, R.N.
-28	103	8	S.E. Africa .			T. E. Buckley.
-28		18	East Africa .			Sir John Willoughby, Bart.
28	9	18	Matabeleland.			Major James Grant.
28	81	131/2	Pungwe	٠		Durban Museum.
28 1	8	$22\frac{1}{2}$	Do			H. R. Holden.
27 3	91	14 3	East Africa .	•		F. J. Jackson, C.B.
-27 3	83	101	Barberton .			Dr. Percy Rendall.
27 5	8	98	Zululand .			Major-Gen. Sir Arthur Ellis, K.C.V.O.
$27\frac{1}{2}$	83	22	South Africa .			D. Norman Ritchie.
$27\frac{1}{2}$	81	18	Pungwe	•		G. L. Bonham.
271	88	16 3	Barotseland .	••		H. Timmins.
271	9 1	13½	East Africa .			J. Gardiner Muir.
27	$9\frac{1}{2}$	13	Zululand .			A. Cameron.
27	8	15\$	Pungwe			Lord Edward Manners.
27	88	171	Danakil		•	Viscount Edmond de Poncins.
27	8	174	Pungwe	•	•	Ford G. Barclay.
27	7 1	$34\frac{1}{2}$	Zambesia .			Comdr. A. T. Hunt.
27	9	44	Matabeleland		•	W. Crosley.
26 <u>3</u>	9	$18\frac{1}{2}$	Pungwe	•		Capt. G. F. Henry.
26 2	81	14	Do	•	•	Col. G. A. Percy.
26 3	6 <u>1</u>	18‡	E. C. Africa .	•	•	Lord Delamere.
26½	8	114	Do		•	Lord Edward Manners.
-26½	81	151	Masailand .	٠		Count Scheibler.
-26	8	171	South Africa .	•	•	Dublin Museum.
-26	81	. 18	Do. .	•	•	James J. Harrison.
-26	71/2	201	Do		•	Dr. W. P. Y. Bainbrigge.
25 7	8 3	171	Sudan			Col. Ralph Vivian.
258	81	11	East Africa .		•	The late Capt. Speke, British Museum.

COMMON WATERBUCK (Cobus ellipsiprymnus)—continued.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.		
$25\frac{1}{2}$	81	19½	British Central Africa	J. E. Gough.		
25 8	7 🖁	18 1	}	Lord Delamere.		
251	84	13 7	East Africa	Sir Robert Harvey, Bart.		
25 1	8	13#	Do	H. C. V. Hunter.		
244	$7\frac{1}{2}$	9#	Somaliland	W. D. James.		
-23.6	8.13	10.1	British East Africa	Prince Henry of Liechtenstein.		
22 3	6	151	Somaliland	Major H. G. C. Swayne.		
221	$7\frac{1}{2}$	12 7	Do	A. H. Straker.		
22	7 1	9 1	Do	Prince Demeter Ghika.		
-21 <u>3</u>	7₺	9 7	Do	Count E. Hoyos.		



Head of Common Waterbuck, from a specimen shot in Somaliland.



Skull of Male Sing-sing Waterbuck, from a specimen in the British Museum.

SING-SING WATERBUCK (Cobus defassa).

Distinguished from the common water-buck (C. ellipsiprymnus) by the presence of a large white patch on the lower part of the buttocks, instead of a white elliptical ring extending higher up. Four more or less distinct local races of this species may be recognised. First, the West African sing-sing (C. defassa unctuosus), of Senegal and Gambia, characterised by its rufous colour and the small amount of white in the region of the eye. Second, Crawshay's sing-sing (C. defassa crawshayi), from British Central Africa, with a dusky coloration. Third, Penrice's sing-sing (C. defassa penricei), from the interior of Benguela, Angola, in which the colour is so dark as to be almost black.

And, fourth, the defassa sing-sing (C. defassa typicus), extending from Western Abyssinia through Sennar, Kordofan, and the valley of the White Nile to Uganda and British and German East Africa, in which the ears are longer and more pointed, and there is more white in the region of the eye than in the other races; the general colour being rufous. The four races are frequently regarded as distinct species, but they are so evidently local modifications of a single somewhat variable form that it appears far preferable to include them all under a single specific heading.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
1 -35½			Near Toru	Major C. G. H. Sitwell.
-29 3	8§	298	Sudan	Count T. Palffy.
-29	91	171	?	Dublin Museum.
-29	81	$28\frac{5}{18}$	Upper Basaland, Abyssinia	J. Menges.
-29	8	6 <u>1</u>	?	Julius Jeppe.
-27 7	81	9 7	Sudan	Count T. Palffy.
278	81	131	?	British Museum.
27 1	7 1	18 <u>‡</u>	Atbara Valley, Abyssinia .	Do.
-271	9	12	East Africa	Lord Delamere.
27 1	$7\frac{1}{2}$	18 1	Atbara Valley, Abyssinia .	British Museum.
27	8	154	Kavalli, Lake Albert	Col. Trevor Ternan.
26 1	74	15 1	Uganda	The late Captain Speke, British Museum.
26 3	8§	151	East Africa	E. Gedge.
-26	78	133	Borders of Lake Tchad .	Paris Museum.
25 3	9	153	Bahr-el-Salam, Sudan .	LieutCol. Hon. W. Coke.
251	81	16 1	Kikuyu	E. Russell.
2518	85	138	British East Africa	Hon. Walter Rothschild.
241	83	16 1	Molo River	G. E. Smith.
23½	8	83	?	Lord Delamere.

¹ Mr. F. R. Hicks favours me with this measurement.

DEFASSA SING-SING (Cobus defassa unctuosus).

See page 178.

Height at shoulder, 47 to 48 ins.—Major A. J. Arnold, D.S.O.

Length on front.	Circum- ference.	Tip to Tip.		Loca	lity.		Owner.
244	81/2		Gambia				Julius Jeppe.
$-24\frac{1}{2}$	9	12	Nigeria		•		Dr. Percy Rendall.
24	7‡	154	Do.				P. A. Clive.
23 3	71	141	Do.				F. Swanzy.
23 3	71/2	131	Do.				Major A. J. Arnold, D.S.O.
23	7	16	Do.				Do.
23	7	134	Boussa,	Niger	ia		A. W. M. Brodie.
22 ⁷ 8	81	103	Nigeria				Sir Edmund G. Loder, Bart.
22	7	171	Do.				Major G. S. C. Jenkinson.

PENRICE'S SING-SING (Cobus defassa penricei).

See page 178.

Length on front.	Circum- ference.	Tip to Tip.		Locality	.		Owner.
29	83	18 <u>1</u>	Benguela				G. W. Penrice.
-28 1	81	1112	Do.				A. Ohlsson.
28	8	$22\frac{1}{2}$	Do.				G. W. Penrice.
$24\frac{1}{2}$	8	91	Do.				Hon. Walter Rothschild.
201	6 7	12‡	Do.				Julius Jeppe.



Horns of Mrs. Gray's Waterbuck, from Lieut.-Col. Hon. W. Coke's specimen.

MRS. GRAY'S WATERBUCK (Cobus maria).

This species is the last of the typical group of *Cobus*, in all the members of which the size is large, the fur grizzled, and the neck maned. From the preceding species Mrs. Gray's waterbuck differs by the presence of an hour-glass-shaped white area on the back of the neck, and the form of the horns, which are highly sinuous and inclined backwards; the general colour of the fur being dark reddish brown. Height at shoulder apparently from 35 to 40 inches.

Distribution.—The swamps bordering the White Nile and its tributaries. Very rare in collections, being represented by complete specimens only in the museums of Vienna and Berlin. It was discovered by Heuglin, but has been confounded with the very different white-eared kob.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-30 1	6 1	17	?	Berlin Museum.
	6 <u>‡</u>	12	?	Hon. Walter Rothschild.
29 ¹ / ₂ 26 ⁷ / ₈	$6\frac{7}{8}$	133	Arwan, Bahr-el-Ghazal	The late Consul J. Petherick, Brit. Mus.
26 1	61	12	White Nile.	LieutCol. Hon. W. Coke.

WHITE-EARED KOB (Cobus leucotis).

The smaller antelopes included in the genus *Cobus* may be popularly termed kobs, and differ from the water-bucks not only by their inferior size, but by the uniformly rufous hue of the upper parts, and the absence of a mane on the neck. From its allies the present species is sufficiently distinguished by the white outer surface of the ears and upper part of the head, including the region of the eyes. Height at shoulder about 34 or 35 inches.

Distribution.—The region of the Upper Nile, including the Sobat, Bahr-el-Ghazal, and their tributaries, and extending to the Niam-Niam country. The first known example was sent to Berlin by the German artist and traveller Werne, who obtained it in Sennar.

Length on front.	Circum- ference.	Tip to Tip.	Locality.		Owner.		
-21 5	7 8	13 18	White Nile			Paris Museum.	
-20	7	71/2	Do.			Berlin Museum.	
197	6 8	71	Bahr-el-Ghazal			British Museum.	
19 §	5₹	91	Do.			The late Consul J. Petherick, British Museum.	
174	6	71	. ?			Sir Victor Brooke's Collection.	



Frontlet and Horns of Buffon's Kob.

BUFFON'S KOB (Cobus cob).

In common with the Uganda kob and the puku (C. vardoni), this species has the back of the ears rufous like the body; and it agrees with the latter of these in having the horns less than twice the length of the head, the hair short, and the front of the fore-legs black; its peculiar distinctive features being the presence of a white line over each eye and its comparatively small size, the approximate height at the shoulder being from 32 to 33 inches.

Although this species was known to the French naturalist Buffon, it is only of late years that its affinities have been fully worked out. The puku differs from both this species and the under-mentioned sunu by its much longer hair and the absence of the black down the front of the legs.

Distribution.—West Africa, from the Gambia to Nigeria.

Length on front.	Circum- ference.	Tip to Tip.	L	ocal	ity.	Owner.
192	6 7	71	Nigeria		•	Captain A. H. Festing.
188	5 1	71/2	Do.			Captain J. S. Brogden.
-171	•••	•••	Do.			Major A. F. Mockler-Ferryman.
16 3	6	7	Do.			Major A. J. Arnold, D.S.O.

BUFFON'S KOB (Cobus cob)—continued.

Length on front.	Circum- terence.	Tip to Tip.	Locality.		Owner.	
163	5½	8	Ibi District .		Carl Jeppe.	
161	6	78	Do		Mr. Justice Hopley.	
16	53	74	?		Sir Edmund G. Loder, Bart.	
15‡	5ē	7	Ibi District .		Julius Jeppe.	
15‡	6‡	8	Gando, Western Sudan		J. W. Carroll.	
131/2	6	7‡	Gaboon		Hon. Walter Rothschild.	
91/2	5	3	Gambia		A former Earl of Derby, British Museum.	
8	48	44	Do		Do.	



Head of Lichi.



Skull of Uganda Kob, from a specimen shot by F. J. Jackson.

UGANDA KOB (Cobus thomasi).

The East African representative of Buffon's kob, from which it is distinguished mainly by its superior size, and the presence of a complete white ring round each eye, instead of having only a white line above the same. General colour rich fulvous, with the muzzle, lips, chin, under parts, and inner surfaces of upper portion of fore-legs and thighs white; front of fore-legs with a black line, and hind-legs with a similar line, which does not, however, ascend within some distance of the hocks. Height at shoulder about $35\frac{1}{2}$ inches.

Distribution.—East Africa, from Kavirondo to Uganda.

Native name, Sunu or N'Sunu.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
201	7	94	North end of Lake Albert	Col. Trevor Ternan.
193	7	97	Victoria Nyanza .	E. Gedge.
18 <u>1</u>	6₽	97 8 5	Lake Albert	Col. F. D. Lugard, C.B., D.S.O.
18 1		•••	Lake Victoria	Major W. H. Williams.
18	7 1	I I 🛓	Uganda	E. Gedge.
171	6 <u>\$</u>	ΙΙÃ	Do	M. F. Ğage.
17	6 <u>}</u>	74	Kavirondo	F. J. Jackson, C.B.
171	7⅓	5 <u>₹</u>	Do	Rowland Ward.
17 Å	7	5	Do	F. J. Jackson, C.B., British Museum.
-17	518	9개 6월	East Africa	Paris Museum.
16 7	6 1	6 <u>1</u>	Uganda	Sir Edmund G. Loder, Bart.
16	6₿	10	East Africa	F. H. Barber.
15‡ 14‡	5 ²	10	Do	Julius Jeppe.
143	6	9₹	Victoria Nyanza .	C. F. S. Vandeleur, D.S.O.



F. H. Barber's Lichi Horns.

LICHI or LECHWE (Cobus lichi).

Lechwi in the Barotse and Lake Leché, Lee-gwee of the Makololo.

Ngami countries.

Njha in the Chilala and Chibisa

countries.

Leché, Lee-gwee of the Makololo.

Inya of the Masubias.

Oonya of the Makubas.

In this handsome antelope the horns are longer than in its allies, considerably exceeding twice the length of the head, while the front of the fore-legs is black, and the hair long. Height at shoulder, 40 or 41 inches.

Like the puku, this kob was discovered by Livingstone and his companions Oswell and Murray during their journey to Lake Ngami in 1849. In point of size it approximates to the true waterbucks, from which, however, in addition to the features already mentioned, it may always be distinguished by its lighter and more graceful build. The general colour is fulvous, of a lighter tint than in the Uganda kob.

The lichi affords excellent and often exciting sport for the gunner. Seldom found very far from water, this antelope is most abundant in the lagoons and swamps created by the annual rising of the Upper Zambesi, the Botletli, Chobe, Tamulakan, and other rivers of the interior, in localities where the surrounding flats are inundated for some part of the year. The lichi seldom ventures into the deep rivers from fear of the crocodiles, but among the reed beds, the shallow lagoons, and flooded flats it is a familiar figure. It is often to be found, in the less accessible regions, in vast herds. An excellent swimmer, it can progress by a succession of splashing bounds at great speed through the lagoons and shallows. It is extremely tenacious of life, and requires very straight powder. The thick rufous-yellow coat of the lichi is extremely handsome, and the skin is greatly prized by the natives.

Distribution.—Zambesia, reaching northwards to Lake Mweru, and to Lake Ngami towards the south-west.

Length on front.	Circum- ference.	Tip to Tip.	Locality.		Owner.
$-33\frac{1}{2}$	7	26 3	Lake Bangweolo		Poulett-Weatherley.
-28 1			Zambesia		F. H. Barber. (See illustration.)
-28	•••	•••	?		T. Poole.
27\$	8	14	Okavango Valley	•	Major H. J. Goold-Adams, C.B., C.M.G.
$27\frac{1}{2}$	102	191	Chobe Valley .		F. C. Selous, British Museum.
271	8	143	Do		F. C. Selous.
-27	71	14	Okavango Valley		Sir Edmund G. Loder, Bart.
-26 1	9	201	?		Julius Jeppe.
26 1	88	21	South Africa .		A. Beit.
-26 1	6 §	16	?		Mr. Justice Hopley.
26 1	8	15 (about)	Linyanti, Chobe Vall	ley	F. C. Selous.
25 3	7 1	123	?		G. Richards.
-253	7 ½	18	Zambesia		T. E. Buckley.
- 2 5∯	71/2	13 8	Do		Dr. Holub, Paris Museum.
-25½	74	16 1	3		Julius Jeppe.
-258	7 1	141	Lake Ngami .		Dr. Percy Rendall.
25 1	7 3	111	?		Mr. Justice Hopley.
24 1	7	19‡	?		R. A. Cooper.
24½	71/2	I I ½	South Africa .		Hon. Walter Rothschild.
248	7∰	13€	Zambesia		The late J. S. Jameson.
-244		•••	Do		C. T. Jones.
24	71	121	?		A. Ohlsson.
231	$7\frac{1}{2}$	141	Chobe Valley .		F. C. Selous.
-23			Do		O. R. Dunell.
23	9	9	Do.		E. Gedge.
228	6 }	123	Barotseland .		R. T. Coryndon.



Head of Puku, from a specimen in the British Museum.

PUKU (Cobus vardoni).

Impookoo of the Masubias.

Muntinya in the Barotse country.

Pookoo in the Lake Ngami country. Seūla in the Chilala and Chibisa countries.

From the other small kobs with the back of the ears rufous, the puku is distinguishable at once by the uniformly foxy colour of the fore-legs, as well as by the greater length of the hair, especially in the region of the back and loins, where it has a tendency to curl. General colour reddish-yellow. Height at shoulder about 39 or 40 inches.

Distribution.—Mainly confined to a narrow belt of country extending along the southern bank of the Chobe for about sixty miles from its junction with the Zambesi. The herds are small, seldom including more than ten or a dozen individuals.

PUKU189

PUKU (Cobus vardoni)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
20 1	81/2	12‡	Luswesi Valley	F. Smitheman.
191	6 3	81	South Africa	J. Carr Saunders.
-185	•••		Njoko Valley	Major A. St. H. Gibbons.
183	6 3	9 1	Barotseland	R. T. Coryndon.
-18	6 1 ⁄2	9	Luwulé Valley, Congo	Poulett-Weatherley.
τ8	6‡	83	Free State Luapulu Valley	F. Smitheman.
171	6 <u>1</u>	7	Bangweolo	Do.
171	6 <u>3</u>	51/2	Barotseland	R. T. Coryndon.
-17½	7	6₫	Choma Valley, B.C.A.	R. H. Ferrers Stranack.
17 #	61	18 malformed	Kabampo Valley .	Major H. J. Goold-Adams, C.B., C.M.G.
163	6 <u>1</u>	83	Do	Do.
-164	6 1	71/2	Luwulé, B.C.A	Poulett-Weatherley.
-16 1	7	6 <u>1</u>	Choma Valley, B.C.A.	Julius Jeppe.
16 1	6월	7 ½	South Africa	British Museum.
16	6 <u>1</u>	6 <u>3</u>	South bank of Chobe Valley	F. C. Selous.
16 8	6≩	57	Choma Valley, B.C.A.	Mr. Justice Hopley.
16	5⅓	41	West Nyasaland	C. C. Bowring.
-16	6 <u>1</u>	8	?	A. Ohlsson.
1-15 4	• • •	•••	Njoko Valley	R. T. Coryndon.
15 8	6 1	51	British Central Africa	Rowland Ward.
15 8	6 1 ⁄2	6 8	Chobe Valley	F. C. Selous.
15 1	6	10	South Africa	J. Carr Saunders.
2-15 8	•••		Njoko Valley	R. T. Coryndon.
- 15	6‡	6 <u>‡</u>	Zambesia	T. E. Buckley.
³ -15			Njoko Valley	R. T. Coryndon.
148	61	71/2	Lofu Valley, B.C.A	John Yule.
141	6‡	$3\frac{1}{2}$	British Central Africa	Julius Jeppe.
14 1	6 8	51	Chobe Valley	F. C. Selous, British Museum.
1 Weigh	nt as he fell,	177 lbs.	² Weight as he fell, 185 lbs.	³ Weight as he fell, 196 lbs.



Head of Vaal Rhebok.

GRAY or VAAL RHEBOK (Pelea capreolus).

Iliza of the Swazis.

Pshiatla of the Basutos.

Peeli of the Bechuanas.

The short, upright, straight, and slender horns, together with the somewhat woolly nature of the hair, serve to differentiate the vaal rhebok from the antelopes of the kindred genera. The tail, like that of the reedbucks, is short and bushy; the ears are tall and narrow; the build is slight and graceful; and the general colour is uniformly pale gray, tending somewhat to fawn on the head and limbs. Height at shoulder about 29 or 30 inches. The chief essential distinctions between this antelope and the reedbucks are the form of the horns and the absence of the bare patches below the eyes.

Distribution.—The open hilly districts of Africa south of the Zambesi. In the mountain ranges of the eastern and northern districts of Cape Colony, and thence onwards to the Zambesi, the vaal rhebok affords good and exciting sport. And although its somewhat stiff

and stilty appearance is not at first suggestive of unusual activity, yet the pace and regularity with which a party of six or eight of these antelopes will race up the steep flank of a mountain are calculated to remove any doubts on this point. The most peculiar feature of the vaal rhebok is the soft and woolly nature of the gray coat; the horns afford but insignificant trophies, and the flesh is of decidedly poor quality for the table.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-1112	•••	•••	Spitzkop .	F. H. Barber.
-11½	•••	•••	Cape Colony	Cape Town Museum.
-11	•••	•••	?	M. E. Bowker.
10 3	2	$2\frac{1}{2}$	Basutoland .	Julius Jeppe.
$-10\frac{1}{2}$	2 1	45	Transvaal .	H. T. and A. H. Glynn.
-101/2	•••		Cape Colony	Cape Town Museum.
-9 1	2 8	28	?	Julius Jeppe.
-91	•••	•••	Transvaal .	C. T. Jones.
-9 ₹	2	41	Basutoland .	Sir A. Milner.
91	21/2	24	?	Dr. Oakshott.
9	25	2	3	Mr. Justice Hopley.
-815	28	Single horn	Barberton .	Dr. Percy Rendall.
-8 3	25	3₽	?	A. Ohlsson.
88	$2\frac{1}{2}$	23	Cape Colony	R. H. Venables-Kyrke.
8§	$2\frac{1}{2}$	25	South Africa	The late Dr. Burchell, British Museum.
81	$2\frac{1}{2}$	2	?.	G. Richards.
88	2	3₺	?	Sir Edmund G. Loder, Bart.
81	2	27	5	F. C. Selous.
-81	218	34	?	Dr. W. P. Y. Bainbrigge.
-8	2	3	Mauchberg .	F. Vaughan Kirby.
7 3	•••	•••	?	O. R. Dunell.
72	$2\frac{1}{2}$	34	?	Hon. Walter Rothschild.
7 1	28	3	3	H. Atkinson.
7‡	21	$3\frac{1}{2}$	Basutoland .	A. Beit.
7	2	$2\frac{1}{2}$? `	Capt. H. A. Livingstone.



Head of Common Reedbuck.

COMMON REEDBUCK (Cervicapra arundinum).

Bemba of the Masaras.

Inzeegee of the Amandebele.

Im-vwee of the Masubias.

Um-vwee of the Makubas.

Inhlango of the Swazis.

Iklabu of the Basutos.

Impoyo of the Lower Zambesi tribes.

Mziki of the Zulus and Matabele. Natafwi in the Mashukulumbwi country.

Mutobo in the Barotse country.

Sibughat in the Lake Ngami country.

Mpoyo in the Chilala and Chibisa country.

All the reedbucks, as the members of the genus Cervicapra may be collectively termed, differ from the waterbucks and kobs by their inferior

size, lighter build, and the presence of a completely bare or very short-haired patch on each side of the head immediately beneath the ear. The tail, too, is more bushy and shorter, the lateral hoofs are relatively smaller, and the horns, which are of medium length and stoutness, curve regularly upwards, and in some cases also forwards. The present species, the true reitbok of the Boers, is the largest of the genus, standing about 36 or 37 inches at the shoulder; and has the ear-patch completely bare, and the horns inclining markedly forwards. Length from nose to tail 37 inches in a specimen shot by Dr. Percy Rendall.

Distribution.—South Africa, extending as far north as Angola on the west, and to Mozambique on the east coast. Although formerly numerous in the eastern districts of Cape Colony and Natal, reedbuck have now become scarce in South Africa, and it is not till Khama's country is reached that they are met with in any numbers. As their name implies, reedbuck frequent the reed-brakes fringing so many African rivers; and some of the rivers where they are still fairly abundant are the Lotsani in Bamangwato and those of Ngamiland and the country between Mashonaland and the east coast. They are generally met with in small family parties; and when excited or alarmed utter a characteristic shrill whistle. As their flesh is of fair quality, and their shooting by no means difficult, they are much sought after by sportsmen.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
1-18	•••	•••	S. Africa	F. Vaughan Kirby.
$-17\frac{1}{2}$	6 1	171	Barotseland	Percy C. Reid.
-16 8	63	191	Mpimbi, B.C.A	Major F. Trollope.
-16	5½	1112	S. Africa	A. Ohlsson.
15 7	6‡	101	Do	C. D. Rudd.
15 7	6∤	141	Do	British Museum.
-15 7	6 7	10 1	Zululand	F. R. N. Findlay.
² 15 ³	6 1	$13\frac{1}{2}$	Transvaal	Julius Jeppe.
154	6 1	•••	South Africa	The late J. S. Jameson.
$-15\frac{1}{2}$		•••	Nyoko Valley, B.C.A	Major A. St. H. Gibbons.
-15½	6 <u>₹</u>	151	Transvaal	F. Vaughan Kirby.

This was seen and measured by Mr. Kirby. It had one horn only; the other was shot off.
2 Abnormal head.

COMMON REEDBUCK (Cervicapra arundinum)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-151		16	?	Lord David Kennedy.
-15½	•••	•••	Damaraland	. Cape Town Museum.
-151	•••		?	F. H. Barber.
15 1	6	111	Manica Plateau, N. o Zambesi	of F. C. Selous.
15	6 1	9 \$	Mashonaland	. Sir John Willoughby, Bart.
15	6	$12\frac{1}{2}$	Barotseland	. H. Timmins.
-15	•••	•••	?	H. T. and A. H. Glynn.
-15	6 <u>1</u>	98	?	T. E. Buckley.
-15	•••		?	O. R. Dunell.
144	6	15 1	Zomba, B.C.A	. C. C. Bowring.
-14 8	6 <u>8</u>	101	Sabi Flats	. Dr. Percy Rendall.
148	7 8	9 1	?	Hon. Walter Rothschild.
141	6 8	7 8	?	Sir Edmund G. Loder, Bart.
141	61	128	Mashonaland	. A. Eyre.
148	5½	16 <u>8</u>	S. Africa	. LieutCol. Hon. W. Coke.
141	5 1	111	British Central Africa	. J. E. Gough.
-142	•••		Transvaal	. C. T. Jones.
14	6 <u>1</u>	111	Barotseland	. R. T. Coryndon.
13 8	54	114	S. Africa	. W. A. Edmonds.
134	6	10 3	Do	. R. A. Cooper.
138	6	82	Pungwe	. H. R. Holden.
13 8	6	12	Northern Rhodesia .	. F. Smitheman.
131	6	91	?	Hon. Walter Rothschild.
131	61	6	Pungwe	. Viscount Ennismore.
131	6 <u>3</u>	6 7	S. Africa	. Major-General Sir F. C. Carrington, K.C.B., K.C.M.G.
$-13\frac{1}{2}$	5 2	$12\frac{1}{2}$	Matabeleland	J. Brander Dunbar.
131	7 1	111	Zululand	. A. Cameron.
131	61	91	S. Africa	. G. Richards.
-13	7	7 ½	Do	. James J. Harrison.

COMMON REEDBUCK (Cervicapra arundinum)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
-13	61	108	?			Mr. Justice Hopley.
12	5₩	8	Zululand			Capt. L. O. Williams.
124	58	81/2	Busi Valley .			E. S. Grogan.
-I2 <mark>3</mark>	42	11	Nyasaland .	•		Major P. W. Forbes.
122	6 1	8 8	Near Bubye River		•	A. M. Sagar-Musgrave.
124	5‡	91	Zambesia	•	•	Comdr. A. T. Hunt.
$12\frac{1}{2}$	6	6 <u>1</u>	S. Africa	•	•	F. E. Potter.
121/2	6	98	Pungwe	•	•	Capt. G. F. Henry.
121/2	54	134	Do			R. K. Micklethwait.
12	6 <u>8</u>	82	Do	•		Capt. F. H. Lehmann.
12	54	9 8	Do		•	Hon. T. Thynne.
12	6	7	Do	•		Col. G. A. Percy.
12	5₩	91	?			Julius Jeppe.
12	5 1	103	S. Africa		•	Montrose Cloete.
112	5₽	101	Benguela			G. W. Penrice.
112	5	9	Pungwe	•	•	C. C. Gouldsmith.
1112	6	13	Do		•	Count E. Hoyos.
114	7	91	Do		•	P. B. Vander-Byl.
-111	6 3	91	?			Dr. W. P. Y. Bainbrigge.
11	61	111	Zululand	•		A. J. Brandon.
10 7	6	81	Pungwe			Lord Edward Manners.
101	5	6	Do			G. L. Bonham.
101	48	6	Do			G. Micklethwait.

BOHOR REEDBUCK (Cervicapra bohor).

Boroufa of the Gallas. Porhé of the Swahilis.

From the common reedbuck (C. arundinum) the present species is chiefly distinguished by its smaller size (height at shoulder about 31 inches), while it is considerably larger than either of the other three members of its genus. Young animals have the horns more hooked at their tips than is the case with the common reedbuck, but with advancing age even this point of distinction becomes little apparent, owing to the wearing away of the tips of the horns. Both the head and body are more uniformly fawn-coloured than in the common species.

Viscount Edmond de Poncins says:—"These antelopes are very numerous in the Galla country near Mount Yokoila; they like open grassy plains more or less dotted with the small mimosa bushes and are found in small herds of from four to eight, sometimes even fifteen or twenty. They are not very wild and may easily be stalked at less than 200 yards in the middle of the day. Old males frequently are found in the long grass quite alone, when if disturbed they gallop through the grass, jumping very high; on the plains they go easy and fast without jumping. The flesh is not bad eating, and the Gallas are very keen about getting the skins. Weight about 80 lbs."

A female shot by him measured from nose to root of tail 51½ inches; height at shoulder, 31½ inches; round the body, 30 inches.

A good male shot in the Galla country south of the Shoa, by the same sportsman, measured from nose to root of tail 57½ inches; tail, 6½ inches; height at shoulder, 34½ inches; round the body, 34½. Distribution.—From Abyssinia through East Africa as far south as Kilimanjaro. Discovered by Rüppell in Abyssinia.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.	
132	5 1	8 1	East Africa .			British Museum.	
131/2	51/2	41/2	Do	•		Major J. R. Macdonald.	
$12\frac{1}{2}$	5∰	5 8	Do			Sir John Kirk, K.C.B., British Museum.	
10§	41/2	51/2	Do			Sir Robert Harvey, Bart.	
-9 1	47	3 \$	Galla country.			Viscount Edmond de Poncins.	
-91^{4}	51/2	61	Do			Do.	
9 1	48	5 1	Usongola .			Col. F. D. Lugard, C.B., D.S.O.	

BOHOR REEDBUCK (Cervicapra bohor)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
91	58	. 8 <u>1</u>	Sudan		The late Dr. Burchell, British Museum.
91	41	4₽	Masailand		
-91 ¹ 6	5 1	5 7	Galla country		Prince de Lucinge.
9	5‡	318	Zanzibar		Sir John Kirk, K.C.B.
9	5	34	Ganda, Nigeria .		J. W. Carroll.
-9	6	5	Shoa, Abyssina .		Prince de Lucinge.
88	5	38	East Africa		Sir Robert Harvey, Bart.
83	47	34	Masailand	•	H. C. V. Hunter.
81	57	5	East Africa	•	R. P. Carroll.
81	$4\frac{1}{2}$	51	Do		Sir Edmund G. Loder, Bart.
8 1	44	4월	Unyoro		Col. Trevor Ternan.
81	44	$2\frac{1}{2}$	Masailand		H. C. V. Hunter, British Museum.
7 3	4 🖁	23	East Africa		F. J. Jackson, C.B.
-7 3	5 1	6	Do		Julius Jeppe.



Head of Mountain Reedbuck.

MOUNTAIN REEDBUCK or ROOI RHEBOK (Cervicapra fulvorufula).

Inhlang-amatshe of the Swazis and Zulus.

This and the nagor are the two smallest representatives of the reedbucks, both measuring about 28 inches at the shoulder. The present one is specially distinguished by the absence of a distinct forward hook to the tips of the horns, and the general grayish fawn colour of the fur, which has, however, in some cases a more or less distinctly rufous tinge. Distribution.—Eastern Africa to the south of the Zambesi, particularly

Natal, Zululand, and Bechuanaland; represented farther north by the under-mentioned form. The rooi (red) rhebok differs from the typical representative of its genus in being an inhabitant of the basal slopes of mountains, at a lower level than the tract frequented by the vaal rhebok. They are generally found in small troops of a dozen head or less; and from their active habits and the difficult nature of the ground they frequent, afford excellent stalking. Although rare in most of the Transvaal, good bags of this antelope have been made in the western districts of that country. In the neighbour-

hood of the tributaries of the Limpopo, on the Mabube, Tamulikan, Machabe, Sunta, and Chobe rivers, as well as the tributaries of the Zambesi east of the Victoria Falls, and throughout Mashonaland and Matabeleland, rooi rhebok are still very abundant. Those reported from the Manica plateau, north of the Zambesi, may be the so-called Chanler's reedbuck.

			· ·	
Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-81	34	41	Transvaal	F. R. N. Findlay.
74	4‡	4 1	S. Africa	Hon. Walter Rothschild. (Shot by F. C. Selous.)
-7 1	48	4	Transvaal	F. Vaughan Kirby.
-7 1	4	31/2	Lebombo Range, S.A.	Count E. Hoyos.
-7 1			5	H. T. and A. H. Glynn.
78	4‡	3	Zululand	A. H. Neumann.
-71			3	O. R. Dunell.
-7‡	41	$2\frac{1}{2}$?	Mr. Justice Hopley.
-7 1	41	53	?	Julius Jeppe.
7	31/2	41	Zululand	A. Cameron.
6 3	4 1	43	3	Bloemfontein Museum.
6 2	48	4 1	?	F. E. Potter.
-6 §	3 1	38	?	Mr. Justice Hopley.
61	4	31	Zululand	LieutCol. Hon. W. Coke.
$-6\frac{1}{2}$	3 4	44	Barberton	Dr. Percy Rendall.
-6½	4	31/2	?	Dr. W. P. Y. Bainbrigge.
6 8	4‡	37	Zululand	A. J. Brandon.
-61	•••		?	C. T. Jones.
-64	41/2	4	?	A. Ohlsson.
-6	37	27	Swaziland	T. E. Buckley.
6	4	38	Zululand	Captain L. O. Williams.

LYDENBURG REEDBUCK (Cervicapra fulvorufula subalpina).

The reedbuck discovered by Mr. Vaughan Kirby in the Lydenburg range of the Transvaal departs from the ordinary type of the mountain reedbuck in exhibiting a tendency to partial albinism. Whether this peculiarity entitles it to be regarded as a distinct race must, for the present, remain a moot point.



Head of Chanler's Reedbuck. From a specimen shot by W. Astor Chanler.

CHANLER'S REEDBUCK (Cervicapra fulvorufula chanleri).

A reedbuck from East Africa described by the Hon. Walter Rothschild from a single skin, as a distinct species, appears to be nothing more than a local race of the rooi rhebok, from which it is even doubtful whether it should be separated as a sub-species. In the type specimen the nose shows a dark streak like the one often seen in the rooi rhebok and common reedbuck. Other skins have been obtained by Mr. F. J. Jackson, C.B.

Distribution.—The mountainous districts of British East Africa to the east of Mount Kenia and adjacent regions. By Mr. H. S. H. Cavendish these antelopes were seen in the Boran country.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
5₩	4	4	?	Lord Delamere.
5∰	31/2	48	?	Major A. E. Smith.
54	3½	37	Zombènè Range	W. Astor Chanler.

NAGOR REEDBUCK (Cervicapra redunca).

Distinguished from the mountain reedbuck (with which it agrees approximately in size) by the distinct forward curvature of the tips of the horns, so as to form a hook. General colour uniformly bright fawn, without darker markings on the limbs; tail only slightly bushy, fawn-coloured above and white beneath.

Distribution.—Senegal and Gambia.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
91/2	5	31 .	Ganda, Nigeria .	J. W. Carroll.
8	5 1	$3\frac{1}{2}$	Ibi, Benue River, West Africa	Julius Jeppe.
8	6	73	West Africa .	Hon. Walter Rothschild.
8	5	58	Gambia	Guy H. Sangster.
71	5 1	2	?	Sir Edmund G. Loder, Bart.
6 8	4 3	2 1 B	Ibi	Capt. A. H. Festing.



Skull and Horns of Blackbuck. From Mr. A. O. Hume's specimen.

BLACKBUCK or INDIAN ANTELOPE (Antilope bezoartica).

Although in former days almost or quite all of the then known species of antelope was included in the genus *Antilope*, the extent of the latter has been gradually whittled down until it comprises the Indian blackbuck alone. In addition to being the sole representative of the genus, this animal is also the type of a sub-family or group of antelopes, embracing among others the saiga, the gazelles, the impala,

and springbuck. All these are small or medium-sized antelopes, with hairy muzzles, generally short tails, and tall, narrow-crowned cheekteeth, like those of sheep. With the exception of the springbuck and the majority of the gazelles, horns are confined to the males. From all the other members of this assemblage the blackbuck is distinguished by the beautiful spiral formed by its horns. It has large face-glands, a short and compressed tail, and well-developed lateral hoofs. Height at shoulder about 32 inches; average weight, 85 lbs. It is only in adult males that the characteristic dark tint from which the species takes its name is developed; young males being uniformly brownish fawn above like the does.

Distribution.—India, from the foot of the Himalaya to Cape Comorin, and from the Punjab to Lower Assam; unknown in Ceylon and the countries to the eastward of the Bay of Bengal.

Length straight.	Circum- ference.	Tip to Tip.	Locality.	Owner.
1-28 <u>3</u>			Jeypore	Major-Gen. Sir B. Blood, K.C.B.
28 1	5	172	Near Delhi	A. O. Hume, C.B. (See illustration.)
-28		20	Do	Major P. Dunell Pank.
-28	5	15	38 miles north of Ahmedabad	T. Le Mesurier.
-27 <mark>7</mark>		•••	Jeypore	Major-Gen. Sir B. Blood, K.C.B.
-27	5	191	Bikanir	Capt. Harry V. Brooke,
26 3	42	251	Bhurtpore, N.W.P.	Col. E. T. H. Hutton.
-26 8	5	191	Bikanir	Capt. Harry V. Brooke.
-26 3	44		Do	Major-Gen. Alexander A. A. Kin-
-26 1	•••		N.W.P	loch. C. B. Oldfield.
26 8	5	177	Sirsa, Punjab	A. O. Hume, C.B.
-26 ₁₈	54	22	Bikanir	LieutCol. G. D. F. Sulivan.
26]	5	23 1	Oudh	E. St. J. Lawson.
26	5	21 7	Bikanir	Capt. H. W. Codrington.
-26			Jeypore	Capt. G. L. Holdsworth.
25 1	•••	15	?	Col. Martin.
$25\frac{1}{2}$	47	18	Punjab	Major R. H. Rattray.
25 8	51/2	141	Sirsa, Punjab	Hume Collection, British Museum.

¹ Measured and recorded by A. O. Hume, but not now in the possession of General Sir B. Blood, K.C.B.

BLACKBUCK or INDIAN ANTELOPE (Antilope bezoartica)—continued.

Length straight.	Circum- ference.	Tip to Tip.	Locality.		Owner.
-25 8	5	19	?		Count E. Hoyos.
25 1	44	19	Jeypore		A. B. Graves.
25 1	5 1	15	Kathiawar .		LieutCol. L. L. Fenton.
25	48	19	Sirsa, Punjab .		Hume Collection, British Museum.
-25	5‡	131	?		Bombay Natural History Society's Museum.
-25	•••		?		Lucknow Museum.
24 2	5	20	Jeypore		Surgeon-Major J. B. Buchanan.
243	$4\frac{1}{2}$	171	Dholpur	•	A. J. Coppinger.
241	58	201	Bikanir		H.H. Maharaja of Bikanir.
24 8	48	184	Do		LieutCol. G. D. F. Sulivan.
241	4 3	19½	?		A. Leslie Renton.
24 <u>‡</u>	5	$18\frac{1}{2}$?		Marquis of Ailsa.
-241	4-33	20.85	Udepur		Viscount Edmond de Poncins.
-24‡	5	194	Punjab	•	Capt. A. Hicks-Beach.
24	5	18 7	Aligahr .		St. George Littledale.
24	48	163	Rajpura		The late J. E. Ubsdell.
-24	48		Punjab		J. Johnston-Stewart.
24	•••	20	?		Duke of Bedford.
-24			?		Otho Shaw.
234	5 8	21	?		H.R.H. the Duke of Saxe-Coburg and Gotha.
23 8	44	134	?		Capt. F. C. Quicke.
-23 8	•••	•••	Central Provinces		Capt. H. M. Biddulph.
23 \$	5	16 1	Punjab		Major J. W. M. Cotton.
23 §	4 7	19‡	Patiala	•	LieutCol. E. E. Carr.
23 §	5 l	143	?		Hon. Walter Rothschild.
23 §	43	16 8	Central Provinces	•	C. D. Twopeny.
1 23½	34	20 1	?		A. O. Hume, C.B.
23½	5 8	16 3	Bikanir	•	Capt. F. E. S. Adair.
23½	41/2	171	Do		Major H. C. Morland.

¹ Malformed, curving backwards like an ibex.

BLACKBUCK or INDIAN ANTELOPE (Antilope bezoartica)—continued.

Length straight.	Circum- ference.	Tip to Tip.	Locality		Owner.
$-23\frac{1}{2}$	5	17½	;		James J. Harrison.
23½	5	$24\frac{1}{2}$	Bikanir .	•	U. O. Thynne.
$-23\frac{1}{2}$	51/2	18	Gaziabad		Major G. Douglas.
231	5½	$18\frac{1}{2}$	Do		Hume Collection, British Museum.
231	43	173	Sirhind .		Col. R. J. Heber-Percy.
23	5 Å	138	North India	•	Sir Victor Brooke's Collection.
23	51	15‡	Indore .	•	Col. Cunliffe Martin.
23	51	15½	?		H. G. Buxton.



Head of Blackbuck.



Skull and Horns of Impala, from a specimen shot by A. H. Neumann.

IMPALA (Æpyceros melampus).

Inzero of the Masubias. Swalah of the Swahilis.

Mpala in the Barotse, Lake Ngami, and the Chilala and Chibisa.

This beautiful impala, or pala, differs from all the members of the gazelle group by the absence of the lateral hoofs; its specific name (black-footed) being taken from the pair of tufts of black hair on each hind foot. The horns of the bucks, which are of considerable length in proportion to the size of the animal, show a characteristic and graceful double curvature; and the bright foxy red hue of the shining fur of the upper parts aids in the easy identification of the species. Height at shoulder about 33 inches. The Boer name of rooibok (red buck) is happily chosen. Weight about 110 lbs.

Distribution.—Southern and South-Eastern Africa, extending as far north as Lower Kordofan. In Angola replaced by a closely allied form (Æ. petersi), usually regarded as a distinct species, but which may be merely a local race; it is distinguished by the presence of a purplish black streak down the middle of the face and another

IMPALA 207

through each eye. In the days of their abundance impala were to be found in big troops; and such are still to be met with on the Upper Zambesi, in East Mashonaland, and parts of British East and Central Africa. Half a century ago, or even less, they were to be found in similar numbers among the covert on the banks of every river in the Transvaal and Bechuanaland; but now it is not till the northern borders of the former country that they are to be met with, and even then only in small parties here and there. Impala are some of the fleetest of all antelopes, and are also in the habit of leaping high in the air; their presence always implies the neighbourhood of water.

Len	gth.				
On front curve.	Straight.	Circum- ference.	Tip to Tip.	Locality.	Owner.
30	24	61	171	E. C. Africa	Lord Delamere.
29	221	58	11	Do	Do.
28	23	5 8	22 3	Do	A. H. Neumann. (See illustration.)
•••	22 7		•••	Masailand	F. J. Jackson, C.B.
28	21	5 7	124	E. Africa	G. E. Smith.
27½	21	5 8	144	South Africa	The late Dr. Burchell, British Museum.
27	23	6	231	Between Lakes Bar- ingo and Rudolph	H. Andrew.
27	$22\frac{1}{2}$	•••	151	Between Lakes Nai- vasha and Baringo	F. J. Jackson, C.B.
27	$22\frac{1}{4}$	6	104	E. Africa	Major A. E. Smith.
-27	211	6	9	Do	E. Gedge.
-26 1	20 ³	•••	•••	South Africa	Grahamstown Museum.
26 8	21	5€	16	Do	Hon. Walter Rothschild.
26 1	22 <mark>3</mark>	5 1	15	Uganda	Col. F. D. Lugard, C.B., D.S.O.
26 1	194	54	221	E. Africa	C. W. Hobley.
25½	211	6 1	127	Do	G. E. Smith.
•••	20½	54	15	Do	W. Astor Chanler.
	20½	5₹	111	Do	H. C. V. Hunter.
-25½	20	6	14	Matamiri Bush .	F. Vaughan Kirby.
25½	20	6	132	E. Africa	Henry Charrington.
25 1	20½	5	125	Do	F. J. Jackson, C.B.
-25.13	20	5.12	15.7	Do	Prince Henry of Liechtenstein.

IMPALA (Æpyceros melampus)—continued.

Len	gth.					
On front curve.	Straight.	Circum- ference.	Tip to Tip.	Locality.		Owner.
-25 1	201		102	3		F. J. Newnham.
25	211	5 2	178	East Africa .		F. J. Jackson, C.B.
24 7	193	5 7	115	Do		Sir Edmund G. Loder, Bart.
24½	191	5 8	128	S. Africa .		British Museum.
	20	•••		Chobe Valley .		F. C. Selous.
	20			E. Africa .		Sir John Willoughby, Bart.
24	20	5 1	16	Do	•	J. Gardiner Muir.
24	20	6	114	Do		E. J. L. Berkeley.
-24	20½	5½	131	S. Africa .		Dublin Museum.
-24	19	51/2	10	Do		Mr. Justice Hopley.
-24	•••	61/2	16 1	E. Africa .	•	T. E. Buckley.
235	192	58	16 8	Do		Capt. R. A. J. Montgomerie, C.B., R.N.
235		5½	9 8	Do		Hon. Walter Rothschild.
$-23\frac{1}{2}$	19		12 §	Lydenburg .		F. H. Barber.
$23\frac{1}{2}$	19	5	8	E. Africa .		C. F. S. Vandeleur, D.S.O.
23 8	20 8	6	141	Do		Sir Robert Harvey, Bart.
23 1	198	58	I 23	Ndi, E. Africa		C. Steuart Betton.
-23	181	5	10	S. Africa .		A. Ohlsson.
22 3	183	6	147	E. Africa .	•	Major W. H. Williams.
$22\frac{1}{2}$	$18\frac{1}{2}$	5½	148	Matabeleland .		Hon. R. A. Ward.
-22½	18		•••	?		O. R. Dunell.
-22	19			S. Africa .		Cape Town Museum.
	19			Njoko River .		R. T. Coryndon.
22	18	6	101	S. Africa .		James J. Harrison.
213	18	5	9‡	Pungwe		Hon. T. Thynne.
211	171	$5\frac{1}{2}$	124	S.E. Africa .		E. Lort-Phillips.
211	183	51	10	Do		Col. G. A. Percy.
211	171	5 1	111	Do		G. L. Bonham.
211	17	5	81	Do		Viscount Ennismore.

IMPALA 209

The following specimens belong to the small form described as AE. *johnstoni*, now regarded as inseparable from the ordinary impala:—

Ler On front curve.	ngth. Straight.	Circum- ference.	Tip to Tip.	L	ocal	ity.		Owner.
198	15 2	47	5€	Choma l	Riv	er, B.C	.A.	R. H. Ferrers Stranach.
19	143	5 1	7	B.C.A.				John Yule.
171	143	44	9₽	Do.				C. C. Bowring.
17	141/2	5 1	8	Do.	•	•	٠	Hon. Walter Rothschild.
		A NTC	OT A NT	TREDAT		/ ED		

ANGOLAN IMPALA (Æpyceros petersi).

See page 206.

23 1	183	6	121	Kaokoland		Capt. F. Cookson.
21	17 2	5 7	114	Angola .		Hon. Walter Rothschild.
-21	17 8	6	93	?		Count E. Hoyos.



Head of Impala.



Horns of Saiga, from the Hon. Walter Rothschild's specimen.

SAIGA (Saiga tatarica).

One of the most remarkable of all antelopes is the Central Asian saiga, which, although belonging to the same group as the blackbuck, differs from most of its kindred by the extraordinary inflated and puffy shape of the nose. This forms a kind of trunk, comparable to that of the dik-diks, with the nostrils directed downwards. The tail is short, and lateral hoofs, which are wanting in the impala, are present. In summer the colour is dull yellowish above and whitish beneath, but in winter the whole fur is uniformly whitish. The short and blunt ears are very thickly covered with hair, and the horns of the males pale amber colour. Height at shoulder about 30 inches.


Distribution.—The open steppes of Southern Russia and South-Eastern Siberia. Formerly the range of this animal was much more extensive in Europe, reaching to the confines of Poland, but nowadays it is restricted to the Kalmuk Steppes between the rivers Don and Volga.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
148	5‡	$3\frac{1}{2}$	Siberia		Hon. Walter Rothschild. (See illustration.)
13#	5	•••	Volga		Sir Edmund G. Loder, Bart.
133	4 3	31/2	?		Sir Victor Brooke's Collection.
13§	5	51/2	Sarepta, South Russia	•	British Museum.
131	58	48	Russia		Hon. Walter Rothschild.

SAIGA 211

SAIGA (Saiga tatarica)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
127	58	single horn	Russia		•	British Museum.
12 7	48	5 8	Sarepta, South Russia			Do.
$-12\frac{1}{2}$	5	44	Siberia	•	•	Dublin Museum.
11	4 7	41/2	Do			British Museum.



Head of Male Saiga.



Head of Chiru, from H. Z. Darrah's specimen.

CHIRU or TIBETAN ANTELOPE (Pantholops hodgsoni).

Among several animals peculiar to the Tibetan plateau none is more interesting than the so-called chiru, whose beautiful horns form some of the most cherished trophies of the sportsman. Although very unlike in general appearance, the chiru is evidently a near ally of the saiga, having the nose even more swollen at the sides, at least in the male, but less bent downwards at the tip. The long black horns, which are somewhat compressed, rise almost vertically from the head, and are slightly divergent, nearly straight below, but evenly curving forwards above, and ridged in front. General colour of fur, which is very dense and short, pale fawn above, with a pinkish suffusion, but the face of the male black. Height at shoulder about 31 or 32 inches; weight from 90 to 120 lbs.

Distribution.—The plateau of Tibet, at elevations of from 13,000 to

16,000 feet, or even more; the animal associates in pairs or small parties.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
274	6 1	131/2	From the Darma Side across the Beansi Pass	A. O. Hume, C.B.
27§	5 1	111	Tibet	Sir Robert Harvey, Bart.
27 1	5 7	15 1	Hills N. of the Beansi Pass	A. O. Hume, C.B.
-27	•••	12	?	J. D. Inverarity.
-26½	•••	•••	Chang Chenmo	Major A. E. Ward.
26 §	57	13§	Ladak	Hon. Charles Ellis.
26	48	127	Tibet	Captain G. Campbell.
25 §	5 1	124	?	Arnold Pike.
25½	5 3	I I ½	Tibet	Captain C. B. Vandeleur.
25 8	5 t	121	Hills north of Leh .	Hume Collection, British
251	52	. 12 3	Hills north of Kumaon	Museum. Do.
25	5₹	111	Chang Chenmo	H. Z. Darrah. (See illustration.)
-251	5	108	Tibet	P. H. G. Powell Cotton.
24 ³	58	1112	?	Sir Edmund G. Loder, Bart.
243	5	11	?	E. L. Phelps.
24 🖁	5€	112	Tibet	Sir Robert Harvey, Bart.
$-24\frac{1}{2}$	44	•••	Do	Major-General Alexander A. A.
24 8	5‡	16 <u>1</u>	?	Kinloch. British Museum.
-24	•••	•••	Changchingris	Otho Shaw.
-24	5 1	12	?	T. E. Buckley.
-24	•••	•••	Chang Chenmo	Captain H. Trevor.
-24	•••	•••	?	Indian Museum.
-24	•••	•••	?	H.H. the Maharaja of Travan- core, G.C.S.I.
-24	51	121	Tibet	Captain C. B. Vandeleur.
-235	5 1	115	Do	H. C. V. Hunter.
-23§	5 1	137	Chang Chenmo .	Paris Museum.
$-23\frac{1}{2}$	5½	81	Do	P. Church.
23½	5	151	Do.	Major H. C. Morland.
23½	5	143	Do	Reginald Beech.
23½	51	127	Do	Captain H. Trevor.
23 ⁸	5 1	I 1 8	Ladak	Major C. S. Cumberland.
231	5 8	148	Do	Hon. Walter Rothschild.
231	5 8	133	Do	Captain H. M. Biddulph.
231		114	Do	Colonel R. J. Heber-Percy.
23 1	5	8‡	Do	G. B. Milne.
23k	5½		Do	Dr. W. P. Y. Bainbrigge.
23	48	114	Do	Captain II. Trevor.
23	5	15	Do	H.R.H. le Duc d'Orléans.



Head of Springbuck, from F. H. Barber's specimen.

SPRINGBUCK (Antidorcas euchore).

Although nearly allied to the gazelles, both in bodily form, coloration, and the shape of the horns, the springbuck presents a peculiarity entitling it to be regarded as the representative of a genus by itself. This peculiarity is the presence of a fold, or narrow pouch, running down the middle of the hinder part of the back, and lined with long pure white hairs. In periods of excitement this pouch is capable of being turned partially inside out, when the long white hairs are erected, and give a totally different appearance to this region of the animal. Owing to the dark nose-streak stopping short at the level of the eyes, the springbuck has more white on the face than is the case with the majority of gazelles; and the white on the rump, which includes the tail and joins that of the middle of the back, also occupies a larger area. Height at shoulder, 30 inches.

Distribution.—The plains of Southern Africa, extending in the central districts of the continent to about latitude 20° S., where its limits are defined by the forests to the south of the Mabebe River; in the west ranging as far north as Mossamedes and Benguela in Southern Angola, and in the east at least up to the Limpopo. Although now never found in the countless thousands which formerly swarmed over the plains of the Transvaal and Bechuanaland, springbuck are still

abundant in many districts. Protected to a certain degree by law, they are to be met with in parts of Cape Colony and the Orange Free State; while on the plains bordering the Botletli and the neighbouring salt-pans, as well as in Great Namaqualand, Damaraland, and the Ovampo Flats, they occur in large numbers. Spring-buck-stalking on the open veldt affords excellent rifle practice; zest being added to the sport from the fact that the venison is most excellent for the table.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
1 19	5	201	Kalahari	The late W. F. Webb.
-17			Middleburg Flats	F. H. Barber. (See illustration.)
-16 1	•••	•••	Bechuanaland	Vryburg Club.
15½	$6\frac{1}{2}$	7‡	?	Hon. T. Thynne.
-15½	6 <u>1</u>		?	Dr. Maloney.
151	6 1	$3\frac{1}{2}$	Cape Colony	C. D. Rudd.
-15½	6	23	S. of Great Namaqualand	Th. Rehbock.
-154	5½	418	?	Dr. W. P. Y. Bainbrigge.
15 1	61	5 7	Ovampo Flats	Capt. F. Cookson.
15 1	5\$	107	?	The late W. C. Oswell.
15	51/2	8‡	?	Capt. E. J. Lugard.
-15	6	6‡	?	A. Ohlsson.
-15			Langberg, Kimberley .	Cape Town Museum.
$-14\frac{1}{2}$?	Alfred Ebden.
141	7	$2\frac{1}{2}$	South Africa	A. W. Davis.
141	61	5	Do	C. Ansell.
14	5≇	41	?	British Museum.
14	5½	5	?.	F. E. Potter.
13 7	6	5#	Griqualand	F. C. Selous.
135	5‡	5₹	Benguela	G. W. Penrice.
135	5 1	7 1 8	?	Sir Victor Brooke's Collection.
131	5 1 2	5	Botletli River, Ngamiland	H. A. Bryden.
131	5 8	34	?	Sir Edmund G. Loder, Bart.

¹ A malformed specimen.

SPRINGBUCK (Antidorcas euchore)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
$-13\frac{1}{2}$		•••	?	O. R. Dunell.
138	5 2	48	South Africa	F. C. Selous.
131	81	41/2	Do	R. A. Cooper.
13‡	6	4	Do	Hon. John Ward.
₽ 13			Middleburg Flats	F. H. Barber.
- 9 13			South Africa	Alfred Ebden.
13	5 1	6	Britstown, South Africa.	W. S. Curtis.
124	51	3 1	Cape Colony	Rowland Ward.
125	5∰	7 1	Orange Free State	Julius Jeppe.
-12½	5	4	Do	Do.
121	51/2	41/2	South Africa	P. H. Illingworth.
-I2 8	418	7 18 .	Transvaal	Dr. Percy Rendall.
-12 8	5 1	48	Orange Free State	Count E. Hoyos.
♀ 12å	27	•••	S. of Great Namaqualand	Th. Rehbock.
-912	•••	•••	Damaraland	F. H. Barber.
-12	•••	•••	?	C. T. Jones.
- 2 114	3	3	3	Dr. W. P. Y. Bainbrigge.
δ 10 3	3 1	5 \$	South Africa	F. C. Selous.
δ 10 β	38	$2\frac{1}{2}$	Ovampoland	Capt. F. Cookson.
♀9 1	38	4 1	Benguela	G. W. Penrice.
. 8§	$2\frac{1}{2}$. 4	Natal	British Museum.
- 2 8 ¹ / ₈	211	3 18	Transvaal	Dr. Percy Rendall.
₽8	31	$2\frac{1}{2}$	South Africa	A. Beit.
₽ 7 1	28	27	Do	F. C. Selous.



GOA or TIBETAN GAZELLE (Gazella picticaudata).

The true gazelles, from which the springbuck is now separated as a genus by itself (Antidorcas), form a very extensive group of delicately built antelopes of easy definition. They are of medium or rather small size, with the muzzle of ordinary shape, the neck not unduly elongated, and no fold containing a crest of long erectile hairs down the middle of the back. To suit the nature of their haunts, their coloration is generally of a sandy hue, with the under parts white, and the face in most cases marked with parallel dark and light longitudinal streaks; dark bands being also frequently present on the rump and on the flanks to separate the tawny of the back from the white beneath. The knees are very generally furnished with brush-like tufts of long, stiff hairs; and the tail is either short or of medium length. With the exception of four species, horns are present in both sexes; those of the males being stout, distinctly ridged, and generally of about the same length as the head, although occasionally much longer. Except at the tips, they curve backwards, so as to present an anteriorly convex lower portion, above which they are generally more or less curved forwards and upwards. Gazelles have a wider geographical distribution than any other genus of antelopes, and are, for the most part, inhabitants of more or less desert regions, or their confines.


The Tibetan goa belongs to a small and aberrant group of the genus in which the tail is very short, the usual dark and light streaks on the face are wanting, and the females are hornless. As a species, it

is distinguished by its comparatively small size, and the strongly marked backward curvature of the horns, which are not hooked at the tips. Height at shoulder about 25 inches. Weight about 45 lbs.

Distribution—The plateau of Tibet and some of the adjacent

Distribution.—The plateau of Tibet and some of the adjacent parts of Central Asia.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
141	35	2	Hanle, Spiti	Hon. Walter Rothschild.
-131/2		•••	?	LieutCol. T. Greenaway.
131	38	5‡	Hills N. of Sikim .	Hume Collection, British Museum.
- 131	4	•••	Tibet	Major-General Alexander A. A. Kinloch.
-13‡	3½	3	Ladak	Capt. H. W. Codrington.
-13 1	4	21/4	Do	Do.
131	4	35	Tibet	H. C. V. Hunter.
-13 1	3 🖁	6‡	S.E. of Hanle .	P. H. G. Powell-Cotton.
-13	37	57	Tibet	Bombay Natural History Society's Museum.
13	4 h	$4\frac{1}{2}$	East Ladak	Col. J. Biddulph.
13	37	1 ½	?	Captain C. B. Vandeleur.
13	•••	•••	?	Indian Museum.
13	34	58	N. Sikim	Surgeon-Captain A. Pearse.
$12\frac{7}{8}$	3 7	47	Hills N. of Kumaon	A. O. Hume, C.B.
127	3#	4#	?	The late B. H. Hodgson, British Museum.
123	37	21/2	Hills N. of Kumaon	Hume Collection, British Museum.
125	47	5 l	Near Hanle	H. Z. Darrah.
121	38	7	?	Arnold Pike.
128	$3\frac{1}{2}$	5½	South of Hanle .	Col. F. C. Lister-Kay.
121	3½	6]	Ladak	David T. Hanbury.
-121	3½	31 ⁸ 0	Tibet	Prince Henri d'Orléans, Paris Museum.
-12g	$3\frac{1}{2}$	61	Do	Sir Edmund G. Loder, Bart.
12	3₹	5	Ladak	Capt. F. E. S. Adair.
-12	3 4	5	Tibet	Capt. H. M. Biddulph.
12	3∰	$2\frac{1}{2}$	Do	H. C. V. Hunter.
12	4	5½	Do	Rowland Ward.
112	34	$3\frac{1}{2}$?	R. Johnstone.
IIĝ	38	5 7	Tibet	The late B. H. Hodgson, British Museum.
117	3€	41	Do	Hon. Walter Rothschild.
111	$3\frac{1}{2}$	41	Hanle	F. W. H. Walshe.
114	3 3	41	?	J. V. Phelps.
11	39	3∦	Ladak	R. Lydekker, British Museum.




Skull and Horns of Przewalski's Gazelle, from a specimen shot by Prince Demidoff.

PRZEWALSKI'S GAZELLE (Gazella przewalskii).

Nearly allied to the goa, from which it differs by its rather larger size, and the distinct hooks formed by the tips of the horns of the bucks. General colour in summer, deep fawn; in winter, pale finely grizzled fawn; white of buttocks running up in an angle on each side of the tail, which is very sharp, and almost concealed by the fur. Front of limbs more or less brown; no tufts of hair on the knees.

Distribution.—Mongolia.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
11	48	48	Altai	St. George Littledale.
10‡	48	$2\frac{3}{4}$?	British Museum.
10	4	48	?	Henri de Bourbon, Comte de Bardi.
10	41 ³ 0	6	Altai	Prince Demidoff.
93	418	48	North of Pekin .	The late R. Swinhoe, British
-811	3 3	48	North China	Museum. Père A. David, Paris Museum.



Skulls and Horns of Mongolian (1, 3, 5) and Persian (2, 4) Gazelle. From specimens in the Collection of A. O. Hume.

MONGOLIAN GAZELLE (Gazella gutturosa).

Jaeisw or Sava-keek in Turkestan. Hwang-yang in Mongolia.

Belonging to the same group of the genus as the goa, this larger species takes its scientific title from the peculiar swollen condition of the throat of the male during the pairing season. In addition to its size (height at shoulder about 30 inches), the species is characterised by the comparatively slight backward curvature of the horns, which are not hooked at the tips; the general colour being pale fawn, with the rump, flanks, and the whole of the limbs white.

Distribution.—Northern and Eastern Mongolia and the southern borders of Russian Transbaikalia. This gazelle was discovered by the Russian traveller Pallas, who described it as long ago as the year 1777.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
152	5	6.34	Lob Nor	A. O. Hume, C.B.
143	48	3	Chinese Turkestan .	Major C. S. Cumberland.
-148	31 ³ 6	43	Lob Nor	Prince Henri d'Orléans, Paris Museum.
14‡	41	3	Wana Plain, Wazir- istan	A. J. Grant.
134	48	$4\frac{1}{2}$	Chinese Turkestan .	E. L. Phelps.
13 3	41/2	6	Saissan, West Siberia.	Dr. O. Finsch, British Museum.
138	4 2	5½	?	Hume Collection, British Museum.
-131	3 3	31	3	Capt. P. J. Gordon.
13 1	48	61	Mongolia	Sir Edmund G. Loder, Bart.
1112	48	3	Chinese Turkestan .	Reginald Beech.

PERSIAN GAZELLE (Gazella subgutturosa).

A near relation of the Mongolian species, but distinguished by the longer tail, of which the upper surface is crested with black, and the presence of dark and light markings on the face. These face-markings differ, however, from those of all other species save the Marica gazelle (G. marica) of Arabia, in that the upper portion of the face, at least in aged individuals, is pure white, so that the central dark band is interrupted on the forehead. From the Marica the Persian gazelle may be distinguished by its superior size (height at shoulder from 26 to 27 inches), and the absence of horns in the female. The larynx is swollen, forming a peculiar prominence on the front of the upper part of the throat.

There are two distinct local races of this gazelle, the ordinary Caspian and Persian G. subgutturosa typica, and the Yarkand G. subgutturosa yarcandensis; the latter being distinguished by its superior dimensions and darker face-markings, as well as by the much smaller degree of divergence of the horns of the males, which are also relatively shorter.

Distribution.—Western Asia from Asia Minor and Caucasia in the west to Turkestan, Yarkand, and Mongolia in the east.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-141	5‡	6	Near Tiflis .	 Clive Phillipps-Wolley.
131	41	4	Khelat .	 A. O. Hume, C.B.
I 2 1 g	4	5‡	Syria	 Hon. Walter Rothschild.
12	4	5	Asia Minor	 Do.
12	4	23	Khelat .	 A. O. Hume C.B.

The undermentioned belong to the Yarkand race.

16	5	31	Yarkand	A. O. Hume, C.B.
157	5	3‡	Do.	Do.
-154		5	Maralbashi .	H. Lennard.
14	41	5.75	Yarkand	A. O. Hume, C.B.
14	5	53	Eastern Turkestan	Col. J. Biddulph.
131	4 8	6	Maralbashi .	David T. Hanbury.

PERSIAN GAZELLE (Gazella subgutturosa)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality	·•	Owner.		
-138	51 ³ 6	7 1 8	Altai			Prince Henri d'Orléans, Paris Museum.	
13	48	7 1 8	Maralbashi			P. Church.	
123	48	38	Altai.			St. George Littledale.	
-12		2	Maralbashi			H. Lennard.	



Head of Springbuck.



Head of Dorcas Gazelle.

DORCAS GAZELLE (Gazella dorcas).

This well-known species brings us to the more typical group of gazelles, in the members of which the central dark face-band is continued uninterruptedly up the forehead, and horns are developed in both sexes. Among the special characteristics of the present species may be noted the circumstance that the white area of the rump does not invade the fawn of the back, the indistinctness of the dark lateral band dividing the fawn of the upper parts from the white beneath, and the perfectly lyrate form of the horns, which are of medium length, with the middle portion twisted outwardly, and the tips converging towards one another. Height at shoulder, 21 to 22 inches; total length about 42 inches. General colour pale fawn, of rather variable tint, with the face-markings distinct.

Distribution.—Morocco, Algeria, and thence eastwards through Egypt into Palestine and Syria. By the Arabs of Algeria this gazelle is known as rhozal or hemar.

DORCAS GAZELLE (Gazella dorcas)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.		Loc	ality.			Owner.
13 1	44	$2\frac{1}{2}$	Algeria					Sir Edmund G. Loder, Bart,
$-12\frac{1}{2}$	38	5 1 8	Do.					Viscount Edmond de Poncins.
121	3₹	3 8	Southern	Sah	ara			J. J. S. Whitaker.
128	3 8	24	Algeria					Sir Edmund G. Loder, Bart.
128	38	•••	Do.					British Museum.
121	4	21	Do.					Rowland Ward.
-121	$3\frac{1}{2}$	34	Do.		•			A. E. Pease.
12	4	3 3	Do.					J. H. Thomas.
11 7	38	41/2	Do.	•		•		Sir Victor Brooke's Collection.
-112	3 3	5 1	Do.		•			Julius Jeppe.
114	31/2	37	Do.	•				E. N. Buxton.
-111	38	318	Do.	•	•	٠		W. E. Pease.
-10 7	4	41	Do.	•		•	•	Count E. Hoyos.
101	4	21/4	Shores of	f Red	l Sea			British Museum.
101	38	23	Nubia	٠	•			Sir Victor Brooke's Collection.
- 2 9 8	•••	•••	Algeria				•	A. E. Pease.
-9 8	4	3 3	Do.		•		•	Dr. Percy Rendall.
- 9 9 1	ış	5½	Do.					A. E. Pease.
-83	4	31/2	Do.					Count Scheibler.
-86			Do.					Viscount Edmond de Poncins.

EDMI or ATLAS GAZELLE (Gazella cuvieri).

From the dorcas the present species may be readily distinguished by its superior size, the imperfectly lyrate form of the horns, which diverge more or less regularly upwards, the presence of a black spot on the tip of the muzzle, and the rough character of the hair. General colour dull fawn, with a very indistinct lateral band, well-defined face-markings, the lower portion of the tail crested with black, and the under parts, buttocks, and inner surfaces of fore-legs pure white. Height at shoulder from 26 to 27 inches.

Distribution.—The mountains of Morocco, Algeria, and Tunisia, where it is known by the name of edml or admi.

Length on front curve.	Circum- ference.	Tip to Tip.		Loca	lity.		Owner.
147	5	3∰	Algeria				Sir Edmund G. Loder, Bart.
$-14\frac{1}{2}$	37	$3\frac{1}{2}$	Do.				A. E. Pease.
142	3 8	5 1	Southern	n Sah	ara		J. J. S. Whitaker.
-14 1	•••	•••	Algeria				A. E. Pease.
♀ 13 8	$3\frac{1}{2}$	7 1	Do.			•	E. N. Buxton.
-13 §	41	51/2	Do.				Viscount Edmond de Poncins.
-13½	4	1/2	Do.				A. E. Pease.
I 2 3	4 3	3∰	Do.				E. N. Buxton.
I 2 🖁	3 8	51/2	Do.				Rowland Ward.
128	31/2	5 1	Do.			•	Hon. R. A. Ward.
\$ 11	$2\frac{1}{2}$	5 8	Do.				J. J. S. Whitaker.
- 8 0 3	2 3	4	Do.				A. E. Pease.
₽8	21	5 1	Do.		•		Capt. G. J. Cuthbert.

ARABIAN GAZELLE (Gazella arabica).

Belonging to the same sub-group of the genus as the edmi, this elegant gazelle may be differentiated from that species by its smaller size, smoother hair, and darker coloration, the general tint of the upper parts being dark smoky fawn, with the central face-band rufous fawn, and a black spot on the tip of the muzzle. Height at shoulder, 24 or 25 inches.

Distribution.—Western Arabia, where it is known as ghasal, its Syrian title being ariel or aiel. In spite of its being one of the commonest of the gazelles, and also one which can easily be obtained in captivity, very little authentic information is forthcoming as to its habits, and even its range is not yet fully determined.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
81	4	2	Mocha, South Arabia		W. T. Blanford, British Museum.
- 27½	2	44	3		Dr. Percy Rendall.
48	3 1	3₿	South Arabia		Sir Victor Brooke's Collection.
₹ 4½	13	$2\frac{1}{2}$?		Major H. G. C. Swayne.



Head of Indian Gazeile, from a specimen shot by Loftus M. le Champion.

INDIAN GAZELLE (Gazella bennetti).

Closely allied to the Arabian species, this gazelle (the chikara of the natives and the ravine-deer of many Anglo-Indian sportsmen) is of smaller dimensions and much lighter colour; the height at the shoulder varying from 23 to 24 inches, and the general colour of the upper parts being dull fawn.

Distribution.—Peninsular India, thence extending westward through Baluchistan to the shores of the Persian Gulf. In the Book of Antelopes Messrs. Sclater and Thomas remark that, like the lion and the hunting-leopard, this gazelle belongs to an African type, and appears to have been originally a migrant from the west into India, whence it has spread over the greater part of the peninsula. It should, however, be remembered that extinct gazelles and hunting-leopards occur in the north of India.

Length on front curve.	Circum- ference,	Tip to Tip.	Locality.	Owner.		
-15	5	•••	Rajputana	H.H. Maharaj Rana Bahadur of Ihalawar.		
141	41	88	?	L. M. le Champion. (See illustration.)		
-14.	•••	•••	Ferozepore District	Captain Harry V. Brooke.		

INDIAN GAZELLE (Gazella bennetti)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-14 <mark>1</mark>			Sirsa District, Punjab	Captain H. Trevor.
14	48	5₹	Ferozepore District .	Captain Harry V. Brooke.
-14	•••	•••	Near Lahore	
-13 7	41	61	Do	Kinloch (measured by). C. P. Down.
13\$	41/2	7	?	Sir Victor Brooke's Collection.
132	48	81	Sind	L. Napier.
13∰	48	71/2	Bikanir	C. F. Vander-Byl.
13‡	4	41	?	Sir Edmund G. Loder, Bart.
131	4	6‡	?	Major O. A. Chambers.
-13‡	4‡		Delhi	Major-General Alexander A. A. Kinloch.
-13 1	41/2	4	Hissar	Captain P. J. Gordon.
13	$3\frac{1}{2}$	6	Bikanir	Major H. C. Morland.
-13	318	3 1 8	Gwalior	Viscount Edmond de Poncins.
-13	44	5	3	Major Colvin Stewart.
13	34	6	?	Major R. H. Rattray.
123	41	5	Goorgaon, Punjab .	A. O. Hume, C.B.
123	48	71/2	?	H. C. V. Hunter.
123	41	5 7	North Punjab	Colonel J. Biddulph.
123	41	7	?	Captain L. I. B. Hulke.
- I 2 3	4 3	63	Bikanir	Captain H. W. Codrington.
I 2 3 4	4	6	Jodhpore	H.H. Maharaja of Bikanir.
121/2	4‡	54	Bikanir	LieutCol. G. D. F. Sulivan.
$-12\frac{1}{2}$	•••	•••	?	H.H. the Maharaja of Travan- core, C.C.S.I.
12 8	4	5½	Simrol, Bengal	Colonel John Evans, British Museum.
128	47	5∰	Bikanir	LieutCol. G. D. F. Sulivan.
123	418	6	Do	P. B. Vander-Byl.
- I 2 3	48	6	Kythal, Punjab	Captain M. M'Neill.
121	48	5‡	Dholpur	A. J. Coppinger.
-12 1	4	7	Bikanir	Captain C. B. Vandeleur.
12	4	7 ½	Punjab	Captain A. Hicks-Beach.

INDIAN GAZELLE (Gazella bennetti)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-12	•••		Kathiawar	LieutCol. L. L. Fenton.
12			Banda District	Indian Museum.
12	41/2	•••	Meerut, N.W. Provinces	C. Hastings-Wood.
12	3∰	6	Khandesh	A. Cumine.
-12	41	68	Deccan	VetyCapt. G. H. Evans.
-12	3 2	7 1	Bikanir	H.H. Maharaja of Bikanir.
115	4	4 ³	?	Do.
115	4	41/2	?	Sir Robert Harvey, Bart.
1112	41/2	5‡	Bikanir	Captain F. E. S. Adair.
118	3₹	41	?	Captain N. C. Taylor.
114	41	5	Nagaon, Central Provinces	Colonel R. J. Heber-Percy.
1118	45	58	Etawah	Hume Collection, British
111	41	5	Jodhpore	Museum. Captain G. J. Fitzgerald.
11	4 .	54	Bikanir	Captain C. F. Pinney.
11	3 2	6	Berar	C. H. Seely.
♀ 7 8	g	21/4	?	Sir Victor Brooke's Collection.
- 2 5½	•••	•••	Banda District	Indian Museum.



Head of Female Speke's Gazelle.

SPEKE'S GAZELLE (Gazella spekei).

Dhero of the Somalis.

From the edmi and its allies, with which it agrees in its leading characteristics, this very peculiar gazelle is readily distinguished by the development of a flabby corrugated elevation on the skin of the nose; the general colour of the upper parts being pale brownish fawn, with the lateral band darker than in the other members of the sub-group. Height at shoulder from 23 to 24 inches.

The protuberance on the nose is probably connected with the sexual function; in dead specimens it exhibits a slight cavity beneath the skin which can be inflated by blowing air into the nostrils, and it is therefore probably capable of distension during life.

Distribution.—The plateau in the interior of Somaliland,

SPEKE'S GAZELLE (Gazella spekei)—continued.


Length on front curve.	Circum- ference.	Tip to Tip.	Loc	ality.			Owner.
-12	3 3	51	Somaliland				Viscount Edmond de Poncins.
-12			Northern Son	nalilan	d.		J. Menges.
រេះ	3 3	48	The Haud		•		Major H. G. C. Swayne.
115	3 3	$2\frac{1}{2}$	Do.				Col. A. Paget.
1112	48	23	Somaliland				T. W. H. Clarke.
117	37	4∯	Do.				Viscount Edmond de Poncins.
114	4	4	Do.	•			W. H. Cobb.
11 1	4	5‡	Do.	•	•		C. Liddell.
11	4	4	Do.	•	•		A. H. Straker.
11	37	5	Do.		•		Major H. C. Morland.
11	41	35	Do				Lord Delamere.
104	31	3 3	Do.			•	Sir Edmund G. Loder, Bart.
108	48	54	Do.				B. R. M. Glossop.
103	4	4	Do.				R. M'D. Hawker.
103	37	3	Do.		•		C. N. Welsh.
103	34	34	Do.			•	E. Lort-Phillips.
10 3	4	34	Do.		•	•	B. Vincent.
108	3 7	41	Do.			•	Capt. G. F. T. Leather.
$-10\frac{1}{2}$	418	31/2	Do.			•	Count Scheibler.
101	34	$2\frac{1}{2}$	Do.		•	•	The late W. Babington.
-10 1	48	34	Do.		•	•	J. Brander Dunbar.
10½	3 3	4	Do.	•	•	•	Capt. G. Campbell.
-108	37	$4\frac{1}{2}$	Do.			•	Count E. Hoyos.
108	37	418	Do.		•		Sir H. D. Tichborne, Bart.
€ 01′-	38	34	Do.		•	•	P. H. G. Powell Cotton.
10}	4	41	Do.				F. G. Gunnis.
101	34	34	Do.			•	P. B. Vander-Byl.
101	37	4	Do.			•	H.R.H. le Duc d'Orléans.
-101		•••	Do.				C. V. A. Peel.
-10 1	4	41	Do.	•		•	J. Johnston-Stewart.

SPEKE'S GAZELLE (Gazella spekei)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Loc	ality.	Owner.	
10	37	$4\frac{1}{2}$	Somaliland			Hon. Walter Rothschild.
10 <mark>1</mark>	38	31/2	Do.			W. D. James.
10	31/2	48	Do.			Count J. de Bylands.
-10	4	41	Do.			A. E. Pease.
-10	4	5 1	Do.			Julius Jeppe.
-10	3 3	21/4	Do.		•	Dr. Percy Rendall.
- 9 9 1			Do.			J. Menges.
♀ 9½	2 7	3 1	Do.			T. W. H. Clarke.
₽9	$2\frac{1}{2}$	5‡	Do.		•	Capt. G. F. T. Leather.
₽8	$2\frac{1}{2}$	4	Do.			B. R. M. Glossop.
₽8	21	3 1	Do.			J. Benett-Stanford.
- 9 7 <mark>3</mark>	2	21	Do.			A. E. Pease.
- 973			Do.	•		C. V. A. Peel.



Head of Male Speke's Gazelle.



Head of Male and Skull of Female Pelzeln's Gazelle.


PELZELN'S GAZELLE (Gazella pelzelni).

This species, the lowland gazelle of Somaliland, is nearly related to the last, from which it differs by the absence both of the corrugated elevation on the nose and of the black spot on the muzzle. It is also slightly larger than Speke's gazelle, the height at the shoulder being about 25 inches; and its colour is somewhat more rufous, the light lateral band being distinct, and the dark band rufous brown only somewhat darker than the back, without any tendency to blackness. The dark and light bands on the cheeks are relatively short and indistinct.

Distribution.—The plains of Northern Somaliland, bordering the sea. Within fifty miles of the shore this gazelle is exceedingly numerous, and may often be seen in large herds. It is termed by the natives dhero, in common with Speke's gazelle.

PELZELN'S GAZELLE (Gazella pelzelni)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Loc	ality.			Owner.
-13 18	•••	•••	Northern So	m a lila	nd		J. Menges.
121	34	41	Somaliland				Major H. G. C. Swayne.
121	418	51/2	Do.				Lord Delamere.
12 1	35	41	Do.	•			The late W. Babington.
-12	31	44	Do.				Count E. Hoyos.
12	34	41	Do.				Capt. G. Campbell.
12	3 3	5 1	Do.				Viscount Edmond de Poncins.
111	$3\frac{1}{2}$	5 1	Do.				T. W. H. Clarke.
111	31/2	41	Do.				Major H. C. Morland.
111	31/2	5	Do.				Capt. W. H. Williamson.
111	48	5 1	Do.				A. H. Straker.
111	37	48	Do.				Rowland Ward.
-111	41/2	5 1	Do.				S. Payne-Gallwey.
-111	34	41	Do.				A. E. Pease.
-114	4	31/2	Do.				Sir Edmund G. Loder, Bart.
-114	3€	5	Do.				J. Johnston-Stewart.
111	3₽	41	Do.				Capt. G. F. T. Leather.
11 1	3‡	3₹	Do.				Col. A. Paget.
-11 1	4	5	Do.	•			J. Brander Dunbar.
11	3 8	41	Do.			•	A. S. Trevor.
11	34	38	Do.			٠	Prince Boris Czetwertynski.
11	3 2	34	Do.				R. M'D. Hawker.
103	34	4 1	Do.				Sir Edmund G. Loder, Bart.
10	34	41	Do.				F. G. Gunnis.
-10½	34		Do.	•			C. V. A. Peel.
10 8	3₿	2 7	Do.	•			Julius Jeppe.
♀85			Do.				J. Menges.
- 9 8‡	21/2		Do.				C. V. A. Peel.
- 2 8 1	•••	•••	Do.				J. Brander Dunbar.
₽8	17	34	Do.	•			Julius Jeppe.
₽ 7∰	13	24	Do.			•	T. W. H. Clarke.



Head of Loder's Gazelle.

LODER'S GAZELLE (Gazella leptoceros).

Arab name, Rhim.

Although originally described as long ago as the year 1842, very little was known of this gazelle till specimens were procured by Sir E. G. Loder, and described as a new species under the name of G. loderi. Further investigation proved them to be inseparable from the species named by F. Cuvier. Agreeing with the preceding members of the group in the white of the rump not invading the haunches, the species is easily recognisable by the long and slender form of the horns, and the very pale tone of the coloration of the upper parts, which may be described as pale sandy fawn, with the characteristic gazelline

markings only indistinctly defined. On the face the dark streaks are sandy instead of rufous; and the light bands on the flanks are almost imperceptible, while the dark ones below them are pale sandy with the very slightest tinge of brown; the tail being sandy at the base and gradually darkening to brownish-black towards the tip. In the male the horns are about twice the length of the head, very slender, and closely and heavily ridged almost to the tips. Height at shoulder about 28 inches. Weight, 34 lbs.

Distribution.—The sandy tracts of the interior of Algeria, Tunisia, and Western Egypt, thence extending southwards into Nubia and Sennar. The native name is rhem or rhim (reem).

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.		
157	31/2	6≩	South of Biskra .		Dr. Dawtrey Drewitt.		
141	31/2	51/2	Do		F. H. Barber.		
1476	37	4 3	Algeria		Julius Jeppe.		
141	31/2	101	Do		Sir Edmund G. Loder, Bart.		
144	3 \$	6 3	Southern Tunisia		J. J. S. Whitaker.		
14	37	3	Algeria		Hon. R. A. Ward.		
-14	3 #	5‡	Tue Erg		A. E. Pease.		
14	3 8	44	Algeria		Mr. Justice Hopley.		
138	3∰	51/2	Do		W. E. Pease.		
13½	3 \$	88	Southern Tunisia		Sir Edmund G. Loder, Bart.		
-13 8	•••	•••	Do		A. E. Pease.		
-13			Do. .		J. Menges.		
-134	34	34	Sahara		Viscount Edmond de Poncins.		
134	31/2	44	Do		Rowland Ward.		
134	31/2	54	Southern Tunisia		J. J. S. Whitaker.		
13 1	3 3	418	Algeria		J. H. Thomas.		
-12½	3 3	61/2	Do		Julius Jeppe.		
-I2 1	31/2	5½	Tue Erg		A. E. Pease.		
12	3 7	44	Algeria	•	Hon. Walter Rothschild.		
9 −1 1 §	•••		Southern Tunisia		J. Menges.		
♀ 10 1	2 1 8	$3\frac{1}{2}$	South of Atlas Mts.		Lord Grantley.		
♀9≸	2 ½	37	Algeria		F. H. Barber.		
₽918	17	3 1	Do		Hon. Walter Rothschild.		
♀9 1	2 1	24	Tunisia		Do.		
9 9	21	12	Southern Tunisia		J. J. S. Whitaker.		

ISABELLA GAZELLE (Gazella isabella).

In all the foregoing gazelles with horns in both sexes, the horns themselves have their tips slightly curved either inwards or upwards, and never bent back so as to form a right angle with the basal portion. On the other hand, the present species and the Muscat gazelle (G. muscatensis), while resembling the foregoing in the shape of the white rump-patch and the comparatively light tint of the flank-band, differ by the tips of the horns being hooked inwards or upwards so as to form nearly or completely a right angle. From its ally the present species differs by the pale fawn colour of the upper parts; the lateral bands and other markings being also fawn, instead of blackish. Height at shoulder about 25 inches.

Distribution.—The Red Sea littoral from Suakin to Massowa, and through the interior to Bogosland, Barca, and Taka.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
101	34	28	Anseba River, Abyssinia	Sir Victor Brooke's Collection.
10 1	34	4	Komayli, Abyssinia.	W. T. Blanford, British Museum.
9 2	34	2 8	Bogosland	Sir Victor Brooke's Collection.
-9.14	4	3	Between Suakin and Kassala	Prince Henry of Liechtenstein.
9.11	•••		Do.	Do.
-8.14	4.2	3.12	Do.	Do.
\$7	2	3	Do.	Sir Victor Brooke's Collection.
₽7	1 7	2 8	Kordofan	British Museum.

HEUGLIN'S GAZELLE (Gazella tilonura).

This well-marked species brings us to a small sub-group in which the dark band on the flanks is very strongly defined, and black in colour; the present species being distinguished from its allies by the abrupt inward hooking of the tips of the horns. The general colour is deep sandy, with the central face-band but little darker than the back, no black nose-spot, and the tail sandy at the base but black elsewhere. Height at shoulder about 27 inches.

Distribution.—Bogosland, in North-Eastern Africa. Very little is known of this rare species.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
111	4	2 ½	Sudan		LieutCol. Hon. W. Coke.
10 <u>8</u>	3 4	41/2	?		E. Lort-Phillips.
103	4	21	Bogosland,		British Museum.
101	48	3 ⁸	Abyssinia Do.		Do.
101	41	278	North Africa .		Col. Ralph Vivian.
9 7	48	1 7	Bogosland,		Sir Victor Brooke's Collection.
-91 ¹ 8			Abyssinia Eastern Sudan		J. Menges.
97		I ½	Anseba River, Abyssinia		Sir Victor Brooke's Collection.
9 1	3 3	31/2	North Africa .		Col. Ralph Vivian.
91/2	41	13	Do		Do.
8‡	38	21/2	?		Lieut. E. Lacy, R.N.
72	37	ış	Bogosland,		British Museum.
♀-6 <u>1</u>			Abyssinia Eastern Sudan	•	J. Menges.
♀ 5 1	13	18	Abyssinia .		Sir Victor Brooke's Collection.

SENEGAL or RED-FRONTED GAZELLE (Gazella rufifrons).

This medium-sized and rather stoutly built species agrees with Heuglin's gazelle in the black flank-stripe, but differs by the absence of a distinct inward hooking of the tips of the horns; the general colour being deep sandy rufous, brightening into rich rufous on the forehead and muzzle, without a black nose-spot. The knee-tufts so generally present in gazelles are wanting; and the tail, with the exception of the sandy upper surface of the basal portion, is black. Horns relatively small, regularly divergent, curving at first slightly backwards and then forwards, heavily ridged till the terminal two or three inches.

Distribution.—Senegal and Gambia.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.		
101/2	48	6 8	Senegal			Sir Victor Brooke's Collection.	
♀6 <u>↓</u>	2 la	2 3	Do.			British Museum.	



Horns and Ears of Female Thomson's Gazelle.



Head of Male Thomson's Gazelle.

THOMSON'S GAZELLE (Gazella thomsoni).

Swahili name, Swallah.

The distinct black nose-patch and the great width of the black flank-band serve at once to distinguish this handsome species from G. rufifrons, with which it agrees in the general form of the horns. The general colour of the upper parts is deep sandy rufous, with all the markings well developed and sharply defined; the central face-streak being a deeper rufous, and the light lateral band present, although only slightly paler than the back. There is also a narrow black band bordering the white on the sides of the rump; and knee-tufts are developed. The horns are long, and rather like those of the Indian gazelle on an enlarged scale. Height at shoulder about 25 inches.

Distribution.—The interior districts of British and German East Africa, from Lake Rudolph southwards to Irangi. Discovered by Joseph Thomson during his expedition to the Victoria Nyanza in 1883-84, and named by Dr. Günther in the latter year.

Measurements of 4 males and 1 female shot by F. J. Jackson, C.B.

	ft. in.	ft. in.	ft. in.	ft. in.	ft. in.
Length	3 10	4 I	3 II]	4 4	$3 8\frac{1}{2}$
Height	2 I	2 3	$2 ext{ } 1\frac{1}{2}$	2 2	1 11 1
Weight	52 lbs.	54 lbs.	56 lbs.	62 lbs.	32 lbs.

THOMSON'S GAZELLE (Gazella thomsoni)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
151	44	3 \$	Masailand .			F. J. Jackson, C.B.
151	41/2	81	Do			E. Gedge.
151	42	48	Do			F. J. Jackson, C.B., British Museum
151	41/2	41	East Africa			Major A. E. Smith.
158	47	5₹	Do.			Lord Basil Blackwood.
-14 7	37	34	Do.		•	Julius Jeppe.
147	41/2	41/2	Do.			Col. Trevor Ternan.
142	42	41/2	Do.			Lord Delamere.
141	4	31	Do.			Sir Edmund G. Loder, Bart.
-14 1	41/2	21/2	Do.	•		Count Scheibler.
-14 1	4 2	5₹	Do.	•	•	T. E. Buckley.
141	41/2	42	Masailand .	•	•	J. Gardiner Muir.
148	41	31	East Africa	•		Sir Robert Harvey, Bart.
141	4‡	5 8	Masailand .	•		H. C. V. Hunter, British Museum.
144	41/2	5	East Africa			Col. Trevor Ternan.
14	41/2	44	Do.			Henry Charrington.
14	41	3	Do.		•	S. L. Hinde.
134	41/2	4 2	Do.	•	•	Col. F. D. Lugard, C.B., D.S.O.
-13 8 .	48	3€	Do.	•	•	Mr. Justice Hopley.
131	41/2	5	Lake Naivasha	•		J. Ponsonby.
13 8	48	4‡	Do.	•	•	Sir Robert Harvey, Bart.
138	41/2	58	Do.	•	•	Hon. Walter Rothschild.
-134	34	7 8	East Africa	•	•	Julius Jeppe.
-131	31/2	5 1	Do.	•	•	A. Ohlsson.
134	4 1	42	Lorogi District	•	•	A. H. Neumann.
131	4	5	East Africa	•	٠	E. J. L. Berkeley.
138	44	51/2	Do.	•	•	Capt. J. W. Pringle.
13	41/2	4	Do.	•	•	W. Astor Chanler.
13	41/2	6	Do.	•	•	J. Gardiner Muir.
13	48	41/2	Do.	•	•	G. E. Smith.
-12-12	4	3.13	Do.	•	•	Prince Henry of Liechtenstein.
♀ 5⅓	ığ	31	Lake Naivasha	•	•	F. J. Jackson, C.B.



Skulls and Horns of Grant's Gazelle (male and female), from A. H. Neumann's specimens.

GRANT'S GAZELLE (Gazella granti).

With this handsome species we reach the last group of the genus Gazella, which includes species, for the most part of large size, characterised by the fawn colour of the back being invaded to a greater or less degree by the white of the rump. In the present animal the fawn of the back is cut off from the tail, which is included in the white rump-patch, and there is a dark streak on the hinder border of each side of this patch. On the neck and back the hair has a peculiar wavy appearance recalling that of watered silk. Height at shoulder about 38 inches; weight of male from 150 to 170 lbs., when cleaned about 115 lbs.

Distribution.—East Africa, throughout Masailand, and round Kilimanjaro, north of Baringo, and around Mount Elgon and the Suk country; generally on open grass-lands. Two well-defined local races of this species may be recognised. First, the typical race (G. granti typica), of the interior of East Africa, in which the dark

flank-bands are either wanting, or are present in young specimens below the light flank-band. Secondly, the northern race (G. granti notata), of Northern British East Africa, in which dark bands are present both above and below the light band, behind which they unite with each other. Although described by O. Thomas in 1897 as a local race, the latter form was raised by him to the rank of a distinct species in the following year.

F. J. Jackson says they are found "throughout the Masai country, up in Turkwel and Suk country. Those from Njemps, Turkwel and Suk are smaller than those from Kilimanjaro, Naivasha, Athi plains, etc. In the Njemp district a 20-inch head is good."

Measurements of 4 male specimens shot by the last-named sportsman.

			ft.	in.	ft.	in.	ft.	in.	ft. in.
Length			5	8	5	7	5	7	5 3 ¹
Height	•		3	$I\frac{1}{2}$	3	$2\frac{1}{2}$	3	$6\frac{1}{2}$	2 II
Horns			:	28]		27	:	24	20
Weight			15	8 lbs.	16	7 lbs.	166	5 lbs.	135 lbs.
Habitat		Lake	Na	ivasha,	Lake 1	Naivash	a, Lake	Naivas	sha, Njemp.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.		
281	6 §	151	East Africa			F. J. Jackson, C.B.		
27 ⁷	6 §	18 3	Do.			H. C. V. Hunter.		
271	61	16	Do.			F. J. Jackson, C.B.		
271	6 <u>1</u>	187	Do.			Sir Robert Harvey, Bart.		
27		16	Do.			Sir John Willoughby, Bart.		
26 <u>3</u>	61/2	131	Shores of Lak	te Nai	ivasha	A. H. Neumann.		
26½	71	18	East Africa	•		Major A. E. Smith.		
26 1	6 8	162	Do.			Sir Victor Brooke's Collection.		
26	6 <u>1</u>	108	N.E. of Lake	Rude	olph	A. H. Neumann.		
252	68	15	E. Africa .	•	•	Lord Delamere.		
25 2	6	17	Do		•	J. Gardiner Muir.		
25½	6 1	15	Do			Lord Delamere.		

¹ This was the old buck of a herd in which there were three or four other bucks. It was in good condition. I saw a lot of other bucks, but none had horns more than 20 inches.

GRANT'S GAZELLE (Gazella granti)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
25 1 8	6 1	158	E. Africa .			Capt. R. A. J. Montgomerie,
-25.11	6.8	13.14	Do			C.B., R.N. Prince Henry of Liechtenstein.
-25.10	6.1	15.14	Do			Do.
25	6 §	123	Nakuru Lake,	E. A	Africa	G. E. Smith.
25	$6\frac{1}{2}$	14	Kilimanjaro			T. W. Greenfield.
243	6 <u>3</u>	121	East Africa			S. L. Hinde.
24.13	6.14	14.10	Do.			Prince Henry of Liechtenstein.
241	6 <u>3</u>	131	Do.			E. J. L. Berkeley.
24½	7 8	14	Do.			Sir Edmund G. Loder, Bart.
241	6 1	112	N.E. of Lake	Rudo	olph	Hon. Walter Rothschild.
24	71	8‡	East Africa			Henry Charrington.
231	6§	13 §	Do.			Capt. E. J. Tickell.
231	$6\frac{1}{2}$	13	Do.			Major W. H. Williams.
23	6 <u>1</u>	13 8	Do.			Col. F. D. Lugard, C.B., D.S.O.
23	71	9	Do.			E. Gedge.
-23	6 1	91/2	Do.		•	Julius Jeppe.
♀ 17½	3 2	9	Do.	•		H. C. V. Hunter.
₽ 17		6	Lake Jipé .			Sir John Willoughby, Bart.
₽ 16⅓	31/2	48	E. Africa .			Lord Delamere.
Q 15	4	63	Do			E. Gedge.



Skulls and Horns of Peters's Gazelle (male and female), from A. H. Neumann's specimens.

PETERS'S GAZELLE (Gazella petersi).

This species belongs to the same group as Grant's gazelle and the aoul, showing the same extension of the white of the rump on to the back. It has a dark band separating the white of the rump from the fawn of the back, and differs from the other species showing the same feature by the fawn colour of the back being continued in the middle line on to the base of the tail, which is otherwise black. Height at shoulder about 26 inches.

Distribution.—Coast districts of East Africa.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
22 1	6 3	5 8	Shererini		•	F. J. Jackson, C.B., British Museum.
21 3	51/2	8‡	Pica Pica, B.E.A.	•	•	L. E. Caine, British Museum.
201	6 1	87	Ndura			C. Steuart Betton.
19	6	54	Taru Desert .		•	Lieut. G. H. Welch, R.N.
174	5‡	75	Near Pica Pica .			A. H. Neumann.
§01 ♀	3	41	Do			Do.



Head of Scemmerring's Gazelle, from a specimen shot by G. H. Cheetham.

SEMMERRING'S GAZELLE (Gazella sœmmerringi).

Somali name, Aoul. Abyssinian name, Meidafihel. Abyssinian (Danakil) name, Maédedo.

In addition to the forward extent of the white rump-patch, the leading characteristics of this splendid species are to be found in the absence of a black streak dividing the sides of the rump-patch from the fawn of the body, the black-tipped tail, and the very distinct inward hooking of the horns. The ears are long and bordered with black externally, and the face markings well-defined and nearly black. Height at shoulder about 30 to 36 inches. Weight clean about 70 to 90 lbs.

Distribution.—The Abyssinian coast of the Red Sea, Berber, East Sennar, Danakil, and Somaliland; in the latter country occurring all over the Haud and Ogaden.

a. Gazella sœmmerringi berberana.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Locality.		Owner.		
207	54	38	Somaliland			G. H. Cheetham.		
-201			Do.			Norman B. Smith.		
20	5₹	4	Do.			LieutCol. J. W. H. Flanagan.		
20	5‡	51/2	Danakil .			Viscount Edmond de Poncins.		
♀-20		5₽	Somaliland	•		A. E. Pease.		
-20	5 7	6 1	Do.			Capt. M. M'Neill.		
192	5₩	51	Do.			Count J. de Bylands.		
-194	51	58	The Haud.		•	P. H. G. Powell-Cotton.		
-19] [•••	N. Somaliland	•		J. Menges.		
19½	51/2	5∰	Somaliland		•	T. W. H. Clarke.		
19½	518	2	Do.			Sir Edmund G. Loder, Bart.		
19½	51/2	4 3	Do.			J. J. Richardson.		
-19 1	6₺	3	Do.		•	S. Payne-Gallwey.		
-19 1	51/2	48	Do.		•	Count J. Potocki.		
19‡	61/2	44	Do.		•	R. McD. Hawker.		
19‡	51/2	6 1	Do.		•	T. W. H. Clarke.		
19‡	51	$4\frac{1}{2}$	Do.	•	•	Capt. B. L. Carew.		
19‡	51/2		Do.		•	Surgeon-Major J. S. Edye.		
19	5	4	Do.		•	Col. A. Paget.		
19	5₺	43	Do.		•	J. Benett-Stanford.		
19	5	71/2	Do.	•		B. R. M. Glossop.		
19	5⅓	5 1	Do.		•	Ford G. Barclay.		
19	5	4	Do.			Capt. F. C. Quicke.		
-19	4	31	Do.			Prince Nicolas Ghika.		
-19	5₹	8	Do.	•	•	J. Brander Dunbar.		
-19	•••	4 ⁸	Do.	•	•	J. D. Inverarity.		
-19	5 å	58	Do.		٠.	Mr. Justice Hopley.		

SCEMMERRING'S GAZELLE (Gazella scemmerringi berberana)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
18 2	51/2	6	Somaliland	•		F. H. Barber.
18 §	5 1	51/2	Do.	•		Capt. G. R. Cuningham.
18 <u>1</u>	51/2	75	Do.	•	•	Capt. G. Campbell.
18 1	51/2	$4\frac{1}{2}$	Do.	•		A. H. Straker.
181	5½	5	Do.			P. B. Vander-Byl.
181	5	5₽	Do.		•	A. Leslie Renton.
18 8	51	5	Do.			Sir H. Tichborne, Bart.
-183	5	5 1	Do.			J. Johnston-Stewart.
♀ 18‡	•••	6	Do.			T. W. H. Clarke.
181	58	41/2	Do.			W. D. James.
18	51/2	42	Do.		•	J. Kenneth Foster.
18	5	7 8	Do.	•		Major H. C. Morland.
18	5½	4	Do.	•	•	T. W. Greenfield.
τ8	5‡	5₹	Do.	•	•	R. Wahrmann.
18	5	6 <u>1</u>	Do.	•	٠	P. R. Denny.
-18	•••	$4\frac{1}{2}$	Do.	•		A. E. Pease.
18	58	7 1	Do.	•	٠	W. W. Ashley.
\$ 18	31/2	81	Danakil .			Viscount Edmond de Poncins.
177	5	5₺	Somaliland		•	W. W. Ashley.
-172	51/2	18	Do.		•	Count Scheibler.
172	5ቔ	4	Do.			Major C. C. Ellis.
-174	47	31	Do.			Julius Jeppe.
-174	6	•••	Do.			C. V. A. Peel.
178	5	48	Do.		•	Prince Boris Czetwertynski.
-175	5	3 3	Do.			A. E. Leatham.
171	51/2	6	Do.	٠	•	Capt. C. S. Timins.
171	51/2	5₹	Do.			J. Byng Paget.
171	54	6₹	Do.		٠	W. R. Bindloss.
17½	5‡	7₹	Do.	•		Digby Davies.
- 17 1	5 1	48	Do.	•	•	A. Ohlsson.

SEMMERRING'S GAZELLE (Gazella sommerringi berberana)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Loc	cality.			Owner.
178	6	4	Somaliland	i			Lord Wolverton.
-17 8	48	48	Do. '				Count E. Hoyos.
- 9 174	•••	•••	Danakil				Prince de Lucinge.
♀ 16 <u>3</u>	31/2	6 1	Somaliland	i			H.R.H. le Duc d'Orléans.
9 16	3 1	6	Do.				Major H. G. C. Swayne.
9 16	31	$10\frac{1}{2}$	Do.			•	R. Wahrmann.
		b. Gas	zella sœm	meri	ingi	ty	pica.
151	58	5	Sudan				Col. Ralph Vivian.
₽ 15‡	2 7	78	Do.				LieutCol. Hon. W. Coke.
143	5½	41	Near Suak	in			B. Cotton.
141	5 l	51/2	Eastern Su	ıdan			Major W. H. Besant.
14	51	5\$	Sudan		•		LieutCol. Hon. W. Coke.
14	5 1 2	51/2	Do.	•	•	•	Col. A. Paget.
-14	5.12	3.13		?			Prince Henry of Liechtenstein.
-14	5.2	3. I		?			Do.
132	51/2	6	Sudan		•	•	W. D. James.
131	51/2	4	Do.	•			Sir Victor Brooke's Collection.
131	58	59	Suakin				Hon. Walter Rothschild.
123	5	5	Abyssinia				Sir Edmund G. Loder, Bart.
Ç 12∰	$3\frac{1}{2}$	81	Sudan				Col. Ralph Vivian.
₽ 12	31	5 1 8	Abyssinia		•		Sir Victor Brooke's Collection.
♀ 8 <u>1</u>	34	5	Sudan				Hon. Walter Rothschild.

ADDRA GAZELLE (Gazella ruficollis).

The last three representatives of the genus Gazella are not only the largest of the tribe, but are distinguished from all their relatives by the white of the rump extending on to the tail (which is either wholly white or merely tipped with fawn) coupled with the complete absence of a black band between the white of the rump-patch and the fawn of the body; the horns being hooked upwards and forwards, and not

distinctly inwards. In the present species, which measures about 36 inches at the shoulder, the neck and front portion of the back alone display a distinctly rufous tint, the lines of division between the fawn of the upper parts and the white beneath being obscure.

Distribution.—Kordofan and Sennar.

Length on front curve.	Circum- ference.	Tip to Tip.		Locality.				Owner.		
121	4	58	Sennar					British Museum.		
1 1 7	5₩	§	Kordofan					Do.		

DAMA GAZELLE (Gazella dama).

Together with its northern representative the mhorr of Morocco, the dama differs from the aoul by the rufous tint extending completely over the body and flanks, and being well defined from the white of the under parts. In the present form, which is from Senegal, the sides of the thighs are white, so that the rufous of the body is cut off from that of the hind-legs. The height is at least 35 inches.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-12 5	5 1	5½	?	Paris Museum.
I 2 ½	5 1	6	?	Mr. Justice Hopley.
-121	5 18	318	Senegambia	Paris Museum.
81	31/2	58	?	Hon. Walter Rothschild.

MHORR or SWIFT GAZELLE (Gazella dama mhorr).

Although commonly regarded as a distinct species, this fine but somewhat leggy gazelle seems to be only a northern race of the dama gazelle of Senegal, which is the largest member of the genus. The mhorr, as it is called by the Arabs, differs from the dama by the sides of the thighs and legs being of the same rufous hue as the body, instead of white. Height at shoulder at least $34\frac{1}{2}$ inches.

Distribution.—The desert districts of South-West Morocco.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
I I 🔏	6	3\$	Wednoon, Mogador	W. Willshire, British Museum.
-4 18	3 11	$2\frac{1}{2}$	Morocco	Dr. Percy Rendall.



Skull and Horns of Dibatag, from a specimen shot by T. W. H. Clarke in Somaliland.

DIBATAG or CLARKE'S GAZELLE (Ammodorcas clarkei).

Although resembling the true gazelles in the face-markings, the dibatag, as it is called by the Somalis, is so different from these animals as to be entitled to constitute a genus by itself; being in many respects intermediate between the former and the gerenuk. The horns, which are present only in the males, are rather short, and have a regular upward and forward curvature, somewhat like those of a reedbuck; they are ridged on the front for a considerable portion of their length. The neck is considerably elongated, and the tail long and thin. The general colour of the upper parts is a deep cinnamon. Height at shoulder about 33 inches; weight from 65 to 70 lbs. When running, the long neck is thrown back towards the tail, which is elevated, so that the two look as though they would touch. Discovered by T. W. H. Clarke.

Distribution.—Central Somaliland, in the eastern districts of the Haud.

DIBATAG or CLARKE'S GAZELLE (Ammodorcas clarkei)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
-13			Somaliland		J. D. Inverarity.
-13			. Do		J. Menges.
125	52	48	Do	•	Hon. Walter Rothschild.
113	48	3₹	Do		Col. Arthur Paget.
114	4 8	3₹	Do		T. W. H. Clarke.
$-11\frac{3}{4}$	41	5	Do		J. Brander Dunbar.
114	41	6	Do	•	R. McD. Hawker.
11	41/2	51	Do	•	Capt. F. C. Quicke.
11	51/2	21	Do		B. R. M. Glossop.
11	5	51/2	Do		S. Payne-Gallwey.
104	41	42	Do,		Mr. Justice Hopley.
-10 3	5 1	4	Do		Sir Edmund G. Loder, Bart.
-10½	41/2	41/2	Do		A. E. Pease.
-101	4‡	48	Eidegalla, Haud .		J. Johnston-Stewart.
-10‡	41/2	41/2	Somaliland		C. V. A. Peel.
-10 1	41	48	Somaliland		Rowland Ward.
10 1	48	$4\frac{1}{2}$	West of Hargeisa .	•	P. II. G. Powell-Cotton.
-10	41/2	41/2	Somaliland		C. V. A. Peel.
10	41/2	4	Do		Count J. de Bylands.
-10	4‡	48	?		Julius Jeppe.
9 8	41/2	5 1	Somaliland		T. W. H. Clarke, British Museum.
94	41/2	42	Do		Major H. G. C. Swayne.
91	41/2	44	Do	•	B. Vincent.
-9 16	418	418	Dalbahanta Country	•	Dr. Percy Rendall.
-91	41	41	?		A. Ohlsson.
9	31/2	3	Dalbahanta Country	•	A. S. Trevor.
9	41/2	41	Do	•	Lord Delamere.
88	48	41/2	Do		W. H. Cobb.
8 §	4	44	Do		T. W. Greenfield.

DIBATAG or CLARKE'S GAZELLE (Ammodorcas clarkei)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
$-8\frac{1}{2}$	4	3	Dalbahanta Country	Count Grudzinski.
8	4	3 8	Do	H.R.H. le Duc d'Orléans.
7 3	4	41/2	Do.	Ford G. Barclay.



Head of Male Dibatag.



Head of Male Gerenuk, from a specimen shot by H.R.H. le Duc d'Orléans.

GERENUK or WALLER'S GAZELLE (Lithocranius walleri).

Somali name, Gerenuk. Danakil name, Gudan Godu.

Even more aberrant than the last is the gerenuk, in which the elongation of the neck attains its supreme development, while the slender legs are lengthened in proportion. Horns are wanting in the females, and those of the males curve forwards at the tips, where the ridges stop, in a peculiarly characteristic manner. The general colour of the upper parts is a deep rufous fawn, but down the middle of the back runs a broad dark-brown band, nearly eight inches in width. The skull is characterised by its dense and solid structure, as well as by its straightness, the shortness of the facial portion, and the very small size of the cheek-teeth. Height at shoulder, 41 inches; weight, 115 lbs. (T. W. H. Clarke).

This curious antelope, which was first described by the late Sir V.

Brooke on the evidence of a flat skin, is reported to rear itself on its hind-legs when browsing, and is thus enabled to reach boughs at a very considerable distance above the ground.

The horns of the specimens from Somaliland are, as a rule, very much longer than those from British East Africa, as will be seen by reference to the list below.

Length on front curve.	Circum- ference.	Tip to Tip.	Loc	ality.			Owner.
17	5‡	34	Somaliland				H.R.H. le Duc d'Orléans.
-16 3	•••	•••	Do.				(See illustration.) LieutCol. H. G. Mainwaring.
-16½	54	3∰	Do.				E. L. Cappel.
16 <u>1</u>	51/2	15	Do.				Mr. Justice Hopley.
-16 1	51/2	2 ⁷ 8	Do.	•		•	The late J. Rose.
-16 1	•••	•••	Northern So	malil	and	•	J. Menges.
16	5#	6	Do.		•		Sir Edmund G. Loder, Bart.
152	6	43	Do.		•		Rowland Ward.
-153	6	23	Do.				A. E. Pease.
151	51/2	3	Do.				J. Kenneth Foster.
-151	54	47	Do.				F. H. Barber.
158	58	4	Do.			•	Major H. G. C. Swayne.
15 8	58	41	Do.		•		C. Liddell.
15 8	5#	42	Do.				Lord Wolverton.
-151	5½	1 §	Do.				J. Johnston-Stewart.
15	5₿	6 8	Do.				F. G. Gunnis.
15	5	44	Do.				Capt. J. M'Call Maxwell.
-15	6		Do.				LieutCol. J. W. H. Flanagan.
-15	5	4	Do.			•	Sir H. B. Meux, Bart.
-1418	51/2	7 1	Do.				H.R.H. le Duc d'Orléans, Paris Museum.
142	5 i	42	Do.				Lord Delamere.
143	5 1	3₽	Do.				J. J. Richardson.
141	5	5	Do.				Count J. de Bylands.
-141	5 1	7 1	Do.			•	Count Scheibler.
$-14\frac{1}{2}$	48	4	Do.		•	•	Count E. Hoyos.
148	6 <u>1</u>	41/2	Do.	•	•		Capt. F. C. Quicke.

GERENUK or WALLER'S GAZELLE (Lithocranius walleri)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.				Owner.
142	54	41/2	Northern So	malil	and	•	A. Leslie Renton.
141.	54	31/2	Do.				W. W. Ashley.
14		3	East Africa			•	Sir John Willoughby, Bart.
14	62	41/2	Somaliland				W. H. Cobb.
14	5	31	Do.				Col. A. Paget.
14	5≇	61	Do.				P. B. Vander-Byl.
14	51	43	Do.				Ford G. Barclay.
14	5 1	5 1	Do.				Viscount Edmond de Poncins.
-14	6	5	Do.				S. Payne-Gallwey.
-14	51	4	Do.		•	•	P. H. G. Powell-Cotton.
-14	51	34	Do.				J. D. Inverarity.
138	48	51	East Africa				H. C. V. Hunter.
132	48	418	Do.				E. Lort-Phillips.
132	5∰	27	Northern So	mali	land		J. Menges, British Museum.
134	58	378	Somaliland		•		Hon. Walter Rothschild.
13 2	51/2	3	Do.				Capt. G. F. T. Leather.
132	5 1	5	Do.			•	Digby Davies.
-13 2	5 1	71/2	Do.				C. V. A. Peel.
-13 2	5∯	5	Do.			•	J. Johnston-Stewart.
-138	5‡	5₽	East Africa				T. E. Buckley.
121	58	44	Do.				F. J. Jackson, C.B.



Skulls of Beira, from specimens shot by Lord Delamere.

BEIRA (Dorcatragus melanotis).

In spite of its presenting a certain superficial resemblance to the members of the oribi group and its allies, this peculiar little antelope is considered to be best placed in the neighbourhood of the gazelles. From all the members of the group in which the latter are included, the beira is at once distinguished by the short and spike-like horns of the bucks. Perhaps its most striking peculiarity is the great size of its ears, which led its discoverer, Herr Menges, to describe it as a species of klipspringer. Another noticeable feature is the large size of the rounded hoofs. In colour, the upper parts and legs are pinkish fawn; a darker band defining the fawn from the white of the under parts, which (the white) is continued down the inner surfaces of the limbs as far as the knees and hocks. The head, from the ears to the nose, is bright rufous. Height at shoulder about 23 inches.

Viscount Edmond de Poncins writes that "they are good hillclimbers, and keep on rocky ground. The general shape is slender, legs are long, the head is kept erect, the ears are very big and broad, shaped like the dik-dik's, eyes big and dark, nostrils black.

"Colour a sort of grayish blue, a bit like what we call in French gorge de pigeon, and exactly matching the colour of the ground, so, unless they are on the move, it is difficult to distinguish them. Unlike a gazelle, its tail is generally kept down. Horns resemble those of the klipspringer, but curve forward slightly more." From nose to root of tail $32\frac{1}{16}$, height at shoulder 21 inches, horns 4 inches; weight about 20 lbs.

BEIRA 259

Distribution.—The interior of Somaliland, where it appears to be rare and local, going about either singly or in pairs, and inhabiting the open desert. The beira was first made known to science in 1894: its habits have lately been well described by Captain P. Z. Cox in the eleventh volume of the Journal of the Bombay Natural History Society.

Length on front.	Circum- ference.	Tip to Tip.	Locality.		Owner.	
4 18 1	17	•••	?		Lord Delamere. (See illust	ra-
44	•••	•••	Northern Somaliland		tion.) J. Menges.	
4	2	216	French Somaliland		Viscount Edmond de Poncin	ıs.
4	2 1 8	2 ³	Somaliland		Sir Edmund G. Loder, Bart	
3 }	18	31	?		Lord Delamere.	
34	1 §	•••	French Somaliland		Viscount Edmond de Poncin	ıs.
31/2	111	21	?		Lord Delamere.	
25	18	14	French Somaliland		Viscount Edmond de Poncin	ıs.

¹ Length straight 32% inches, along curves 35% inches; height at shoulder, 25% inches; girth, 19% inches; weight, 23 lbs. as he fell.



Head of Grant's Gazelle.



Skull and Horns of Sable Antelope, from R. T. Coryndon's specimen.

SABLE ANTELOPE (Hippotragus niger).

Impengo of the Masubias.
Impalampala of the Swazis and Zulus.
Ookwa of the Makubas.
Potoquane of the Southern Bechuanas.
Pala-hala of the Swahilis.
Pala-pala of the Makalakas.

Qualata inchu of the Bamangwatos and Makololos.

Solupe of the Masaras.

Utjiele of the Amandebele.

Qualata and T'choo in the Barotse country.

Qualata and Tshumu in the Lake Ngami country.

Nkwalandi in the Chilala and Chibisa countries.

The group of antelopes which includes the present species, the roan antelope, the gemsbuck, addax, etc., differs very markedly from all those before mentioned. Among the leading features are the scimitar-shaped, conical, or spiral horns, which are placed just over the eyes, and are present in both sexes, the hairy muzzle, the absence of face-glands, and the long, tufted tail. Even more remarkable are the upper molar teeth, which have square grinding-surfaces and tall crowns, like those of oxen.

From the other members of the group the beautiful sable antelope and its near ally the roan antelope are well distinguished by the scimitar-shaped horns, which arise at an angle with the plane of the face, as well as by the maned neck, the tufts of long hair below the eyes, and the large size of the ears. By far the handsomer of the two is the present species, whose sable coat and great length of horn render the buck the most striking of its tribe. Other distinctive features are the continuance of the white eye-stripe to the muzzle, the length of the mane, and the relatively moderate size of the ears. Height at shoulder about $4\frac{1}{2}$ feet. A single horn in the Florence Museum measured by Mr. F. C. Selous is 61 inches on the front curve.

Distribution.—From about the centre of the Transvaal northwards to Nyasaland and the adjacent districts of South-East and East Africa. Still abundant in parts of Eastern Mashonaland, and thence towards the coast, as well as on the Manica plateau to the north of the Zambesi. Scarcer in Central East Africa and Mozambique. Admired by all who have seen it in its native haunts, the sable antelope when wounded is a dangerous antagonist, to be approached with extreme caution. It runs with considerable speed, and possesses much staying power.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-47 8	•••	•••	Rhodesia	Measured by F. C. Selous (Fielden's specimen).
471	9 1	26½	?	R. T. Coryndon. (See illustration.)
-46	•••	•••	Eastern Border of Transvaal	W. Russell Bowker.
45½	9‡	151	Mashonaland	Major G. A. L. Carew, D.S.O.
458	10	142	Lebombo Mountains	F. Vaughan Kirby.
454 .	91	124	S.E. Mashonaland	J. G. Millais.
-45 1	•••	121	Pandamatenka	R. T. Coryndon.
-451	91/2	$20\frac{1}{2}$?	Mr. Justice Hopley.
-45			Batoka Country	R. T. Coryndon.
-45	101		South Africa	Earl of Dartmouth.
-45			Angwa River	Capt. J. A. Spreckley.
44 ³	108	131	Barotseland	H. Timmins.
441	91	16	?	R. T. Coryndon.
441	9	17	Transvaal	J. P. Fitzpatrick.
-44½			?	H. T. and A. H. A. Glynn.
-44½	(single l	horn)	Batoka Country	F. V. Worthington.

SABLE ANTELOPE (Hippotragus niger)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
448	91	14	Eastern Transvaal	F. Vaughan Kirby.
441	91/2	8‡	Eastern Mashonaland .	F. C. Selous.
-44			Batoka Country	F. V. Worthington.
-44		•••	Lydenburg	F. H. Barber.
-44	10	32½	?	Berlin Museum.
-43 3	97	8	Matabeleland	Count E. Hoyos.
43½	91/2	$9\frac{1}{2}$?	Sir Edmund G. Loder, Bart.
43½	9 2	15	Matabeleland	The late J. S. Jameson.
434	91	101	Mashonaland	S. Chillingworth.
43	108	16 1	Chobe Valley	F. C. Selous.
43	9₽	6 1	Do	F. C. Selous, British Museum.
43	•••		Do	Major Hon. R. T. Lawley.
43	10‡	13	Mashonaland	Major H. J. Goold-Adams, C.B., C.M.G.
42 7	94	478	Eastern Transvaal	F. Vaughan Kirby.
42 3	98	121	Mashonaland	J. A. Jameson.
428	9 1	91	South Africa	Bethnal Green Museum
421	9 8	6	Mashonaland	G. H. M. Banks.
-42½	•••	•••	?	Cape Town Museum.
421	91	5	Rhodesia	W. W. Ashley.
-42	10	31/2	South Africa	Dublin Museum.
-42	•••	•••	Transvaal	Cape Town Museum.
-42	•••	•••	S.E. Mashonaland	J. G. Millais.
414	98	12	Do	A. Beit.
412	•••	91	Do	A. C. Fountaine.
412	91	9	Muchinga Plateau, Northern Rhodesia	F. Smitheman.
411	9	5‡	Do	C. D. Rudd.
411	9 1	16½	?	Col. F. Rhodes, D.S.O.
411	10	127	S.E. Mashonaland	Hon. Walter Rothschild.
411	108	$11\frac{1}{2}$	Sabi River	Earl of Dunmore.

SABLE ANTELOPE (Hippotragus niger)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
411	9	101	South Africa	R. A. Cooper.
-411	8 <u>1</u>	172	Do	Julius Jeppe.
41	91	103	Matabeleland	W. Van Ness.
41	10	12	Do	Hon. R. A. Ward.
41	91/2	125	Do	Capt. Sir K. Fraser, Bart.
41	91	12	Do	Major James Grant.
-41	91	9	Pungwe	James J. Harrison.
40½	8	13	N.E. Transvaal	A. M. Naylor.
-40 8	91	15	Sabi Flats	Dr. Percy Rendall.
40 8	98	16	Upper Zambesi	M. C. Greaves-Bagshawe.
401	10	13	?	F. Struben.
401	91	13	Matabeleland	Hon. R. A. Ward.
-40 1	9‡	15‡	South Africa	Dr. W. P. Y. Bainbrigge.
40	10	22	Matabeleland	Abe Bailey.
40	91	101	Near Ruo River, S.E. Afric	a C. C. Bowring.
-40	•••	•	Nyasaland	S. Pulley.
-40	10	131	South Africa	A. Ohlsson.
-39 1	•••	•••	?	O. R. Dunell.
-39 1			Mashonaland	J. Vaughan.
♀ 39 1	61	6 §	Do	F. C. Selous.
39	10	74	Do	F. B. Dunsford.
39	9	10 <mark>8</mark>	?	Sir Victor Brooke's Collection.
38 2	9 3	$2\frac{1}{2}$	Matabeleland	Rev. Dr. R. J. Nevin.
38 <u>1</u>	8 2	13	Pungwe	Capt. F. H. Lehmann.
38 1	91	9	Matabeleland	Duke of Roxburghe.
38 1	9	151	Do	Hon. C. Greville.
38	9 8	147	Mashonaland	D. Norman Ritchie.
37	9	14	Matabeleland	Hon. John Ward.
-37	•••		?	C. T. Jones.

SABLE ANTELOPE (Hippotragus niger)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
321	9	153	British Central Africa	ι.	•	J. E. Gough.
♀ 35 8	6₿	•••	Lebombo Mountains			F. Vaughan Kirby.
♀34₺	6 <u>₹</u>	10	Matabeleland .			Hon. R. A. Ward.
♀338	6 3	121	?			G. Richards.
♀ 33₺	6 §	3 3	Sabi Flats			Dr. Percy Rendall.
♀32		•••	Mashonaland .			F. C. Selous.
-♀32			?			H. T. and A. H. A. Glynn.
♀ 31½	7	125	Algoa Bay			British Museum.
₽ 31	7	71/2	Matabeleland .			J. A. Pease.
₽ 30	61/2	$11\frac{1}{2}$	Do			Lord Brackley.



Head of Sable Antelope, from a specimen shot by F. C. Selous.



Head of Roan Antelope, from a specimen shot by F. C. Selous.

BOAN ANTELOPE (Hippotragus equinus).

Ee-taka of the Amandebele.
Ee-pala-pala chena of the Makalakas.

Impengo eetuba of the Masubias.

Kwar of the Masaras.

Klabakila of the Basuto.

M'pelembe in the Chilala and
Chibisa countries.

Mtagaisi of the Swazis and Zulus. Oo-ka-mooh-we of the Makubas. Qualata of the Northern Bechuanas.

Qualata and Etsetla in the Lake Ngami country.

Qualata and Tseu in the Barotse country.

Tai-hait-sa of the Southern Bechuanas.

In spite of its larger size (height at shoulder about 4 feet 9 inches), the shorter horns and mane, the larger ears and eye-tufts, and, above all, the grizzled roan coat, render the present species a much less striking animal than its sable cousin. A marked character of the face of the roan antelope is the cutting-off of the white eye-stripe from the muzzle by a transverse dark bar connecting the dark nose-streak with the brown of the cheeks; while the dark nose-streak itself likewise stops short of the muzzle, which is thus wholly white.

Distribution.—From north of the Vaal and Orange Rivers through East and East Central Africa to the Sudan and Abyssinia, and westward to Angola, Nigeria, Gambia, and Senegambia. Recently the name H. rufopallidus has been applied to an antelope from East Africa, but it would seem unlikely that this can be anything more than a

local race of the present species. Gordon Cumming shot the roan antelope just north of the Orange River in Griqualand West, where it has for many years been exterminated. Nowhere abundant, the species is most plentiful in Mashonaland and neighbouring districts; in the Transvaal it is only sparsely distributed. South of the Orange River this group of antelopes was formerly represented by the much smaller blaubok, or blue antelope (H. leucophæus), long since exterminated.

Circum-	Tip to Tip.	Locality.	Owner.
	81	South Africa	A. Ohlsson.
	_		Major H. J. Goold-Adams, C.B.,
	•	3 ,	C.M.G. A. C. Fountaine.
			Julius Jeppe.
ŭ			F. C. Selous.
		•	Major A. St. H. Gibbons.
			Sir Edmund G. Loder, Bart.
			F. C. Selous.
	_		Major P. W. Forbes.
_	-	•	J. A. Jameson.
10	_		Mr. Justice Hopley.
10	_	Cul	Count T. Palffy.
7			F. C. Selous.
•		South Africa	G. Richards.
10		Portuguese East Africa	F. Vaughan Kirby.
87	-	J	Count E. Hoyos.
-	•	?	O. R. Dunell.
	_	Matabeleland	W. Van Ness.
•	,,,	Mashonaland	Hon. Walter Rothschild.
	10	Do	Sir John Willoughby, Bart.
9 1	4	Bahr-el-Salam, Sudan.	LieutCol. Hon. W. Coke.
• •	•	?	Cape Town Museum.
			F. V. Worthington.
8 §	121	C1!-	Dr. Percy Rendall.
	Ference. 91/2 9 81/2 91/2 91/2 81/2 10 7 91 10 81/2 10 7 91 10 81/2 10	ference. The form, ference. The first fig. 9\frac{1}{2}\$ 9\frac{1}{2}\$ 8\frac{1}{2}\$ 9\frac{1}{2}\$ 12 10 13\frac{1}{2}\$ 9\frac{1}{2}\$ 11\frac{1}{2}\$ 8\frac{1}{2}\$ 12\frac{1}{2}\$ 8\frac{1}{2}\$ 13 10 6\frac{1}{2}\$ 10 14\frac{1}{2}\$ 7 7\frac{1}{2}\$ 9 11\frac{1}{2}\$ 8\frac{1}{2}\$ 10 11\frac{1}{2}\$ 10 11\frac{1}{2}\$ 10 11\frac{1}{2}\$ 10 11\frac{1}{2}\$ 11\frac{1}{2}\$ 10 11\frac{1}{2}\$ 11\frac{1}{2}\$ 11\frac{1}{2}\$ 12 13 14\frac{1}{2}\$ 11\frac{1}{2}\$ 14 11\frac{1}{2}\$ 11\frac{1}{2	South Africa Sout

ROAN ANTELOPE (Hippotragus equinus)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
28§	9 3	9§	Mashonaland	F. C. Selous, British Museum.
$28\frac{1}{2}$	8 <u>1</u>	•••	Sudan	Col. Ralph Vivian.
28 1	8 <u>§</u>	9 1	Mashonaland	S. Chillingworth.
28]	94	128	Matabeleland	Hon. R. A. Ward.
- \$ 28 1	6 <u>1</u>	101	?	Dr. W. P. Y. Bainbrigge.
28	91	111	Upper Zambesia	M. C. Greaves-Bagshawe.
28	9	111	Lo Magondi's Country	Earl of Dunmore.
28	98	5	Nigeria	A. W. M. Brodie.
27 <mark>1</mark>	83	81	?	Sir Edmund G. Loder, Bart.
27 3	8 7	61	Karonga, B.C.A	C. C. Bowring.
27 3	93	8 3	Sudan	Hon. Walter Rothschild.
27 \$	91	9	?	J. Carr Saunders.
$27\frac{1}{2}$	9	$11\frac{1}{2}$	Matabeleland	G. H. M. Banks.
278	9	10	Barotseland	H. Timmins.
27 8	9₹	8 1	South Africa	A. Beit.
271	101	6 3	Matabeleland	Rev. Dr. R. J. Nevin.
27 1	9	10 <u>1</u>	Dahomey	F. C. Fuller.
-271	91	10 §	Noggara, Abyssinia .	J. Menges.
27	9	•••	?	Dr. W. P. Y. Bainbrigge.
♀ 27	7	8 <u>3</u>	British Central Africa.	J. E. Gough.
26 7	85	101	Gambia	Dr. Percy Rendall, British Museum.
26 <u>8</u>	83	5₹	Mashonaland .	J. G. Millais.
26 1	9 8	142	Matabeleland	Hon. John Ward.
26 <u>1</u>	9 7	12	Muchinga Plateau, Northern Rhodesia	F. Smitheman.
-26.3	8. 14	13.8	North of Kassala .	Prince Henry of Liechtenstein.
26	8 7	7 1	Gambia	The late Earl of Derby.
♀ 25 7	6 <u>3</u>	7	Mashonaland	F. C. Selous.
25 1	91/2	10½	Nigeria	Major A. J. Arnold, D.S.O.
♀ 25 1	6 7		Do	J. W. Carroll.
25 1	94	58	Sudan	Sir Edmund G. Loder, Bart.

ROAN ANTELOPE (Hippotragus equinus)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
25	8 1	20 3	Gambia		H. C. Goddard.
♀ 24½	6 <u>1</u>	108	Nigeria		A. W. M. Brodie.
24	91/2	14‡	Benguela		G. W. Penrice.
♀24	6 8	$6\frac{1}{2}$	Gambia		H. C. Goddard.
♀22 ‡	7	812	Songwe, B.C.A.		C. C. Bowring.
Ŷ 22 1	6 <u>3</u>	812	Sudan		Hon. Walter Rothschild.
22 lg	7 1	8	Do		W. D. James.
-22	9	9‡	Nigeria	•	B. R. M. Glossop.
21 7	7	9 1	Abyssinia		E. Lort-Phillips.
- 9 21½	•••		Nigeria		B. R. M. Glossop.
- 9 20	61	103	?		T. E. Buckley.
194	8	83	Nigeria		Lieutenant-Colonel T. D. Pilcher.
172	6 1	9‡	Abyssinia		British Museum.
I 24	7	7 1	British East Africa		S. L. Hinde.



Skull and Horns of Gemsbuck, from F. H. Barber's specimen.

GEMSBUCK (Oryx gazella).

The long, straight, spear-like horns, of which even the lion fights shy, render the gemsbuck and its more immediate allies an easily recognised sub-group. And even when, as in the white oryx, the horns are scimitar-shaped, they differ from those of the sable antelope by starting in the plane of the face. It is also a noteworthy fact that in

the present species the horns of the females are longer and finer, and therefore more prized, than those of the bulls. An adult gemsbuck stands about 3 feet 9 inches at the shoulder. In addition to the length of its horns, the species is sufficiently characterised by the presence of a tust of hair on the throat, and the cutting-off of the white eye-stripe from the muzzle by the union of the dark central nose-streak with the black of the cheeks.

Distribution.—The desert regions of South-Western Africa, from the northern Karoos of Cape Colony through the Kalahari and Damaraland to Southern Angola in Mossamedes and perhaps Benguela. North of the Chobe and eastwards of Khama's country the species appears to be unknown. So late as 1846 Gordon Cumming found gemsbuck abundant on the northern Karoos of Cape Colony; and even now a few linger on the plains to the south of the lower reaches of the Orange River. In the northern Kalahari, where they exist for months without water, they are still abundant. The splendid horns of the gemsbuck are always regarded as prime trophies of the hunter's skill; the finding, riding-down, and shooting of one of these wary and enduring desert-bred antelopes being a feat of which any man, however well mounted, may be deservedly proud.

Length on front.	Circum- ference.	Tip to Tip.	Locality.		Owner.	
47½	6 3	171	South Africa			The late J. S. Jameson.
- ♀ 45½			Bechuanaland			F. H. Barber. (See illustration.)
♀ 45 1	6 1	33 \$	Lake Ngami			Major H. J. Goold-Adams, C.B.,
45	81	18 <u>1</u>	Do.			C.M.G. Hon. Walter Rothschild.
-4418			South Africa			Dr. A. Schopf.
-44	61/2	17	?			A. Ohlsson.
-43 4	7	21	?			Dr. W. P. Y. Bainbrigge.
43 8	6 3	187	Nata River			F. C. Selous.
43 1	68	20 7	?			Sir Victor Brooke's Collection.
-43 1	8	16	?			James J. Harrison.
43 1 8	6 7		Botletli River			F. C. Selous.
43	7	18 3	South Africa			Rowland Ward.
-43	7 ½		?			Earl of Dartmouth.
43	6 1	22	?			W. Y. Campbell.

GEMSBUCK (Oryx gazella)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
42 8	7	14	?		A. Beit.
42½	7	20½	Bechuanaland		A. Neilson.
421	7		Do		Major H. J. Goold-Adams, C.B., C.M.G.
42	61	19½	Do		Capt. F. H. Lehmann.
-42 1	7	$23\frac{1}{2}$?		A. Ohlsson.
-42	6 §	22	?		Julius Jeppe.
42	6 1	20 3	?		A. Ryley.
$-41\frac{7}{16}$	8	15 8	Great Namaqualand .	•	Th. Rehbock.
411	7	22 3	?		C. D. Rudd.
9 41	61/2	22 <mark>3</mark>	?		Mr. Justice Hopley.
9 41 1	5 18	29 8	Great Namaqualand .	•	Th. Rehbock.
♀ 41	7	19	3.		Major R. Hayes-Sadler.
♀40 ¾	6 8	171	Cunene River	•	Capt. F. Cookson.
-40 2	•••		?		O. R. Dunell.
40 8	6 8	16 1	?		Lewis Atkinson.
-40 <u>1</u>	6 1	181	South Africa		R. C. Peake.
- 9 40 <u>1</u>	•••	•••	,		O. R. Dunell.
40	6 2	181	South Africa		Sir Edmund G. Loder, Bart.
- 9 40	•••		Do	•	Cape Town Museum.
- 9 39 8	6 1	151	Do		Julius Jeppe.
394	6 <u>8</u>	18	Bechuanaland	•	LieutCol. W. Sitwell.
39	71	22	?		R. A. Cooper.
39	61	16	South Africa	•	A. J. Forbes.
♀ 38 1	6₽	15 8	Kalahari	•	H. A. Bryden.
38	•••	183	Bechuanaland	•	A. C. Fountaine.
38	81/2	81/2	South Africa	•	F. E. Potter.
-38	7	18	Do	•	A. E. Pease.
-38		•••	?		C. T. Jones.
♀37	61	171	South Africa		Dr. W. P. Y. Bainbrigge.
♀ 36 2	6§	•••	North Bechuanaland .	•	F. C. Selous, British Museum.



Head of Beisa.

BEISA (Oryx beisa).

Beida of the Somalis. Sala of the Abyssinians of Danakil.

The beisa may be regarded as the north-eastern representative of the gemsbuck, from which it may be distinguished at a glance not only by the absence of a tuft of hair on the throat, but also by the separation of the black nose-stripe from the eye-stripes. There is also no black on the haunches or thighs, and the horns are considerably shorter and less divergent. Height at shoulder reaching to about 4 feet. Weight 458 lbs. (F. J. Jackson, C.B.).

Distribution.—North-East Africa, from Suakin through Abyssinia to Berbera in Somaliland, and south to Lake Baringo and the Equator.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
40		•••	?	Measured by A. O. Hume, C.B., at Aden.
♀ 39	51/2	8	Near Hargeisa, Somali- land	
371	7	11‡	Somaliland	G. D. E. Chapman.
36 \$	$6\frac{1}{2}$	63	Do	Prince Boris Czetwertynski.
- ♀ 36½			Northern Somaliland .	J. Menges.
36	7	6‡	Somaliland	G. D. E. Chapman.
36	6 <u>3</u>	74	East of Lake Rudolph	A. H. Neumann.

BEISA 273


BEISA (Oryx beisa)—continued.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
36	6 <u>₹</u>	101	50 miles from coast of	Capt. J. T. Brinkley.
36	6	81/2	Somaliland Somaliland	Count J. Potocki.
- ♀ 36			Do	Dr. Donaldson Smith.
354	6 <u>3</u>	9 1	Do	Lord Delamere.
♀ 35¥	5₹	8	Do	R. Wahrmann.
351	6 <u>1</u>	8 <u>\$</u>	Do	Prince Nicolas Ghika.
♀ 35₺	6	101	East of Lake Rudolph	A. H. Neumann.
35	6	9	Somaliland	Col. Arthur Paget.
♀ 35	51/2	7 \$	Danakil	Viscount Edmond de Poncins.
35	5 3	78	Somaliland	A. S. Trevor.
-35	6§	61	Do	J. Johnston-Stewart.
♀ 34¥	5 1	81/2	Do	G. M. Norrie.
♀ 34½	6	101/2	Do	Capt. J. M'Call Maxwell.
34½	6 2	83	Do	Alex. R. Alston.
34½	7	6	Do	Ford G. Barclay.
♀ 34½	5♣	83	Do	R. McD. Hawker.
♀ 34½	5븀	13	Do	T. W. H. Clarke.
34½	6 <u>8</u>	81	Do	E. Lee Townshend.
34½	5 1	8	Do	Sir H. D. Tichborne, Bart.
-34 8	6	104	Do	Count Scheibler.
-34 8	8 7	115	Do	T. W. H. Clarke.
-341	71/2	102	S.E. Somaliland .	S. Payne-Gallwey.
- ♀ 34 1 8	5₹	71/2	Burgo	P. H. G. Powell-Cotton.
34	6 1	74	Somaliland	E. N. Buxton.
34	5	9	Do	Lord Wolverton.
- ♀ 34	6	11	Do	S. Payne-Gallwey.
- ♀34	5	10	Do	A. E. Pease.
-33 ‡	6 1	8	Do	Sir Edmund G. Loder, Bart.
33♣	6	9	Do	Capt. C. S. Timins.
33 ‡	6	9	Do	Digby Davies.

T

BEISA (Oryx beisa)—continued.

Length on front.	Circum- ference.	Tip to Tip.	Locality.		Owner.
♀ 33 8	4 1	8	Somaliland		J. Arkcoll.
-33 1	6 1	88	Do		A. E. Leatham.
331	6 <u>3</u>	98	Do		A. H. Straker.
33∯	5 \$	81	Do		Major H. G. C. Swayne.
$33\frac{1}{2}$	5½	7 §	Do		Col. Arthur Paget.
$33\frac{1}{2}$	$6\frac{1}{2}$	10]	Do		G. M. Norrie.
$-33\frac{1}{2}$	6 1	8춁	Do		Count E. Hoyos.
- 9 33½	5½	$7\frac{1}{2}$	Do		Capt. M. M'Neill.
$-33\frac{1}{2}$	•••	•••	Do		J. Menges.
33#	48	7 8	Do		W. D. James.
33 ‡	6 1	9	Do		Major C. C. Ellis.
-331	5 2	54	Do		C. V. A. Peel.
33	6 §	7 1	Do		Capt. B. L. Carew.
♀ 33	5	6 3	Do		Capt. G. Campbell.
-33			Njemps, E. Africa	•	F. J. Jackson, C.B.
-33	7	111	Somaliland		LieutCol. J. W. H. Flanagan.
32 3	6 3	121	?		Lord Delamere.
32 3	71/2	88	Somaliland		Capt. F. C. Quicke.
32 3	6 1	81	Do		J. Byng-Paget.
32‡	6 <u>1</u>	111	Danakil		Viscount Edmond de Poncins.
♀ 32 ફ	5	8	Somaliland		Capt. C. H. Villiers.
$3^{2\frac{1}{2}}$	6	8	Do	•	P. B. Vander-Byl.
32½	6	7	Do		Count J. de Bylands.
32½	6 <u>1</u>	83	Do		Major H. C. Morland.
$-32\frac{1}{2}$	6 3	8 <u>1</u>	Do		W. W. Ashley.
$32\frac{1}{2}$	5‡	7 1 2	Do		A. Leslie Renton.
$-32\frac{1}{2}$	6 1	10 <u>1</u>	Do		Major G. Douglas.
32	61	61	Do		Lord Delamere.
♀ 328	5½	11	Do		Julius Jeppe.
32 1	$6\frac{1}{2}$	8 1	Do		B. Vincent.
32 1	6 <u>1</u>	9	Do		J. J. Richardson.
32	6 3	6	Do		T. W. Greenfield.
♀ 32	5 1	6	Do		E. T. Marshall.
32	6 <u>‡</u>	83	Do		G. H. Cheetham.
-311	47	114	Do.		Dr. Percy Rendall.
314	5 1	81/2	East Central Africa	•	
₽ 31	5 1	7	Somaliland	•	Major W. L. H. Paget.
31	5₹	$7\frac{1}{2}$	Do	•	B. R. M. Glossop.



Head of Fringe-eared Beisa.

FRINGE-EARED BEISA (Oryx callotis).

Distinguished from the ordinary beisa by the fringe of long hairs surmounting the ears, by the extension of the eye-stripe to the lower jaw, along which it runs to join the throat-stripe, by the absence of any black on the front of the legs below the knees, and by the rich fawn of the ground-colour of the upper part of the face. Height at shoulder, 48 inches.

Distribution.—East Africa, from Kilimanjaro and the Galla country to Masailand and the adjacent territory, probably not so far north as the Athi plains. Although considered a distinct species, it might perhaps be better to regard this animal as a well-marked local race of the ordinary beisa. It is generally found in bush-covered country, either singly or in small herds, and is of a remarkably shy disposition. The Swahili name is cheroa.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-3 2		•••	Athi River	R. Cator.
30½	5₹	101	East Africa	F. J. Jackson, C.B.
30 1		6	Sabaki River District.	T. W. Greenfield.
-30		10	East Africa	Sir John Willoughby, Bart.
29½	58	78	Do	Sir Robert Harvey, Bart.
29	6	61	Masailand	LieutCol. Hon. W. Coke.
29	51	1112	Do	Sir Edmund G. Loder, Bart.
283	5	6	Do	H. C. V. Hunter.

FRINGE-EARED BEISA (Oryx callotis)—continued.

Length on front.	Circum- ference.	Tip to Tip.	Loc	ality.			Owner.
♀ 285	4 3	124	Masailand				Henry Charrington.
27 3	6 7	10‡	Do.				F. J. Jackson, C.B.
27 1	6 <u>7</u>	81	Do.				Sir Robert Harvey, Bart.
27	6	9	Do.				E. Gedge.
27	7 1	78	Do.				H. C. V. Hunter.
26 3	7	9	Do.				J. Gardiner Muir.
ହ 26≩ୁ	51/2	10	Do.				Capt. R. A. J. Montgomerie, C.B., R.N.
26§	7	13	Do.		•		Sir Robert Harvey, Bart.
26 1	$6\frac{1}{2}$	9‡	Do.	•	•	•	C. Steuart Betton.
-24	7	81/2	Do.		•		Count Scheibler.
21	6₹	10	Do.			٠.	British Museum.

BEATRIX ORYX (Oryx beatrix).

This oryx is a much smaller animal than the beisa, measuring about 2 feet 8 inches at the shoulder, and is of a whitish colour, with a dark spot on the face, and a large dark patch on each cheek, which meets its fellow beneath the throat; the knees and the front of the lower portion of the legs being blackish brown, and the tail-tuft black. Distribution.—The interior of Arabia, especially the Nejd districts and the confines of the great desert south of Orman, and, it is said, the Bushire district at the head of the Persian Gulf. This antelope has very rarely been killed by British sportsmen.

Length on front.	Circum- ference.	Tip to Tip.	I	ocality.	•		Owner.
-26] }	44	10‡	Arabia		•		Paris Museum.
23 1	5	91	Do.				Sir Edmund G. Loder, Bart.
₽ 15	38	41/2	Head of	Persia	n Gu	lf.	B. T. Ffinch, British Museum.



Head of White Oryx.

WHITE ORYX (Oryx leucoryx).

A very distinct species of the genus, agreeing approximately in size with the beisa, but with long recurving scimitar-shaped horns and a generally whitish coloration, showing a more or less distinct chestnut tinge. The chestnut shows itself chiefly on the neck, shoulders, under parts, and upper portions of the limbs; and in addition to this there are six brownish patches or streaks on the face, two of which are situated in the middle line, while two form eye-stripes, the other pair being between the horns and the eyes.

Distribution.—North-Western Central Africa, from Sennar and Kordofan to parts of Nubia and the Sudan. Rare in collections and menageries.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
39₽	4 7	14 1	?	British Museum.
1 39	5‡		N. of Sokoto	Capt. P. S. Wilkinson.
37∄	5₹	71	?	British Museum.
358	5 1	11 1	?	J. Carr Saunders.
35	5 1	12	?	Sir Edmund G. Loder, Bart.
33 7	51/2	8	?	Hon. Walter Rothschild.
331	5	8‡	?	Duke of Bedford.
243	41	8‡	?	Sir Edmund G. Loder, Bart.

¹ Single horn.



Skull and Horns of mounted specimen of Addax presented to the British Museum by J. J. S. Whitaker.

ADDAX (Addax nasomaculatus).

This antelope is another member of the oryx group, but is referred to a genus apart, of which it is the sole representative. Its most distinctive features are the spirally twisted and closely ringed horns, which recall those of the Indian blackbuck, and the heavy mass of long hair clothing the neck and shoulders and forming a forelock on the forehead. The general colour is yellowish white in summer and grayish in winter, but the head, neck, and mane are brown, although a streak across the face below the eyes, the lips, and a spot on the outer surface of each ear are white. Hoofs very wide and shallow, almost like those of the reindeer. Height at shoulder about 3 feet 6 inches.

Distribution.—North Africa and Arabia.

ADDAX 279

ADDAX (Addax nasomaculatus)—continued.

Len	_	Circum-	Tip to	Locality.		Owner.
On front curve.	Straight line.	ference.	Tip.	ixtanty.		Owner.
38½	30½	61/2	$12\frac{1}{2}$	South Tunisia.		J. J. S. Whitaker, British
37 2	314	54	$23\frac{1}{2}$	Sahara		Museum. (See illustration.) W. Barry.
35₹	28	6 3	134	North Africa .		Hon. Walter Rothschild.
35	29	6 <u>1</u>	14	Do		W. Barry.
34½	27	61/2	171	South Tunisia .		J. J. S. Whitaker.
-341	273	6 <u>1</u>	17	Do. .		A. E. Pease.
33 2	28 3	5 7	22	Do		Sir Edmund G. Loder, Bart.
33½	27 1	61/2	174	Do. .		J. J. S. Whitaker.
$33\frac{1}{2}$	26	5½	121	Do		Hon. R. A. Ward.
321	26 <u>l</u>	61	131/2	Do		Rowland Ward.
315	261	6	173	Do		F. H. Barber.
31	25½	5‡	13	Do		J. H. Thomas.
-3 1	28	•••	•••	Do		Sir H. H. Johnston, K.C.B.
♀ 30 1	271	43	174	South Algeria		Hon. Walter Rothschild.
- ♀ 30½	$27\frac{1}{2}$	••		South Tunisia		Sir H. H. Johnston, K.C.B.
29½	24 ³	6‡	14	Sahara		Hon. Walter Rothschild.
29 1	255	5‡	19½	South Algeria		Julius Jeppe.
283	24 7	5½	12	Do		British Museum.



Skull of Male Nilgai.

NILGAI (Boselaphus tragocamelus).

This somewhat ungainly and decidedly small-horned antelope is the single Oriental representative of an important group whose other members are African. Most are of large size, and, with the exception of the elands, lack horns in the females. In the males the horns are angulated in front and generally spirally twisted, but in no case ridged. There are small face-glands, the muzzle is naked, and the tail is long and tufted. Except in the nilgai, the cheek-teeth have short and broad crowns.

From the other members of the group the nilgai is readily distinguished by its short, upright horns, which, although angulated in front, show no distinct spiral twist. With a long and pointed head, this antelope has the fore-legs considerably longer than the hinder pair; and it is chiefly to this peculiarity that its ungainly appearance is due. Both sexes have a mane on the neck, but the bulls alone possess a tuft of long hair on the middle of the throat. The general colour of the adult bull is dark gray, tinged with blue or brown; but the mane and tufts of long hair are black, and streaks and patches on the face, ears, and throat, as well as the under parts, the lower surface

NILGAI 281

of the tail, and a ring above and below each fetlock, are white. Height at shoulder from 4 feet 4 inches to 4 feet 8 inches. Mr. A. O. Hume shot a specimen in the Aligurh district in 1855 whose horns measured $11\frac{3}{4}$ along the front curve, and had a circumference of 9.5. They were unfortunately destroyed in the Mutiny.

Distribution.—The peninsula of India, from the foot of the Himalaya to the south of Mysore; common in parts of the Eastern Punjab, the North-West Provinces, Guzerat, and the Central Provinces.

Length on front.	Circum- ference.	Tip to Tip.	Locality.	Owner.
9 ⁷ 8	7 3	5 1 2	?	Sir Edmund G. Loder, Bart.
91/2	7½	6 1	3	J. Whitaker.
-9 1	8 <u>1</u>		Bhurtpur .	Capt. E. R. Gordon.
9‡	63	4 7	?	British Museum.
9	7	9	?	J. Whitaker.
9	6∯	3½	?	A. O. Hume, C.B.
8 <u>7</u>	6 <u>1</u>	51	Central Provinces	C. D. Twopeny.
83	6‡	5 \$	Bhurtpur .	Major J. M. Fawcett.
-83			Jumna Valley .	Indian Museum.
81/2	$7\frac{1}{2}$	47	Central Provinces	Capt. E. H. R. Hibbert.
$8\frac{1}{2}$	5½	41	Do	Marquis of Dufferin and Ava.
$-8\frac{1}{2}$	6 <u>1</u>	6	Do	Dublin Museum.
81/2	7 ½	5 1	Do	Dr. W. P. Y. Bainbrigge.
$-8\frac{1}{2}$	6	5 1	Do	Count J. Potocki.
-8 8	7 1	$6\frac{1}{2}$	Do	Dr. Percy Rendall.
8‡	7 l	7³	North India .	Sir Victor Brooke's Collection.
-8.25	7·8o	6.70	Ulwar	Viscount Edmond de Poncins.
81	•••		Kathiawar .	LieutCol. L. L. Fenton.
-81.	•••		Kota	H.H. Maharaja of Bikanir.
8 <u>1</u>	6 1	5	?	Hume Collection, British Museum.
8	6 3	6‡	Nimar, Cent. Prov.	LieutCol. H. Wade-Dalton.
7 1	7 ½	5 \$?	H. G. Buxton.
7₹	7	6	Central Provinces	Major A. Colville.
78	6	7 1	?	L. M. Le Champion.



Head of Bushbuck shot by A. M. Naylor.

BUSHBUCK (Tragelaphus scriptus).

Assali, Danakil name.

Bawala in the Chilala and Chibisa countries.

Boschbok of the Dutch.

Dol of the Somalis.

Ibawara of the Lower Zambesi natives.

Imbabala of the Swazis and Matonga.

Inkonka (male), Imbabala (female) of the Zulus.

M'babala in the Barotse country.

M'babala and Serolo buchuhu in the Lake Ngami country.

Mbawara of the Swahilis.

Scrolobutuku of the Bamangwatos.

Ungurungu of the Makubas.

The bushbucks and harnessed antelopes form an extensive group of species nearly allied to the kudus, but usually displaying great sexual differences in the colour of the coat, and generally having a simpler spiral to the horns. The females are nearly always striped with white on a chestnut ground, but the bucks may be darker and more uniformly coloured. As in the kudus, the females are hornless. The true or lesser bushbuck is the smallest and at the same time the most widely spread member of the group, having several local races. The height at the shoulder ranges from $2\frac{1}{2}$ to 3 feet, and the weight from 100 lbs. to 170 lbs. In the Abyssinian bushbuck (T. scriptus decula), which ranges

into the dense forests bordering the Webbe River in Somaliland, and is locally known as dol, the build is low and stout, and the general colour yellowish, the light stripes being nearly obsolete. In the West African bushbuck (T. scriptus typicus), from West, Central, and South-Central Africa, the ground-colour is bright rufous, and the spots and stripes are very conspicuous. On the other hand, in the East African race (T. scriptus roualeyni) the bucks are dark brown, with only faint indications of white markings. Finally, in the Cape bushbuck (T. scriptus sylvaticus) the colour is dark brown without transverse white stripes, and the spots reduced to a few indistinct ones on the haunches.

In the southern districts of Cape Colony bushbuck are still to be found in abundance, even in the neighbourhood of large towns like Port Elizabeth. Recourse is, however, had to preserving for the greater part of the year, by which means the well-known Easter Hunts, when driving is practised and large bags are made, are still kept up. In more inland districts either the does or both sexes are periodically protected. A wounded bushbuck will often make a desperate and dangerous charge.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
-1918	54	11	Lower Zambesia .	•	Surgeon W. H. S. Stalkartt, R.N.
177	61	5₹	East Africa		Sir John Kirk, K.C.B.
$-17\frac{1}{2}$	$6\frac{1}{2}$	78	Mount Zomba, B.C.A.		D. MacAlpine.
174	6	5‡	N.E. Gazaland.		A. M. Naylor. (See illustration.)
17	5½	8 <u>1</u>	Mount Zomba, B.C.A.		C. C. Bowring.
16 §	5 7	61	Manda Island, B.E.A.		British Museum.
16 1	5 \$	(one horn)	Pungwe		Col. G. A. Percy.
-16 1	5½	5½	?		Julius Jeppe.
-16 1		•••	South Africa		A. Bowker, Grahamstown Museum.
-16½	•••	•••	?		O. R. Dunell.
-16 1	61/2	7 1/2	Natal		General A. W. Drayson.
-16	61/2		Kalamba Hills		F. Vaughan Kirby.
153	6	58	Okavango River .		Major H. J. Goold-Adams, C.B., C.M.G.
154	6	81	Zululand		A. H. Neumann.
15%	5	34	Do		Capt. L. O. Williams.
-15 3		•••	South Africa		F. H. Barber.
155	5½	3	British East Africa .		G. E. Smith.

BUSHBUCK (Tragelaphus scriptus)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	_	Owner.
15§	5‡	5	Pungwe		lion. T. Thynne.
151/2	5½	3∰	Transvaal		C. F. Eustace.
151	58	41/2	South Africa		F. C. Selous.
-151			Do		H. T. and A. H. Glynn.
15 8	68	$7\frac{1}{2}$	Do		Major-General Sir Arthur Ellis,
1 5 g	5½	7 l	Zululand		K.C.V.O. Hon. Charles Ellis.
154	6	6 3	Pungwe		H. R. Holden.
-154	7 1	6 <u>1</u>	Nyasaland	•	Alex. R. Alston.
15 <u>1</u>	5½	81/2	. ?		Mr. Justice Hopley.
142	58	6 <u>3</u>	Shiré River, B.C.A		H. H. Williams.
144	5‡	6 <u>1</u>	Zululand		LieutCol. Hon. W. Coke.
144	5	$6\frac{1}{2}$	South Africa	•	R. A. Cooper.
-144	5 3	34	Nyasaland	•	Alex. R. Alston.
148	5 2	23	Matabeleland		G. H. M. Banks.
148	$5\frac{1}{2}$	7	Zambesia		G. Richards.
$-14\frac{1}{2}$			Cape Colony		F. II. Barber.
-141			Transvaal		C. T. Jones.
-14 1	5	7	Pungwe		Julius Jeppe.
14			East Africa		F. J. Jackson, C.B.
14	57	54	Matabeleland		H. and C. Beddington.
14	5 7	61	South Africa	•	C. D. Rudd.
14	5 1	10]	Pungwe		G. Micklethwait.
14	5‡	7	Ngamiland		Capt. E. J. Lugard.
-14	6	$7\frac{1}{2}$	South Africa		James J. Harrison.
137	53	5½	Do		H.R.H. the Duke of Saxe-Coburg and Gotha.
13 3	5 3	6 <u>1</u>	Do		Hon. Walter Rothschild.
133	51/2	58	Do		Sir Edmund G. Loder, Bart.
-134	6‡	6	Natal		T. E. Buckley.
138	5½	5½	East Africa		H. C. V. Hunter.
13 8	58	48	?		Sir Victor Brooke's Collection.

BUSHBUCK (Tragelaphus scriptus)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
$13\frac{1}{2}$	51/2	7	South Africa .			F. C. Selous.
131	51	5 8	Do	• .		British Museum.
$13\frac{1}{2}$	5 1	48	Do			Sir John Willoughby, Bart.
-138	5 7	7 1	North of Pungwe	•		Count E. Hoyos.
-134		•••	Cape Colony .			C. T. Jones.
13	6 1	5 1	East Africa .			F. J. Jackson, C.B.
13	51/2	7	Benguela			G. W. Penrice.
13	5 1	5	Pungwe . :			G. L. Bonham.
13	54	5 ½	British Central Afric	a		J. E. Gough.
-13	•••	•••	Zambesia		•	C. T. Jones.
12 <u>3</u>	5	41/2	South Africa .			A. E. Capell.
-12 <u>8</u>	54	8 1	Barberton			Dr. Percy Rendall.
12‡	48	5 1	3			Dr. W. P. Y. Bainbrigge.
128	5 à	48	Chobe Valley .			F. C. Selous, British Museum.
121	61	7 1	South Africa .			E. G. Christian.
-I2 1	5	51	Matabeleland .		•	J. Brander Dunbar.
I 14	58	4 1	Pungwe			Hon. T. Thynne.
112	5 1	41/2	Uganda	•		The late Capt. Speke, British Museum.
118	5	6 8	East Africa .			Sir Robert Harvey, Bart.
-11 8		•••	Adda Bush .	•		Cape Town Museum.
102	41/2	5	Nigeria			Major A. J. Arnold, D.S.O.
101	41/2	21	Gambia	•		A late Earl of Derby, British Museum.
-9 ₿	41/2	6 1	Songwe, Nyasaland			John Yule.



Head of Abyssinian Bushbuck shot in the Webbe Shebayle River District, Somaliland, by Major H. G. C. Swayne.

The following specimens belong to the Abyssinian race (*T. scriptus decula*); the body measurements being those of one shot by Viscount Edmond de Poncins on the Hawash River, 1898:—

From nose to root of tail	48\frac{3}{6} ins.
Tail (hair included) .	10 3 "
Height at withers	$26\frac{\bar{3}}{4}$ "
Round the body	30 § "

	_						J-g "
Length on front curve.	Circum- ference.	Tip to Tip.	Localit	y.			Owner.
-17			Somaliland				Major H. G. C. Swayne.
-16 1	6 1	5	Do.				A. H. Straker.
16	6	7±	Do.				Major H. G. Swayne.
14 1	6 7	5	Do.				A. H. Straker.
$-13\frac{3}{4}$	$5\frac{1}{2}$	5 1	Harar, Abyssini	a			Prince de Lucinge.
-13	$5\frac{1}{2}$	5‡ 6դ8	Do.				Do.
- 12 7	5½ 5½	34	Near Hawash R sinia	liver.	Ab;	ys-	Viscount Edmond de Poncins.
125	5	51	Abyssinia .				W. D. James.
12	51	5 1 5 1 5	Do				British Museum.
II		5	Settite River, A	byssi	nia		Col. Ralph Vivian.
11	5분 4분	2 §	Northern Abyss	inia	•		British Museum.




Horns of Male Bongo, from the type specimen in the British Museum.

BONGO or BROAD-HORNED ANTELOPE (Tragelaphus euryceros).

This West African species is the largest of the harnessed antelopes. In addition to its large size, it is characterised by the bright chestnut colour of the males, marked with a number of narrow transverse white stripes, the shortness of the hair, the absence of a throat-fringe, and the smooth and stout horns, of which the tips become yellow by wear. The markings on the face take the form of a pair of white spots below the eyes, and there is a white crescent on the breast. Height at the shoulder probably about 4 feet.

Distribution.—West Africa, from Liberia, through Fanti to the Ashkankolu Mountains and the Gaboon. It is doubtful if this rare and beautiful antelope has ever been killed by British sportsmen. The original specimens were obtained by P. Du Chaillu, and are in the British Museum.

Len On front curve.	gth. Straight.	Circum- ference.	Tip to Tip.	Locality.	Owner.
	271	9 1	16 1	Gaboon	Sir Edmund G. Loder, Bart.
32‡ 31‡ 30‡	26 1 24 1	9 ‡ ₹0 ₹	12 12	West Africa Do	Sir Victor Brooke's Collection. Do.
30	24 l	9₹	111	Ashkankolu Mountains	British Museum.
29½	25 8	94	10g	Gaboon	P. Du Chaillu, British Museum.
27 중	23 3	10]	10 \$	Fanti	Do.



Horns of Nyala, from a specimen in the possession of F. H. Barber.

NYALA (Tragelaphus angasi).

The next African representative of the larger harnessed antelopes is a smaller and more delicately built animal than the bongo, standing about 3 feet 6 inches at the shoulder. Weight about 250 lbs. to 300 lbs. In both sexes the hair is very long and coarse, but in the male its colour is dark grayish brown with a small number of indistinct white stripes, while in the female it is bright reddish chestnut with clearly defined stripes. The males have a fringe of long hair on the neck and the under parts of the body, their horns being much rougher than those of the bongo. As in the latter, the hoofs are short.

Distribution.—South-East Africa, including Zululand, Delagoa Bay, and Nyasaland; on the West Coast it has been reported from Angola, although this form may indicate a distinct local race.

Length.		Circum-	Tip to	Locality.	Owner.	
		ference.	Tip to Tip.	Locality.		
31 1	24½	8‡	$3\frac{1}{2}$	Shiré River, B. C. A.	Fergus Maclagan.	
-294	24		1112	?	F. J. Newnham.	
-29 1	•••	•••		Delagoa Bay	F. H. Barber. (See illustration.)	

NYALA 289

NYALA (Tragelaphus angasi)—continued.

Length.		Circum- Ti	Tip to	T andies	0.000	
On front curve.	Straight.	ference.	Tìp.	Locality.	Owner.	
-29 1				?	O. R. Dunell.	
28 1	23 3	8	124	Zululand	Hon. Walter Rothschild.	
-28½	•••			?	W. Russell Bowker.	
$-28\frac{1}{2}$		•••		Delagoa Bay	F. H. Barber.	
$-28\frac{1}{2}$		•••		Do	Prince Boris Czetwertynski.	
28	24	8	15	Katanga, B.C.A	John Yule.	
28	23	7 1	134	Near Chiromo, B.C.A.	Staff-Surgeon J. Dowson, R.N.	
28		8 1	10	South Africa	British Museum.	
-28	241	7 ½	112	Delagoa Bay	C. T. Jones.	
-28	23 \$	8	11	?	Dr. W. P. Y. Bainbrigge.	
27 🖁	22½	7 1	91	South Africa	Sir Victor Brooke's Collection.	
27 §	23	7 8	8‡	Do	Earl of Dunmore.	
-27½	23	81	13	Do	E. D. Scott.	
271	22 <u>1</u>	7∰	81	3	C. D. Rudd.	
27½	23½	71/2	15	Delagoa Bay	Major H. J. Goold-Adams, C.B., C.M.G.	
278	•••	8 1	10 1	South Africa	Sir Edmund G. Loder, Bart.	
-271	•••	7 1	10 §	?	Julius Jeppe.	
-27 l s	23½	•••	14	Delagoa Bay	H. T. and A. H. Glynn.	
27		8	91	Shiré River, B.C.A.	F. Vaughan Kirby.	
27	225	72	111	3	J. R. Buckler.	
26 ‡	221	$7\frac{1}{2}$	11	3	F. E. Potter.	
26 3	228	7	113	Amatongaland .	H. A. Bryden.	
26 1	22	7 1	7	Delagoa Bay	Frank Harris.	
-26½	22½	7 §	6 g	Do	Dr. Percy Rendall.	
26 1	22	6 7	ΙΙ§	Do	F. C. Selous.	
26	•••	7 8	87	Zululand	Hon. Charles Ellis.	
25	21	8	98	?	Capt. A. M'Lean Wait.	
-25	22½	8	124	3	A. Ohlsson.	
24½	20 <u>1</u>		. •••	Amatongaland .	Cape Town Museum.	
238	20	7 5	115	Zululand	Durban Museum.	
23 1	201	7 1	71/2	Do	G. Richards.	
23 1	19	6 ₹	7	Do	A. H. Neumann.	
23	•••	7	5₹	Do	British Museum.	
-22 1	19½	7	9 1	Delagoa Bay	Count E. Hoyos.	
-22	•••			St. Lucia Bay	G. F. Angas.	
21	19	9	•••	Do	LieutCol. Hon. W. Coke.	

U



Horns of West African Harnessed Antelope, from the Gambia.

WEST AFRICAN HARNESSED ANTELOPE (Tragelaphus gratus).

Together with its near ally the situtunga, this species differs from all the other members of the group by the extreme elongation of the hoofs, which are thus adapted for supporting the weight of the body on the spongy soil of the marshes in which these antelopes dwell. The lateral hoofs, too, are much more developed than in other harnessed antelopes. The general coloration of the West African species is very similar to that of the nyala, the ground-colour of the coat of the male being olive, and that of the female bright rufous, marked in both sexes with white stripes on the body and spots on the face. There is, however, no fringe of long hair on the throat. Height

at shoulder about 3 feet 7 inches. Both in this species and the situtunga the horns of the males are longer and more twisted than in the other members of the genus, and thus come very close to those of the kudus. In the situtunga the coloration of the adult is uniform grayish brown. The West African species ranges from the Congo to the Gaboon and Cameroon districts. It has but seldom been collected by sportsmen, so that very little is known of its habits in the wild state. Several specimens have been bred in captivity at the Zoological Gardens, Amsterdam.

Length.		C '	T:						
On front curve.	Straight line.	Circum- ference.	Tip to Tip.	Locality.			Owner.		
30	26 <u>‡</u>	7 3	148		?			Rowland Ward.	
29½	$24\frac{1}{2}$	81	14	Gaboon			•	Sir Edmund G. Loder, Bart.	
$28\frac{1}{2}$	23	8	8	Gambia	•			Rowland Ward. (See illustra-	
25‡	211	6₹	121	Do.	•			Guy II. Sangster.	
$25\frac{1}{2}$	22	8	•••	French Co	ongo			Hon. Walter Rothschild.	
-24 11	211	7흥	6 <u>₹</u>	West Afri	ca			Dr. Percy Rendall.	
$-23\frac{1}{2}$	20 l	8 <u>1</u>	91 ¹ 8	Do.				Do.	
•••	19		•••	Gaboon				British Museum.	
$23\frac{1}{2}$	18	$7\frac{7}{8}$	113	. Do.				Sir Victor Brooke's Collection.	
23	20½	6 7	9 <u>1</u>	West Afri	ca			Hon. Walter Rothschild.	
22 4	19½	7.	10 3	Nigeria				J. A. Burdon.	
228	193	7	111	Gambia				Rowland Ward.	
-21.5		7.8			?			Berlin Museum.	
-17	141 ³ 6	6 §	•••	Ogooné				Paris Museum.	



Frontlet and Horns of Situtunga, from John Yule's specimen.

SITUTUNGA (Tragelaphus spekei).

Situtunga in the Barotse country. Zowè in the Chilala and Chibisa countries.

Nakong of the Batauwani at Lake Ngami.

Situtunga, Puvula, Unzuzu of the tribes on the Chobe and Central Zambesi.

N'zoé of the natives of Lakanga River north of the Zambesi.

This species, also known as the nakong, has the same habits and make as the West African harnessed antelope, from which it differs by its uniformly grayish-brown colour; the young alone being faintly barred and spotted. Another peculiarity is to be found in the length and

silkiness of the hair. The horns, which are nearly smooth and strongly keeled, form nearly two complete turns, and thus approximate to those of the kudu.

Distribution.—The situtunga is an inhabitant of the dense reed-swamps bordering the rivers of Central, South-Central and East Africa; and is semi-aquatic in its habits, frequently burying itself up to the eyes in the water. Consequently it is one of the most difficult of all antelopes to kill; and has even baffled the energy of Mr. Selous. By firing the reed-beds in the dry season, the natives are able to spear the situtunga as they cross open water. At night these antelopes leave the reed-brakes for the islands in the rivers, but before dawn return to their impenetrable covert. Mr. A. B. Phipps, in a letter dated October 1895, states that they have become very rare in the swamps bordering the Okavango River, owing to that river having changed its course and ceased to flow into the Botletli. Consequently they go down to the latter for water, and are shot. A few are found on a bush-clad rocky island far out in the Victoria Nyanza; in which neighbourhood the species was first discovered by the late Captain Speke.

Length.					
On front curve.	Straight line.	Circum- ference.	Tip to Tip.	Locality.	Owner.
35	28	7½	1 5 ½	South end of Lake Tanganyika	John Yule. (See illustration.)
-33½	27 §	7 1	19	?	Julius Jeppe.
33 1	26]	8	16§	Okavango Valley .	Major H. J. Goold-Adams,
-33	•••		•••	?	C.B., C.M.G. Alfred Ebden.
32	27	71/2	16 8	Chobe Valley	F. C. Selous.
-32½		•••	,	?	C. T. Jones.
31 7	25 4	81	171	Near Linyanti	F. C. Selous.
3112	24 8	7	16 1	Chobe Valley	F. C. Selous, British Museum.
-311	25 1	$7\frac{1}{2}$	13‡	Bangweolo	F. Smitheman.
-311	2510	7 1	14 8	Congoland ·	S. de Brazza, Paris Museum.
31	23 1	8	9 ₹	3	G. Richards.
-30½	•••	•••		?	O. R. Dunell.
-30½	$23\frac{1}{2}$	812	151	?	Julius Jeppe.
-30	25	78	18	?	Mr. Justice Hopley.
29 <mark>3</mark>	25	83	17	? .	R. A. Cooper.

SITUTUNGA (Tragelaphus spekei)—continued.

Len	gth.				
On front curve.	Straight line.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-29½	26	7 3	16	?	A. Ohlsson.
-28 3	26}	78	148	Lake Mweru, B.C.A.	R. H. Ferrers Stranack.
271	23 1	7 1 8	10	B.C.A	Rowland Ward.
271	$21\frac{1}{2}$	8	6	Barotseland	H. Timmins.
27 1	228	7 5	14 7	Lake Mweru, B.C.A.	Rowland Ward.
26 3	213	71	113	?	W. Van Ness.
26½	213	7 1	17	Chobe Valley	Frank Harris.
2 6	22§	7	171	?	Sir Victor Brooke's Collection.
	25			Okavango Valley .	A. B. Phipps.
25 7	221	7‡	198	Chobe Valley	F. C. Selous, British Museum.
25½	21 ³ / ₄	7	138	Bangweolo	F. Smitheman.
25	21	8	16 1	Botletli Valley	Hon. Walter Rothschild.
24	20	71/2	111	Barotseland	E. D. Scott.
23§		7‡	10	?	Major H. J. Goold-Adams,
23	20 1	•••	5 7	?	C.B., C.M.G. F. H. Barber.
22 ³	19½	10	7 ½	?	Hon. Walter Rothschild.
20 ³	18 1	6‡	10	Benguela	G. W. Penrice.
20 ³	17‡	6‡	5	Do	Do.
198	16 7	6 8	91	Barotseland	R. T. Coryndon.
-171	•••	54	7	Victoria Nyanza	E. Gedge.
158	141	5#	83	Do	Hon. Walter Rothschild.



Head of Male Greater Kudu.

GREATER KUDU (Strepsiceros kudu).

Eebala-bala of the Amandebele. Ngoma in the Chilala and Chibisa Ee-zilarwa of the Makalakas. countries. Dwar of the Masaras. Noro of the Mashonas. Godir of the Somalis. Tolo of the Bechuanas. Itolo of the Basutos. Tolo in the Barotse and Lake Ngami countries. Itshongonons of the Swazis. Muzeeloua of the Batongas. Unza of the Mazubias.

Unzwa of the Makubas.

A male shot by Dr. Percy Rendall, in Nyasaland, measured:

	inches.			inches.
Nose to tail	91 1	Girth of barrel		. 72
Height at shoulder .	55	" before hips		. 54
Point of shoulder to nose	32	" of fore-leg	•	$14\frac{3}{4}$
Length of tail	$17\frac{1}{2}$	" "thigh		$19\frac{1}{2}$
Girth of neck (min.)	30	" " neck (ma	x.)	. 43

Although rather less brilliantly coloured than some of the harnessed antelopes, the kudus are among the handsomest of all antelopes, their spiral horns, striped coat, and noble carriage rendering them really magnificent creatures. Their chief difference from the bushbucks is to be found in the fuller spiral formed by the horns: both sexes being nearly similar in colour. The special characteristics of the greater or true kudu are the large size (height at shoulder reaching to 4 feet 10 inches or 5 feet), the presence of a thick fringe of long hair on the throat, and the open spiral of the horns of the bull. The colour is too well known to require description.

Distribution.—The kudu, in suitable localities, ranges over the greater part of Africa south of the Sahara, extending from Abyssinia and Somaliland through East and Central Africa to the Cape, and westward across the continent to Angola, where the Congo apparently forms its northern limits. In spite of its bulk, it is an adept at concealment; and this trait, coupled with its general wariness and acute sense of smell and hearing, has largely contributed to its survival in districts where it is much hunted. Except in the Uitenhage jungles, where it is preserved by English farmers, the kudu has, however, been exterminated from Cape Colony. Mashonaland it is still abundant, as it is in the highlands of Somaliland, in which country it is rarely met with on the plains.

Although an adept at getting across rocky hills, this animal is by no means a good performer on the flat (where it seldom allows itself to be surprised), and can be ridden down without much difficulty by a fairly well-mounted hunter.


Horned females occur very rarely; but three specimens with horns are recorded on p. 302.

Leng	gth.		. оп р	J		
On outside curve.	-	Circum- ference.	Tip to Tip.	Locality.		Owner.
	48 7			Ngamiland		F. H. Barber. (See illus-
63	48 1	$12\frac{1}{2}$	49	Macloutsie River .		tration.) E. W. Tompson.
-63	44½		42 8	Nr. Tete, Zambesi.	•	Major P. W. Forbes.
-63	39	101	12	3		Mr. Justice Hopley.
	45½		•••	Delagoa Bay	•	F. H. Barber.
61 <u>1</u>	-45 1		•••	South Africa		O. R. Dunell.
60 <u>3</u>	441	12	29	Do		Frank Harris.
60∰	458	1112	33	Macloutsie River .		F. C. Selous.
-6 0₿	44	•••	31	South Africa		H. T. and A. H. Glynn.
-60½	47	101	44 ²	Do		A. Ohlsson.
-60	3 9	10	7 §	Lebombo Mountains		F. Vaughan Kirby.
-594	4518		39 8	Pungwe Valley .		Major A. St. H. Gibbons.
59 1	421	II§	19	?		G. Richards.
-58‡	46 1	1112	412	Transvaal		F. Van Zeller.
-58 1	45		46	?		O. R. Dunell.
-58½	45	1112	44 ³	3		Mr. Justice Hopley.
-58 1	46	10	39 2	South Africa	•	Julius Jeppe.
58	46	115	39	S.E. Mashonaland.		J. G. Millais.
-58	46	•••	•••	Zomba, B.C.A		D. MacAlpine.
58	431/2	101	$32\frac{1}{2}$	Okavango River .		Major H. J. Goold-Adams, C.B., C.M.G.
-58	44	11	44	Lebombo Mountains	. '	
-58	41		•••	South Africa		Cape Town Museum.
57 1	41	112	21½	N.E. Transvaal .		A. M. Naylor.
-57½	39 \$	117	15	South Africa		Julius Jeppe.
-57 8	39	98	32	Somaliland		J. Johnston-Stewart.
571	458	103	35	B. C. Africa		H. C. Macdonald.

Len	gth.	Circum- Tip to	mi			
On outside curve.	Straight line.	ference.	Tip to	Locality.		Owner.
57‡	44 8	112	348	South Africa		Sir Edmund G. Loder, Bart.
57 l	43	ΙΙ∯	34 3	Mashonaland	•	H. and C. Beddington.
57	43	14	32	South Africa		A. Moseley.
57	42‡	11	31.	Do		Capt. M. D. Graham.
57	39½	9§	36 <u>1</u>	Somaliland		H.R.H. le Duc d'Orléans.
56 1	431	111	26½	Matabeleland .		W. Van Ness.
56 1	42 §	10 8	382	?		Sir Victor Brooke's Collec-
56 <u>1</u>	41½	101	23 1	Matabeleland .		Hon. C. Greville.
56 <u>1</u>	383	91	28 3	Somaliland		J. Benett-Stanford.
56 1	35	10 1	30	Do		Sir Edmund G. Loder, Bart.
56 1	441	118	33 1	South Africa		Hon. Walter Rothschild.
56	43	103	43	Mashonaland .		F. C. Selous.
56	44	103	39	South Africa		F. B. Dunsford.
554	41	103	36	Do		F. Struben.
55½	39 1	103	26½	Mashonaland .		A. Neilson.
55½	$42\frac{1}{2}$	1112	37	Matabeleland .		Lord Brackley.
55½	421	101	38	Do		Major James Grant.
55 1	401	11	27‡	E. C. Africa		H. S. H. Cavendish.
55	411	112	28	Matabeleland .		Major R. Hayes-Sadler.
55	40½	121	321	Zululand		C. D. Rudd.
-55	41	10	41	?		The Maclaine of Lochbuie.
541	39 1	101	$27\frac{1}{2}$	Upper Shiré, B.C.A.		C. C. Bowring.
541	40½	11	$21\frac{1}{2}$	Mashonaland .		S. Chillingworth.
-54½	39	11	$27\frac{1}{2}$	South Africa		Dublin Museum.
- 54½	38 1	11	26	Do		Durban Museum.
- 54½	32½		•••	N. Somaliland .		J. Menges.
541	40 1	101	39½	Zululand		C. H. Akroyd.
- 54 1	38 7	101	22§	Somaliland		P. H. G. Powell-Cotton.
54	40	101	32	Do		Captain C. S. Timins.

GREATER KUDU (Strepsiceros kudu)—continued.

Length.		C'	TP:- 4-						
On outside curve.	Straight line.	Circum- ference.	Tip to Tip.	Locality.			Owner.		
54	41	11	37 3	?			Sir Edmund G. Loder, Bart.		
54	42	103	354	Zululand .			A. Cameron.		



Frontlet and Horns of Greater Kudu, from F. H. Barber's specimen.

54	42	114	39 3	Mashonaland		Earl of Dunmore.
-54	40½	101	314	South Africa .		Dr. W. P. Y. Bainbrigge.
54	40	114	24	Do.		LieutCol. W. Sitwell.
-54	39	101	32	Somaliland .		Norman B. Smith.
-53 ²	411	101	$32\frac{1}{2}$	S.W. Somaliland		C. V. A. Peel.

Length.		C'	~				
On outside curve.	Straight line.	Circum- ference.	Tip to Tip.	Locality.		Owner.	
53₩	41	10	25	Mashonaland .		J. A. Jameson.	
- 53½	36 <u>1</u>	91	23½	Somaliland		J. Brander Dunbar.	
53½	39	101	31‡	Zululand		LieutCol. Hon. W. Coke.	
53 1	40½	111	40½	Lebombo Mountains		Count E. Hoyos.	
53½	37	9§	23	Somaliland		E. Lort-Phillips.	
53	411	101	28‡	South Africa		W. Y. Campbell.	
•••	40.7	8.8	30.9	Bahr Setit		Prince Henry of Liechten-	
52 1	40	10]	37	Delagoa Bay		stein. Sir John Kirk, K.C.B.	
52 3	39 8	9 4	25 <u>Ř</u>	Mashonaland .		Sir John Willoughby, Bart.	
-53½	36½	91	231	Do		J. Brander Dunbar.	
52½	35½	9 1	22	Somaliland		The late W. Babington.	
521	37‡		•••	Do		Major H. G. C. Swayne.	
52 8	40 1	10	36½	Zululand		Major-Gen. Sir Arthur Ellis, K.C.V.O.	
521	42	1112	35	Natwani River .		R. J. Cuninghame.	
-52½	44	11	437	Lebombo Mountains		Count E. Hoyos.	
52 1	41 2	104	39₿	Zululand		Hon. Charles Ellis.	
52 1	37½	9 1	$23\frac{1}{2}$	Somaliland		T. W. H. Clarke.	
52	42	10	33	South Africa		The late W. F. Webb.	
52	36 1	10	39 1	Somaliland		E. N. Buxton.	
-52	4318	10	398	Damaraland		Th. Rehbock.	
52	382	101/2	35	South Africa		James J. Harrison.	
~ 5 2	38 1		•••	Do		C. T. Jones.	
511	36 1	11	29	Zululand		A. J. Brandon.	
511	39½	10	311	Somaliland		W. H. Cobb.	
511	39 2	10 1	28₫	South Africa		W. P. Rylands.	
511	38 1	10	26½	British Central Africa		John Yule.	
511	36 1	11	29	Zululand		A. J. Brandon.	
514	40½	103	28 <u>1</u>	Mashonaland		Col. F. Rhodes, D.S.O.	
50 <u>\$</u>	38	9‡	34	Somaliland		J. Benett-Stanford.	

Length.		Circum-	Tip to			
On outside curve.	Straight line.	ference.	Tip.	Locality.		Owner.
50½	381	10	36 §	Somaliland		Digby Davies.
50½	38	91/2	30 2	Do		Captain G. Campbell.
50½	38	103	22 1	South Africa		R. A. Cooper.
•••	38	12	36	?		Earl of Dartmouth.
50 1	$38\frac{1}{2}$	81	29 1	Somaliland		W. D. James.
50 1	35	9‡	241	Do		B. R. M. Glossop.
-50	39	101	481	Khama's Country .		H. A. Bryden.
50	35	91	131/2	Somaliland		Captain J. M'Call Maxwell.
-50	45	11	37	Abyssinia		Prince de Lucinge.
50	371	10	28 1	South Africa		E. D. Scott.
49 7	39 ⁷	107	40]	Limpopo River .	•	R. M. Sagar-Musgrave.
49 4	401	91	33 2	Zomba, B.C.A		C. C. Bowring.
-49½	37 3	101	191	Somaliland		C. V. A. Peel.
491	36	91	314	Do		Major C. F. Blane.
·49½	40½	113	47	Matabeleland .	٠	Hon. John Ward.
49	38	87	29 8	Somaliland	•	C. Liddell.
-49	39	113	25	Nyasaland		Alex. R. Alston.
48 1	38	10	32	Sabi River	•	Sir Thomas Fowler, Bart.
48	37	10]	27 1	British Central Africa	•	Commander A. T. Hunt.
48	351/2	10	31	Somaliland		R. McD. Hawker.
48	35	$10\frac{1}{2}$	30	Do	•	W. W. Ashley.
-48	•••	10	33½	Do		A. E. Pease.
47 ē	35 3	98	24 §	Do	•	A. H. Straker.
47½	36 1	101	27 §	South Africa	•	British Museum.
47½	35½	91/2	26½	Somaliland	•	Prince Nicolas Ghika.
47½	35	83	38 8	Sudan		Col. Ralph Vivian.
47‡	37 1	101	35½	South Africa		H.R.H. the Duke of Saxe- Coburg and Gotha.
47‡	34 3	9‡	25	Somaliland		Lord Delamere.
47 1 8	344	10	21	Do.		Major C. C. Ellis.

Length.		Circum-	Tip to						
On outside curve.	Straight line.	ference.	Tip.	Loca	lity.		Owner.		
47	35 3	101/2	$33\frac{1}{2}$	Somaliland				J. Kenneth Foster.	
47	35 \$	10	31	Do.				T. W. Greenfield.	
47	$35\frac{1}{2}$	10 <u>1</u>	314	Do.				P. B. Vander-Byl.	
47	35	83	231	Do.				Captain B. L. Carew.	
46 3	37½	94	39 1	Do.	•			F. G. Gunnis.	
46 <u>1</u>	342	10	25‡	Sudan .				LieutCol. Hon. W. Coke.	
46½	34½	9 1		Abyssinia				British Museum.	
461	33\$	91	$24\frac{1}{2}$	Somaliland				Major H. G. C. Swayne.	
46½	34½	10	31	Do.				P. R. Denny.	
461	38	103	411	Do.				A. J. Forbes.	
46	36	92	27 8	Do.				G. H. Cheetham.	
-46	36	91	16	Do.				Count Scheibler.	
-45 2	33₺	91	25 §	Sudan .				Count T. Palffy.	
45½	311	91	181	Somaliland				Lord Delamere.	
-45 8	341	91/2	1110	Do.				Paris Museum.	
45	35	98	32 1	Kaokoland				Captain F. Cookson.	
43½	$34\frac{1}{2}$	9 3	29	Benguela				G. W. Penrice.	
♀ 37	10		•••	South Africa	ι.			F. C. Selous.	
♀ 27½		34	83	Do.				Julius Jeppe.	
9 See	Field, 41	h February	1893	Do.				F. C. Selous.	
- ♀ 26 3		•••	36		?			C. T. Jones.	



Head of Lesser Kudu, from a specimen shot by Mr. Norman B. Smith.

LESSER KUDU (Strepsiceros imberbis).

Godir of the Somalis. Sara of the Abyssinians of Danakil.

Kungu of the Swahilis.

An old male shot by Viscount Edmond de Poncins, 18th Nov. 1897, Digago, Somaliland, measured as follows:—

					inches.
From nose to roo	t of	tail		•	67
Tail					I I 7/8
Height at withers	i .			•	41 है
Round the body					4 I
" neck	•	•	•	•	$20\frac{1}{2}$

Except for its brighter colour, the closer spiral and smaller divergence of the horns, and the absence of a fringe of long hair on the throat (whence the name *imberbis*), this beautiful little antelope might almost pass for a miniature of its larger relation. Height at shoulder about 3 feet 5 inches.

Distribution.—North-East Africa, from Somaliland to German and British East Africa. This antelope generally goes in pairs or threes, and is partial to the cover of thick bush, from which it seldom emerges except for the purpose of feeding. Till the opening up of its habitat, it was very rare in collections.

Leng	gth.	0 :	m *		
On outside curve.	Straight line.	Circum- ference.	Tip to Tip.	Locality.	Owner.
35 1	26 1	71/2	16 <u>1</u>	N. Somaliland .	
34	26 1	•••	•••	Do	tion.) Major H. G. C. Swayne.
34	26	6 3	141	Do	W. W. Ashley.
34	27	6 <u>8</u>	11	Do	R. McD. Hawker, British
33	2 6	71/2	9 1	Do	Museum. Norman B. Smith.
321	25 1	7	8	Somaliland	Capt. T. R. Harkness.
321	254	7	17	Do	Sir Edmund G. Loder, Bart.
-32	26	61	101	Hawash, Abyssinia .	Prince de Lucinge.
-317	25 §		117	N. Somaliland .	J. Menges.
31 2	241	6 <u>‡</u>	141	Do	Lord Delamere.
3112	25 1	6 8	118	Do	Sir John Kirk, K.C.B.
-31½	24	6 <u>1</u>	12	Do	T. W. H. Clarke.
311	24	63		Do	Major C. C. Ellis.
314	25	7	10	Do	Capt. G. F. T. Leather.
314	24½	63	141	Do	Major H. G. C. Swayne.
31	25	7‡	91	Do	W. H. Cobb.
31	24	6 <u>a</u>	91/2	Do	G. H. Cheetham.
-30 3	25	61/2	9₽	Do	Viscount Edmond de Poncins.
30 3	24 8	6 §	117	Do	W. D. James.
30 3	24	6 7	9	Do	E. P. Hare.
30 5	25½	6 8	111	Do	A. H. Straker.
-30½	23	61/2	81	Teita, East Africa .	E. Gedge.
-30½	24	6	121	Somaliland	T. W. H. Clarke.
301/2	24 3	6 §	101	Do	Lord Delamere.
	24			Do	Bombay Natural History Society's Museum.


LESSER KUDU (Strepsiceros imberbis)—continued.

Len	gth.	C '	an .				
On outside curve.	Straight line.	Circum- ference.	Tip to Tip.	Locali	ty.		Owner.
301	24	6₫	117	Somaliland		•	H.R.H. le Duc d'Orléans.
-30	25½	7	14	Do.	•		Capt. M. M'Neill.
29 ³	$22\frac{1}{2}$	$6\frac{3}{4}$	131	Do.			R. Wahrmann.
-29 ³	$21\frac{1}{2}$	$6\frac{1}{2}$	101	Do.	•		Count J. de Bylands.
29½	23	$6\frac{1}{2}$	132	Do.	٠		Prince Demeter Ghika.
298	$22\frac{1}{2}$	$6\frac{3}{4}$	137	Do.	•		E. Lort-Phillips.
-29 1	25	7	I 2 4	Do.	•		A. H. Straker.
29	$23\frac{1}{2}$	6 <u>3</u>	131	Do.			J. Kenneth Foster.
29	23½	$6\frac{1}{2}$	111	East Africa			Sir Robert Harvey, Bart.
-29	22	6 1	12	Somaliland			A. Ohlsson.
29	23 1	6	121	Do.		•	R. McD. Hawker.
28 7	217	$6\frac{1}{2}$	9	East Africa			H. C. V. Hunter.
283	$22\frac{1}{2}$	6 <u>3</u>	11	Somaliland			Sir H. D. Tichborne, Bart.
283	2 I ½	5	9	East Africa		•	Sir Edmund G. Loder, Bart.
28 3	22	6‡	13	Somaliland	•		R. McD. Hawker.
-28 §	23	7	11	Do.			Dr. Percy Rendall.
281	231	7 1	10	Do.			T. W. Greenfield.
-281	$20\frac{1}{2}$	7	131/2	Do.		•	Count E. Hoyos.
28¦	23	5½	115	Do.			Sir John Kirk, K.C.B., British Museum,
•••	23	6 <u>₹</u>	103	Do.		•	Col. Arthur Paget.
28	23		15	Do.			LieutCol. J. W. H. Flanagan.
28	23	6‡	111	Do.			P. B. Vander-Byl.
28	22	6 <u>3</u>	91	Do.			Captain C. H. Villiers.
28	22	6 <u>3</u>	131	Do.			Dr. Donaldson Smith.
27 3	24	6 1	11	Do.			Ford G. Barclay.
	-24	$6\frac{3}{4}$	131	Do.		•	C. V. A. Peel.
271	215	61/2	123	South Somali	iland	•	Sir John Kirk, K.C.B.
271	22	61	7	Somaliland		•	Hon. Walter Rothschild.
-27	•••	6 1	13	Do.	•	•	A. E. Pease.

 \mathbf{x}

LESSER KUDU (Strepsiceros imberbis)—continued.

On outside curve. Straight curve. Circum-ference. Tip. Tip. Locality. Owner. -261 1/8 24 6 1/8 24 6 1/9 10 1/8 24 6 1/9 10 10 10 10 10 10 10 10 10 10 10 10 10	Length.		6 .	Tip to					
26\frac{3}{4} 19\frac{3}{4} 6\frac{1}{4} 10\frac{1}{4} Do. . Edinburgh Museum. 26\frac{1}{2} 22 6 14 Do. . Captain C. S. Timins. 26\frac{1}{4} 21\frac{3}{4} 6\frac{1}{2} 12\frac{3}{8} Do. . Major H. C. Morland. 26\frac{1}{8} 21\frac{1}{2} 6\frac{1}{2} 12\frac{3}{4} Do. . F. G. Gunnis. -25\frac{3}{2} 21\frac{1}{4} 6\frac{1}{2} 12\frac{3}{4} Do. . Count Scheibler. 21 Do. . J. D. Inverarity. -25\frac{3}{8} 20\frac{1}{6} 6\frac{1}{8} 11 Do. . Paris Museum. -25 21\frac{3}{4} 6\frac{1}{2} 10\frac{1}{2} Do. . Digby Davies. 24\frac{3}{4} 19\frac{1}{4} 5\frac{3}{4} 9\frac{3}{4} Do. . Captain W. H. Williamson. 24 20 6 14 East Africa . W. Astor Chanler. -24 19 6 14\frac{1}{2} Somaliland . J. Brander-Dunbar.					Locality.			Owner.	
26½ 22 6 14 Do. Captain C. S. Timins. 26½ 21½ 6½ 12½ Do. Major H. C. Morland. 26½ 21½ 6½ 12½ Do. F. G. Gunnis. -25½ 21½ 6½ 12½ Do. Count Scheibler. 21 Do. J. D. Inverarity. -25½ 20½ 6½ 11 Do. Paris Museum. -25 21½ 6½ 10½ Do. Digby Davies. 24½ 19½ 5½ 9½ Do. Captain W. H. Williamson. 24 20 6 14 East Africa W. Astor Chanler. -24 19 6 14½ Somaliland J. Brander-Dunbar.	-26 }	24	63	13	Somaliland	•	•	Prince Demeter Ghika.	
26½ 21¾ 6½ 12½ Do. . Major H. C. Morland. 26½ 21½ 6½ 12¾ Do. . F. G. Gunnis. -25¾ 21½ 6½ 12¾ Do. . Count Scheibler. 21 Do. . J. D. Inverarity. -25½ 20½ 6½ 11 Do. . Paris Museum. -25 21¾ 6½ 10½ Do. . Digby Davies. 24¾ 19½ 5¾ 9¾ Do. . Captain W. H. Williamson. 24 20 6 14 East Africa . W. Astor Chanler. -24 19 6 14½ Somaliland . J. Brander-Dunbar.	26 3	194	6 <u>1</u>	101	Do.	•		Edinburgh Museum.	
26 1/2 61/2 121/2 Do. F. G. Gunnis. -25 1/2 21 1/2 61/2 121/2 Do. Count Scheibler. 21 Do. J. D. Inverarity. -25 1/2 20 1/2 61/2 11 Do. Paris Museum. -25 21 1/2 61/2 101/2 Do. Digby Davies. 24 19 1/2 51/2 91/2 Do. Captain W. H. Williamson. 24 20 6 14 East Africa W. Astor Chanler. -24 19 6 141/2 Somaliland J. Brander-Dunbar.	26 <u>1</u>	22	6	14	Do.			Captain C. S. Timins.	
-25\frac{3}{4} 21\frac{1}{4} 6\frac{1}{2} 12\frac{3}{4} Do Count Scheibler. 21 Do J. D. Inverarity. -25\frac{3}{8} 20\frac{3}{6} 6\frac{3}{8} 11 Do Paris Museum. -25 21\frac{3}{4} 6\frac{1}{2} 10\frac{1}{2} Do Digby Davies. 24\frac{3}{4} 19\frac{1}{4} 5\frac{3}{4} 9\frac{3}{4} Do Captain W. H. Williamson. 24 20 6 14 East Africa . W. Astor Chanler. -24 19 6 14\frac{1}{2} Somaliland J. Brander-Dunbar.	2 6‡	214	6 <u>1</u>	125	Do.	•		Major H. C. Morland.	
21 Do J. D. Inverarity. -25\(\frac{8}{2}\) 20\(\frac{1}{6}\) 6\(\frac{8}{8}\) 11 Do Paris Museum. -25 21\(\frac{3}{4}\) 6\(\frac{1}{2}\) 10\(\frac{1}{2}\) Do Digby Davies. 24\(\frac{3}{4}\) 19\(\frac{1}{4}\) 5\(\frac{3}{4}\) 9\(\frac{3}{4}\) Do Captain W. H. Williamson. 24 20 6 14 East Africa . W. Astor Chanler. -24 19 6 14\(\frac{1}{2}\) Somaliland . J. Brander-Dunbar.	26 1	$21\frac{1}{2}$	6 <u>1</u>	123	Do.	٠	•	F. G. Gunnis.	
-25\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	-25 8	214	6 <u>1</u>	I 23	Do.	•		Count Scheibler.	
-25 21½ 6½ 10½ Do Digby Davies. 24½ 19½ 5½ 9½ Do Captain W. H. Williamson. 24 20 6 14 East Africa . W. Astor Chanler. -24 19 6 14½ Somaliland . J. Brander-Dunbar.		21	•••	•••	Do.			J. D. Inverarity.	
24 ² 19½ 5 ³ 9½ Do Captain W. H. Williamson. 24 20 6 14 East Africa . W. Astor Chanler24 19 6 14½ Somaliland . J. Brander-Dunbar.	-258	20	6 8	11	Do.	•		Paris Museum.	
24 20 6 14 East Africa . W. Astor Chanler. -24 19 6 14½ Somaliland . J. Brander-Dunbar.	-25	2I4	6 <u>1</u>	101	Do.	•	•	Digby Davies.	
-24 19 6 14½ Somaliland J. Brander-Dunbar,	24 <u>1</u>	19‡	5\$	9 \$	Do.	•		Captain W. H. Williamson.	
•	24	20	6	14	East Africa			W. Astor Chanler.	
-24 203 5½ 9½ Do Julius Jeppe.	-24	19	6	141	Somaliland			J. Brander-Dunbar.	
	-24	202	51/2	91	Do.		•	Julius Jeppe.	



Head of Cow Eland shot by F. C. Selous.

ELAND (Taurotragus oryx).

Doo of the Masaras.

Ee-pofo of the Makalakas.

Eland of the Dutch and English.

Impofo of the Amandebele.

Insefo of the Masubias and

Batongas.

Mofo of the Mashonas.

Mpofu of Swahili.

Mpofu in the Barotse country.

Mpofu in the Lake Ngami country.

Oo-schefo of the Macubas.

Pofo of the Bechuanas.

Tsefu in the Chilala and Chibisa countries.

In addition to being the largest of all antelopes, eland are distinguished from their immediate relatives by the presence of horns in both sexes; these forming a close spiral like a screw, with an upward and outward direction. Female horns are more slender than those of the bulls. Other distinctive features of the genus are to be found in the large dewlap and the long, tufted, ox-like tail.

Bulls of the common eland stand from 5 feet 9 inches to as much as $6\frac{1}{2}$ feet at the shoulder. They have a large tuft of brown hair on the forehead, and the horns are of moderate length and stoutness. The typical race (T. oryx typicus), which formerly extended from the Cape nearly to the Zambesi, has a uniformly tawny skin, without transverse white stripes or a dark brown band above the knees; and appears to be the largest form. Apparently somewhere in Rhodesia a dark brown band is assumed by immature bulls. And as we go northward towards the Zambesi, and thence north and east into the heart of the continent, the bulls have not only this dark leg-band, but the body in both sexes is marked by fine vertical white lines. As this striped variety was discovered by Livingstonianus. Westward the species ranges into Angola.

Throughout Southern Africa, largely owing to the skin-hunters, eland are now becoming exceedingly scarce; and they have already more or less completely disappeared from Cape Colony, Natal, the Orange Free State, Griqualand West, and the Transvaal. In the northern Kalahari, where they subsist for a great part of the year without water, large herds are still to be met with. No species of large game is more easily approached than eland, and, as a rule, none succumbs more speedily to the bullet. Occasionally female eland develop horns in which the spiral is almost obsolete and the length exaggerated; these have been supposed to indicate a distinct species (Antilope triangularis).



Head of Bull Eland.

ELAND (Taurotragus oryx)—continued.

Length straight line.	Circum- serence.	Tip to Tip.	Locality.	Owner.
- 9 35 💏	8,2	20 1 g	Zomba Plain, B.C.A	Dr. Percy Rendall.
· ¥ 33½			?	F. H. Barber.
♀ 32‡	63		Near Chobe River .	M. C. Greaves-Bagshawe.
1 - 9 321	63	54	South Africa	Julius Jeppe.
-321	•••	•••	?	Carl Hagenbeck.
1 9 321	78	83	?	Mr. Justice Hopley.
- 32	•••	•••	?	J. Benett-Stanford.
312	121	223	?	Hon. Walter Rothschild.
¥ 312	7 2	17	Matabeleland	Major R. Hayes-Sadler.
Q 312	71/2	151	?	F. Struben.
Q 313	7½	181	Mashonaland	S. Chillingworth.
318	10}	$25\frac{1}{2}$	East Africa	F. J. Jackson, C.B.
311	123	121/2	Near where Salisbury now stands	F. C. Selous, British Museum.
\$ 31 <u>4</u>	81	20 8	Mashonaland	Do.
₹ 314	7	171	?	Julius Jeppe.
31	13	12	Mashonaland	F. C. Selous.
31	91	21	Nyasaland	F. Vaughan Kirby.
9 31		•••	?	J. Benett-Stanford.
-30 2	1112	282	Nyasaland	Major P. W. Forbes.
302	81	25	?	R. A. Cooper.
♀ 30 ₽	71	213	Lake Ngami	Major H. J. Goold-Adams, C.B., C.M.G.
-30 1	•••	•••	?	O. R. Dunell.
₽ 30	7‡	201	Hanyani River	Capt. M. D. Graham.
-29 3	91	9	British Central Africa.	H. C. Macdonald.
292	91	9	Do	Dr. Percy Rendall.
29 j	8	22	Do	Commander A. T. Hunt.
29½	71/2	8	S. Africa	C. D. Rudd.
-29 1	151	123	Do	A. Ohlsson.
- 9 291	81	17	Do	Mr. Justice Hopley.

¹ The so-called A. triangularis.

ELAND 311

ELAND (Taurotragus oryx)—continued.

Length straight line.	Circum- ference.	Tip to Tip.	Locality.	Owner.
29	10	17	Near Lake Ngami .	Major H. J. Goold-Adams, C.B., C.M.G.
♀ 29	71	181	Matabeleland	Hon. C. Greville.
29	121	14	Do	Abe Bailey.
29	8‡	132	Benguela	G. W. Penrice.
-29			Pungwe	Count E. Hoyos.
28 3	131	41/2	Matabeleland	W. Van Ness.
28₹	1 I ½	24	?	F. Struben.
♀ 28 <u>1</u>	7 1	6 <u>1</u>	Kalahari	H. A. Bryden.
28½	111	88	British Central Africa.	Rowland Ward.
28 <u>1</u>	11	183	Barotseland	R. T. Coryndon.
28 1	10 7	121/2	?	Sir Edmund G. Loder, Bart.
28	10]	19}	British Central Africa.	J. E. Gough.
28	12	19	Upper Shiré, B.C.A	C. C. Bowring.
-28	81	151	E. C. Africa	James J. Harrison.
- 9 28			?	Cape Town Museum.
♀28	8 1	19½	Barotseland	R. T. Coryndon.
27 1	11	16	Mashonaland	Sir John Willoughby, Bart.
$27\frac{1}{2}$	10 1	18	B. E. Africa	Lord Delamere.
$27\frac{1}{2}$	113	121/2	?	R. A. Cooper.
- ♀ 27 ½	$7\frac{1}{2}$	16	?	James J. Harrison.
♀ 27½	81	5	Rhodesia	W. W. Ashley.
$27\frac{1}{2}$	83	10	Matabeleland	Capt. Sir K. Fraser, Bart.
271	11	151	Pungwe	P. B. Vander-Byl.
♀ 27 1	13	20	Matabeleland	Major James Grant.
- ♀ 27	•••	10	East Africa	Sir John Willoughby, Bart.
27	11	103	Pungwe	Marquis of Hamilton.
27	1112	13	Do	F. J. Mitchell.
♀ 26	7 4	13‡	East Africa	Lord Delamere.
26	8‡	101	Kaokaland	Capt. F. Cookson.



Frontlet and Horns of Senegambian Eland, from the Gambia.

SENEGAMBIAN ELAND (Taurotragus derbianus).

Although living specimens appear formerly to have been in the collection at Knowsley, this magnificent species is chiefly known in England by the horns. These are longer and stouter than those of the common species, and thus confirm the statements as to the larger size of the western animal. The bulls are reported to have no dark fore-lock, but a dark-brown mane. Horns of cows are relatively small.

Distribution.—The open districts of the Interior of Senegambia and Gambia.

SENEGAMBIAN ELAND (Taurotragus derbianus)—continued.

Length straight line.	Circum- ference.	Tip to Tip.	Locality	7.	Owner.
-40	131	Single horn	?		Berlin Museum.
-39 8	•••	•••	?		Paris Museum.
341	148	$22\frac{3}{4}$	Gambia .		Sir Edmund G. Loder, Bart.
-34	131	•••	?		F. Coburn.
34	I 23		?		J. Carr Saunders.
-34	1112	17	Gambia .		Dublin Museum.
32½	121	29 ⁷ 8	Do		A late Earl of Derby, British Museum.
32 1	91	123	?		Sir Edmund G. Loder, Bart.
321	13	231	Gambia .		II.R.H. le Duc d'Orléans.
32	13	24 1	?		Hon. Walter Rothschild.
31 8	118	28§	Gambia .		A late Earl of Derby, British Museum.
314	121	151	Do		HC. Goddard.
31 1	12	18§	?		Sir Victor Brooke's Collection.
-31	1112	•••	Gambia .		Sir R. B. Llewelyn, K.C.M.G.
30 8	108	26 ³	?		F. W. Reade, British Museum.
-27 9	138	23∰	Senegambia		Paris Museum.
241	9	6 1	Do		F. W. Reade, British Museum.



Frontlet and Horns of Count Arpad Teleki's Chamois.

CHAMOIS (Rupicapra tragus).

With the chamois we take leave of the antelopes properly so-called. and come to a group connecting the former to a considerable extent with the true goats, one member of the group being indeed commonly designated the Rocky Mountain goat. Most of these ruminants are more or less goat-like in general appearance, having narrow goat-like teeth and short or moderately long tails. Their horns, which are black in colour, are, however, quite distinct from those of the goats. being for the most part cylindrical in section, and curving backwards. From the other members of the group the chamois is at once distinguished by the sharp hook formed by the backward curvature of the horns, which rise almost vertically from the head. The coloration is too well known to need description, although attention may be directed to the dark streak running from the eye to the sides of the muzzle. Height at shoulder reaching 32 inches; weight of male generally about 65 lbs. (although 125 lbs. has been recorded in one case), that of female from 45 to 50 lbs.

Distribution.—The mountains of Central and Southern Europe, from the Pyrenees to the Caucasus. Probably three distinct races may be recognised, namely, the comparatively small izard of the

Pyrenees, the true chamois, or gems, of the Alps, and a third form in the Caucasus.

Length on front curve.	Circum- ference.	Tip to Tip.	Weight.	Locality.	Owner.
1-12	48		1231 lbs. not clean	Retyezat, Carpathians	Count Arpad Teleki. (See illustration.)
-12		91	103 lhs. clean	Tyrol	W. A. Baillie-Grohman.
-12	4	•••		Hinter Riss	H.R.H. the Duke of Saxe-Coburg and Gotha.
-12	•••		•••	Tyrol	Count Arco.
111	3ቔ	6 7	•••	Albreis Morteratch, Engadine	A. E. Pease.
111	31/2	6 <u>1</u>		Engadine ?	Col. Howard.
-11 1		415		Retzezah, Transylvania	Baron A. Nopcsa.
- 11	4				W. A. Baillie-Grohman.
-11	3\$	5½		?	Count John of Meran.
103	31/2	410		Tyrol	Sir Edmund G. Loder, Bart.
103	31/2	410	•••	Do	Do.
-103	34	7 1		?	Eberhard Hollinek.
-10 ¹ / ₂	$3\frac{1}{2}$	58		Austria	H.R.H. the Duke of Braganza.
-10 ¹ / ₂	21	5		3	Count John of Meran.
-10 3	3	48	•••	? .	Marquis Ivrea.
101	31/2	51	•••	?	E. N. Buxton.
101	41	38		Tyrol	Sir Edmund G. Loder, Bart.
-101	3	5#		3	Count Palffy.
-101			•••	Transylvania	C. G. Danford.
- 9 10 1		•••	•••	Grindelwald	F. A. Labouchere.
101	3 1	41/2		Austrian Tyrol .	G. A. Goldschmidt.
10	3∰	4 3		Transylvania	J. G. K. Young.
-10		•••		Grindelwald	F. A. Labouchere.
9 7	31	41	•••	Tyrol	H.R.H. the Duke of Saxe-Coburg and Gotha.
9 3	3 1	37		Transylvania	C. G. Danford, British Museum.
93	3 3	3		W. Caucasus	St. George Littledale.

¹ Greatest width, 71 inches.

CHAMOIS (Rupicapra tragus)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
98	3‡	41/2	Caucasus .			St. George Littledale.
98	31/2	4	Austrian Tyrol			Capt. R. A. J. Montgomerie, C.B.,
98	3	3₿	?			R.N. Hon. Walter Rothschild.
-9§	31/2	31/2	Valtelline Alps			Major C. C. Ellis.
-9½	3	31/2	Alps	•		Dublin Museum.
9-05	3.15	4.75	Savoy .	•		Viscount Edmond de Poncins.
94	3 1	31/2	W. Caucasus			St. George Littledale.
-9 1	28	31/2	Austria .			H. Brinsley Brooke.
9	3	51	Pyrenees .	•		Sir Victor Brooke's Collection.
9	3 1	4	Tyrol .			Julius Jeppe.
₽9	23	41/2	Styria .			Major-Gen. Sir Arthur Ellis, K.C.V.O.
-9	31/2	23	Valtelline Alps			Count Scheibler.
9	3	34	Zillerthall .	•		Noel Fenwick.
83	38	41/8	Austrian Tyrol	•	•	Major-Gen. Sir Arthur Ellis, K.C.V.O.
83	38	3	Caucasus .	•		St. George Littledale.
8 2	3 ½	34	Austria .	•	•	P. B. Vander-Byl.
83	3	44	Caucasus .			Prince Demidoff.
88	31/2	2 3	Do		•	St. George Littledale.
♀8§	•••	318	Bavaria 1870		•	F. C. Selous.
81	2§	4	Alps			The late J. Gould, British Museum.
♀88	21	48	Pyrenees .		•	Sir Victor Brooke's Collection.
81	31	38	?			Sir Clement Hill, Bart.
₽ 8 1	3	2	Caucasus .	•		St. George Littledale.
8	2§		Asturias .			H. Brinsley Brooke.
8	21/2	I 1/2	Caucasus .			Prince Demidoff.
♀ 7½	$2\frac{1}{2}$	7 7	Do. .			Do.



Head of Himalayan Goral. From a specimen shot in Nepal, in the British Museum.

HIMALAYAN GORAL (Cemas goral).

Goral of the W. Himalaya. Pÿ, Pÿar, Rai, Rom of Kashmir. Sáhare, Sarr of the Sutlej valley.

The gorals are near relatives of the serows, from which they may be distinguished by their inferior size, shorter horns, and absence of face-glands, as well as by certain differences in the conformation of the skull. The horns curve regularly backwards, are conical in form, and marked by small irregular ridges for the greater part of their length. The hair is somewhat rough and shaggy, and the tail considerably longer than in the chamois. In general colour the Himalayan goral is brown, tending more or less to rufous or grayish, with the face paler and rufescent, a black line from the nape down the back to the tail (which is also black), and a dark stripe down the front of each leg. Height at shoulder from 26 to 28 inches; weight from 58 to 63 lbs.

By English sportsmen the goral is commonly termed the Himalayan chamois, its habits being very similar to those of the true chamois.

Distribution.—The Himalaya, from Bhutan and Sikim to Kashmir, at elevations from 3000 to 6000 feet.

HIMALAYAN GORAL (Cemas goral)—continued.

Length on front curve.	Girth.	Tip to Tip.	Locality.	Owner.
$-8\frac{1}{2}$		• • •	Bissahir	Major A. E. Ward.
♀-81		•••	Dalhousie	J. Johnston-Stewart.
-8			Kumaon	Major A. E. Ward.
-8	•••	•••	Chamba	Capt. C. B. Vandeleur.
8	3 2	318	Do	Capt. Cecil Levita.
78	34	31/2	?	Hume Collection, British Museum.
$7\frac{1}{2}$	3	21	Near Musuri	A. O. Hume, C.B.
-7½	•••	•••	Mountains near Dal- housie	Capt. F. E. S. Adair.
718	31/2	24	housie	A. H. Ogilvy Spence.
71	3	3	Do	F. W. H. Walshe.
78	31	21/2	Do	Sir Edmund G. Loder, Bart.
-7 1	34	36	Jhelam Valley, Kash- mir	P. H. G. Powell-Cotton.
7	24	34	?	Major J. A. Orr-Ewing.
-7	31	31	?	Capt. H. W. Codrington.
-7	•••	•••	Near Almora, N.W.P.	Capt. B. H. Boucher.
-7	•••	•••	?	Indian Museum.
-7	•••		Punjab	Capt. A. Hicks-Beach.
6 7	41	3	Kumaon	A. S. Crum.
6 7	31/2	27	?	Owen Bevan.
-6 <u>a</u>	•••	•••	?	Major R. H. Rattray.
6₹	3 2	3 1	?	Capt. H. C. Copeman.
-6 2	31/2	2	Dalhousie	J. Johnston-Stewart.
6 §	38	2 3	?	Dr. W. P. Y. Bainbrigge.
6 <u>1</u>	38	31	West of Musuri	B. R. M. Glossop.
6 <u>‡</u>	3	418	?	Capt. L. I. B. Hulke.
6	4	24	?	J. E. Vaughan.
- 9 5½	$2\frac{1}{2}$	13	Jhelam Valley, Kash- mir	P. H. G. Powell-Cotton.
9 4₹	21/4	$2\frac{1}{2}$?	E. L. Phelps.

OTHER GORALS.

	Species.		Collected by	Length on front curve.	Girth.	Tip to Tip.	Loc	ality.		Owner.
Cemas	edwardsi	•	Père A. David	8 լ լ 8լ գ	7½ 7Å	4 2	Moupin	ı, Ti	bet	Paris Museum.
,,	argyrochœti	15	Do.	91	7 1 8	4	Ta - tsie Szec	en - lo huan		Do.
,,	caudatus		Do.	618 61	3 18 4	$3\frac{1}{8}$ $3\frac{1}{2}$	North o	of Pe	kin	Do. British Museum.
,,	cinereus		Do.	$7\frac{1}{2}$	31/8	318	Tibet	•	•	Paris Museum.
,,	griseus	•	Do.	4 3	31	2볗	Do.	•		Do.

JAPANESE SEROW or GORAL (Nemorhædus crispus).

A small species serving in some degree to connect the gorals with the serows, since it agrees approximately in size with the former, but in other characters with the latter.

Distribution.—The southern islands of Japan; an allied species (N. swinhoei) inhabiting the island of Formosa.

Length on front curve.	Circum- ference.	Tip to Tip.		Localit	ty.		Owner.
54	31	31	Japan				British Museum.



Skull and Horns of Himalayan Serow (A. O. Hume's record specimen).

HIMALAYAN SEROW (Nemorhædus bubalinus).

Serow, serowa of Gurhwal and many parts of the Himalaya.

Ramu-Halj, Sálábheer.

Yamu, Kulu.

Aimu, Kumaon.

Goa, Chamba.

A large and clumsily built ruminant, with the ears long, the hair coarse, rather thin, and elongated into a crest from the nape to the withers; the under-fur found in the gorals being absent. The general colour of the upper parts is black or dark gray, with a grizzled appear-

ance, owing to the whitish bases of the hairs; the under parts, shoulders, and thighs being rusty red. Although by no means large, the shining black, somewhat rugose horns form decidedly handsome trophies. Height at shoulder from 33 to 37 or 38 inches; weight from 120 to 190 lbs.

Distribution.—Throughout the Himalaya from Kashmir to the Mishmi Hills, at elevations of from 6000 to 12,000 feet; also recorded from Yunnan.



Head of Male Himalayan Serow. From a specimen shot by W. T. Blanford in Sikim, now in the British Museum.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
-12.25	6.5	2.75	Gurhwal		A. O. Hume, C.B. (See Illus-
-12			Himalaya		tration on page 320.) Major A. E. Ward.
-11			?		Major R. H. Rattray.
10]	5#	31/2	?		Hon. Charles Ellis.
-10½	58	3½	Gurhwal		A. P. Davis.
-10 1			North of Musuri .		Capt. Harry V. Brooke.
-10	54	5½	Mishmi Hills .		A. O. Hume, C.B.
10			Kumaon	•	Major A. E. Ward.

Y

HIMALAYAN SEROW (Nemorhædus bubalinus)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
9 7	5₩	41/2	?		Sir Victor Brooke's Collection.
9₹	5 1 8	6	Pir Panjal		R. Lydekker, British Museum.
91	58	4 1	Kashmir		St. George Littledale.
91	47	31	Nepal	•	The late B. H. Hodgson, British Museum.
91	5∰	3₹	?		Sir Edmund G. Loder, Bart.
- 2 9	5 1	4	Sind Valley .		P. H. G. Powell-Cotton.
9	41/2	$2\frac{1}{2}$	North of Musuri.		B. R. M. Glossop.
-8_{10}^{9}	51 ³ 0	3	Gurhwal		LieutCol. L. L. Fenton.
81	4 3	4	Pir Panjal		Capt. N. C. Taylor.
81	5≹	41/2	Kashmir		Col. R. Pole Carew, C.B.
81	41/2	54	Ranikhet		D. R. Napier.
8	48	43	Near Darjiling .		R. Lydekker, British Museum.
♀ 7월	5	$2\frac{1}{2}$	Gangutri		Col. R. J. Heber-Percy.
71/2	48	3	?		Dr. W. P. Y. Bainbrigge.

BURMESE SEROW (Nemorhædus bubalinus sumatrensis).

Although commonly regarded as a distinct species, there can be little hesitation in classing this animal as a local race of the last, more especially as intermediate forms appear to occur in the neighbourhood of Darjiling. The chief distinction is the more decidedly rufous tint of the hair of the eastern race. Height at shoulder about $34\frac{1}{2}$ inches. Distribution.—From the Eastern Himalaya, Moupin in Tibet, and Yunnan, to Sumatra, occurring throughout the Assam Hills, Burma, Siam, and the elevated tracts of the Malay Peninsula.

Length on front curve.	Circum- ference.	Tip to Tip.	Localit	у.		Owner.
9.25	5	2.75	Garro Hills			A. O. Hume, C.B.
-9. 5	5	5-5	Arakan Hills			Do.
9	5½	5	Burma .	•		The late A. G. Trapmann.
9	5	2	Sumatra .			Hume Collection, British Museum.
-9	5	5	Burma .			VetCapt. G. H. Evans.
-9			Arakan Hills			Indian Museum.
83	51	41/3	?			Hume Collection, British Museum.
8 <u>a</u>	5 ·5	4‡	Muleyit .	•	•	A. O. Hume, C.B.
-8 1	48	2 1	Perak .		•	Perak Museum.
$-8\frac{1}{2}$	5 7	6	Burma .			VetCapt. G. H. Evans.
8 1	5	31/2	Upper Burma			Hon. Walter Rothschild.
-81	51	5 1	Perak .		•	Perak Museum.
7 3	48	41	Assam .		•	A. M. Long.
71/2	4 3	5	?			Duke of Bedford.
7 1	48	3 8	Garro Hills			Dr. Percy Rendall.
57	3 3	3₽	Moulmein, To	enassei	rim	The late Capt. R. C. Beavan, British Museum.



Skull and Horns of Male Takin. From Col. J. Biddulph's record specimen.

TAKIN (Budorcas taxicolor).

Although recently suggested to be related to the musk-ox, this strange ruminant is more generally regarded as an aberrant relative of the serows. It is heavily built, with stout limbs, large lateral hoofs, a short tail, a convex profile, and an almost completely hairy muzzle. But its most remarkable feature is to be found in the horns, which are large, massive, and bent somewhat after the fashion of those of a gnu, curving at first outwards and somewhat downwards, and then bending abruptly upwards about the middle of their length. Height at shoulder about $3\frac{1}{2}$ feet. Horns are present in both sexes, as in the serows and gorals, those of the female measuring about a foot in length, and lacking the curvature of those of the males.

Distribution.—Typically the Mishmi Hills on the northern frontier of Assam, but represented by a variety or allied species in Tibet, and perhaps a third in China. Although living within sight of Indian territory, it does not appear that takin have ever been killed by English sportsmen; and specimens are very rare in collections.

TAKIN 325

TAKIN (Budorcas taxicolor)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.		
-24 1			Mishmi Hills			Col. J. Biddulph. (See illustration.)		
-24.25	12.75	12.75	Do.			Indian Museum.		
228	10 §	143	Assam .		•	British Museum.		
-22.5	13.5	10.75	Mishmi Hills			Indian Museum.		
-22	128	121	Do.		•	Bombay Natural History Society's Museum.		
-21 11	1118	113	Do.	•		Dr. Percy Rendall.		
21	12	13 1	Do.		.•	A. J. Walter.		
20 <mark>7</mark>	118	1 1 7	Assam .			Hume Collection, British Museum.		
20 2	117	121	Mishmi Hills			A. O. Hume, C.B.		
20 3	10	121	Do.	•		Hon. Walter Rothschild.		
20½	107	128	5			The late B. H. Hodgson, British Museum.		
193	11	1112	Mishmi Hills	•	•	Col. J. Biddulph.		
198	11	15	,			Sir Edmund G. Loder, Bart.		
-19	•••	•••	?			Capt. E. T. Dalton, Indian Museum.		
18	10‡	8	?			British Museum.		
151	9 8	5 3	Mishmi Hills		•	The late B. H. Hodgson, British Museum.		
148	83	8 7	Upper Assam	•	•	LieutCol. Graham.		
§ 10 1	8	13	Mishmi Hills	•	•	A. O. Hume, C.B. (See illustration.)		



Frontlet and Horns of Female Takin. From A. O. Hume's specimen.

TIBETAN TAKIN (Budorcas taxicolor tibetanus).

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
-19 ğ	1112	138	Tibet		Père A. David, Paris Museum.
	KA	NSU TAI	KIN (Budorcas	taxic	olor sinensis).
20½	11½	113	Kansu, China		Hon. Walter Rothschild.

BOCKY MOUNTAIN GOAT (Haploceros montanus).

This is one of the very few mammals that are permanently white or whitish at all seasons; and although commonly termed a goat, it really belongs to the same group as the serows, which it closely resembles in the form and colour of the horns. In winter the hair is very long and pure white in colour; along the back it is erect, and much elongated on the withers and haunches, so as to give to the animal the appearance of possessing a pair of humps. The summer coat is comparatively short, and has a yellowish tinge. Height at shoulder just short of 3 feet; weight from 180 to 300 lbs.

Distribution.—North America; throughout the Rocky Mountains, from about lat. 36° in California at least as far north as lat. 60°. By American naturalists the proper generic name of the animal is considered to be Oreannos instead of Haploceros.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
-111		•••	British Columbia		Clive Phillipps-Wolley.
-11	•••	•••	Kutenay, B.C.		John T. Fannin (measured by).
$-10\frac{1}{2}$	5 2	• •	Montana	•	Walter James.
10}	5 ‡	51/2	British Columbia		R. Rankin.
-10 1	6 1	•••	Similkameen River, British Columbia		Arthur Pearse.
10 8	5	6 1	?		E. N. Buxton.
- ♀ 10 <mark>8</mark>	41	•••	British Columbia		Capt. A. Egerton.
10	58	6 8	Do		J. V. Colby.
-9 1	5	•••	Montana		Theodore Roosevelt.
9 3	51/2	61	N.W. Territories		S. Ratcliff.
94	5₺	6	Do		H.R.H. le Duc d'Orleans.
98	5 1	6 1	Do		Sir Edmund G. Loder, Bart.

ROCKY MOUNTAIN GOAT (Haploceros montanus)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
9½	5 1	6 1	Alaska	St. George Littledale.
9½	41/2		North America	J. D. Cobbold.
Չ 9 ½	41	5½	British Columbia .	P. B. Vander-Byl.
6^{7}	51	6 8	East Kutenay, British Columbia	A. E. Butter.
-9 1	5‡	$6\frac{1}{2}$	Bitter Root Mts., U.S.A.	James J. Harrison.
- 오 9흵	41/2	5₿	British Columbia .	A. E. Leatham.
-98	58	6₹	Do	T. W. H. Clarke.
9 1	5 ½	5 1	Do	J. Turner-Turner.
9‡	5 ½		North America	Earl of Lonsdale.
9‡	$5\frac{1}{2}$	53	British Columbia .	G. Lloyd Graeme.
9 1	•••	6	Montana	Thomas Bate, British Museum.
9 1	5 1	5	British Columbia .	Sir Peter Walker, Bart.
9	4 1	6	Do	T. P. Kempson.
$-8\frac{7}{8}$	5½	4 1	Do	Count E. Hoyos.
8‡	48	5 1	Do	Count Scheibler.



Head of Rocky Mountain Goat. From a specimen shot by Sir Edmund G. Loder, Bart.



Head of Male Himalayan Tahr.

HIMALAYAN TAHR (Hemitragus jemlaicus).

The three species of tahr are the first representatives of that great group of ruminants which also includes the true goats and sheep. In all these animals horns are, as a rule, present in both sexes, and are generally more or less distinctly angulated; while the cheek-teeth have tall and narrow crowns like those of the serows, which the tahr serve to connect with the true goats. From the latter tahr are readily distinguished by the small size of their horns, which exceed but little in length the head, as well as by the absence of a beard on the chin of the males. The Himalayan tahr is easily recognised by the great length of the hair of the body, which, although in museum specimens generally combed straight, is very shaggy in nature; and likewise by the form of the horns, which have a knotted sharp keel in front. at shoulder from 36 to 40 inches; weight about 200 lbs. Mr. Wilson, "Mountaineer," had a pair of horns, length 16.5 and girth 10.5, which Mr. A. O. Hume measured.

Distribution.—The Himalaya, from Bhutan to Kashmir.

HIMALAYAN TAHR (Hemitragus jemlaicus)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
-144			Chamba			A. H. Ogilvy Spence.
-14 5			?			II. Vansittart.
14 <mark>16</mark>	83		Chamba			J. S. Rivett-Carnac.
$-14\frac{1}{2}$	9 3	6 <u>3</u>	Do			A. H. Ogilvy Spence.
141	87	48	• 3			A. Cadell.
148	8 7	6 §	Kumaon			A. O. Hume, C.B.
-14 1			?			A. B. Lindsay.
14	•••		?			Capt. R. L. Tottenham.
-14	•••	•••	?			Major A. E. Ward.
137	9	88	Chamba		•	Hugo de Burgh.
137	•••		Do	•		Col. R. J. Heber-Percy.
134	9	111	?			Hume Collection, British Museum.
134	88	6 7	Danpur, Kumaon	•	•	A. O. Hume, C.B.
134	81	7 1	?			I. Morse.
13 <u>\$</u>	8	6 1	?			Hume Collection, British Museum.
-13½	10	•••	?			Major C. F. Blane.
138	9 1	5₹	?			Hon. Walter Rothschild.
134	8 7	5≇	?			Sir Victor Brooke's Collection.
134	81	6 7	Nepal (?)	•	•	Sir Edmund G. Loder, Bart.
131	81/2	•••	?			E. L. Phelps.
-131	9	5	?			P. H. G. Powell-Cotton.
-13	9	•••	?			Indian Museum.
- 13	9 \$	•••	Chamba	•	٠	Capt. A. Hicks-Beach.
-13	91	•••	Himalaya	•	٠	Major-General Alexander A. A. Kinloch,
-123	•••	•••	Chamba	•	•	Col. R. J. Heber-Percy.
-124 121	9 8 <u>1</u>	7 4 10	Do	٠	•	H. L. S. MacLean, Officers' Mess, Queen's Own Corps of Guides. The late J. E. Ubsdell.
121	8 7	43	?			Sir Robert Harvey, Bart.
121	81	61/2	?			H.R.H. the Duke of Saxe-Coburg and Gotha.



Skull and Horns of Male Arabian Tahr. (From Thomas, P.Z.S. 1894.)

ARABIAN TAHR (Hemitragus jayakeri).

A much smaller and rather shorter-haired species than the last, of a generally tawny brown colour, with relatively longer and more slender horns, which are less boldly knotted on the front edge. Height at shoulder about $24\frac{1}{2}$ inches.

This species, of which the type specimen is in the British Museum, was first obtained by Dr. A. S. G. Jayakar, in honour of whom it was named by Mr. O. Thomas in 1894.

Distribution.—Jebel Taw, and probably some of the other ranges of Oman, in South-East Arabia.

Length on front curve.	Locality.		Owner.
-115	Oman Mountains		Surgeon Lieut,-Col. A. S. G. Javaker.



Head of Male Nilgiri Tahr.



Although commonly designated by British sportsmen the "Nilgiri ibex," this species (the warriattu of the native shikaris) is really a member of the same genus as the Himalayan tahr. From that species it is at once distinguished by its short hair and the form of the horns; the latter having the outer surface convex instead of flat, and lacking the knotted front keel. The general colour is dark blackish-brown, old males developing a light saddle-like patch on the back. Height at shoulder from 39 to 42 inches in the males, and up to 35 inches in the females.

Distribution.—The hill-ranges of Southern India, including the Nilgiris, Anamalais, and the Western Ghats nearly to Cape Comorin; usually at elevations of from 4000 to 6000 feet, but occasionally descending to lower levels. This interesting species has been greatly reduced in numbers by constant persecution, but as the herds have now been placed under Government protection, and a special permit is necessary for shooting, there is hope that it may shortly increase.



NILGIRI TAHR (Hemitragus hylocrius)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
-17.5	9.9	6	Nilgiris			Rhodes Morgan.
-17	9 2	•••	Do			Measured by General MacMaster,
162	8 <u>7</u>	5€	?			1869. A. O. Hume, C.B.
-16 1			Nilgiris			St. George Littledale.
16 <u>1</u>	81	6	Do			S. G. Bird.
-16	9.5	5	Do			Rhodes Morgan.
I 5 ½	8	77	Do			Sir Edmund G. Loder, Bart.
158	84	6 3	Do			Martyn Kennard.
15	81	6	Do			St. George Littledale.
15 8	88	4 2	Do			Martyn Kennard.
15 1	85	48	Do			St. George Littledale.
-15	8 <u>‡</u>	51	Do			Sir H. D. Tichborne, Bart.
142	81	5₽	Do		•	Capt. C. S. Timins.
142	8 7	51/2	Do			R. A. Sterndale.
142	8‡	71	Do	•		M. Loam.
141	8 <u>‡</u>	6 1	Do	•		British Museum.
141	85	5 1	Do			Sir Victor Brooke's Collection.
148	8	4 §	Anamalai Hills			T. W. Greenfield.
-141	8 1	4	Travancore .	•		J. D. Inverarity.
-141	8	41/2	Do	•		A. E. Leatham.
14	74	5	Do	•		Capt. Hon. E. Baring.
14	7 1	61/2	Do	•		Major G. S. Rodon.
14	7 f i	6 8	Do			Col. J. Biddulph.
♀ 128	51/2	28	Anamalai Hills	•		M. Loam.
Ç 11 2	5 8	418	Do			Hon. Walter Rothschild.
♀11.25	5.25	4	Do			A. O. Hume, C.B.
9 11 g	54		Travancore .			Col. Hon. Francis C. Bridgeman.
\$ 10	5#	4	Nilgiris		•	St. George Littledale.
♀9 			Do	•	•	Indian Museum.
♀9 <u>₹</u>	5½	31/2	Do			LieutCol. Hon. W. Coke.



Horns of Pir Panjal Markhor. Shot by Major J. C. Shirres, in the Kajnag, April 22, 1884.

ASTOR and the PIR PANJAL MARKHOR

(Capra falconeri typica, and C. falconeri cashmiriensis).

The markhor, of which there are four local races, probably passing more or less completely into one another, introduces us to the true goats (Capra). The males differ from the tahr by their much larger horns, whose length greatly exceeds that of the head, and likewise by the presence of a more or less distinct beard on the chin. In all the varieties of the markhor the beard is extended so as to form a fringe on the throat and chest; the hair on the body being also Moreover, the horns form a spiral, quite unlike the elongated. scimitar-shape characteristic of the ibex and common goat. In the Astor (C. falconeri typica) and Pir Panjal (C. falconeri cashmiriensis) races of the markhor the horns take the form of an open corkscrewlike spiral, with comparatively few turns, recalling those of the kudu: the spiral being most open, the spread widest, and the turns fewest in the Astor variety. Height at shoulder of a Gilgit specimen, 381 inches; of Kashmir specimens, from 38 to 41 inches; weight from

about 200 to 240 lbs. The Astor race is found in Astor and Baltistan, while the Pir Panjal variety inhabits the Pir Panjal and Kajnag ranges of Kashmir, and extends to the north-west into Hazara and Gilgit, where it probably intergrades with the former.



Horns of Astor Markhor. From A. O. Hume's specimen.

Leng	gth.	Circum-				
On outside curve.	Straight line.	ference.	Tip to Tip.	Locality.	Owner.	
-63	•••		•••	?	Major - General Alexander A. A. Kinloch. 1	
-60	•••		•••	Astor	Lucknow Museum.	
-59		12	36	Kajnag	Major J. C. Shirres, D.S.O. (See illustration on p. 333.)	
-59			•••	Pir Panjal	Lucknow Museum.	
-58½	40½	10	38½	Gilgit	Marquis of Lansdowne, K.G., K.C.M.G.	
-56½	38	11	38	Kashmir	J. Benett-Stanford.	
-56	•••	•••		Astor	Capt. H. Trevor.	
-56	36₿	13%	34∯	Do	Viscount Edmond de Pon- cins.	
55	44 1	11	47	?	East India Club.	

¹ Picked up either on the Pir Panjal or Kajnag Ranges by the late Col. Cuppage, and measured by Major-General Alexander A. A. Kinloch.

ASTOR and the PIR PANJAL MARKHOR (Capra falconeri typica and C. falconeri cashmiriensis)—continued.

Leng On outside curve.	-	Circum- ference.	Tip to Tip.	Localit	y.		Owner.
-55	•••			Astor .	•		Otho Shaw.
541		101	26 1	?			A. O. Hume, C.B.
54		Io	33 3	Astor .			Martyn Kennard.
53 1 €	42½	$11\frac{1}{2}$	52 about	Do	•		Sir Victor Brooke's Collec-
53	36 1	10 1	30 1	?			A. O. Hume, C.B. (See
-53		9 1	26	?			illustration on page 337.) Major A. E. Ward.
53	383	9₺	351	Gilgit .			Hon. Walter Rothschild.
52 3	39 \$	12 1	33 2	Astor .			Martyn Kennard.
52	35	91/2	421	Gilgit .	•		Major F. H. Taylor.
51 8	37 1	10	33	Pir Panjal			Sir Edmund G. Loder, Bart.
51 l		118	498	Astor .			Martyn Kennard.
-51 1		12	441	Haramosh		•	P. H. G. Powell-Cotton.
50	38	114	36	Astor .			Sir Edmund G. Loder, Bart.
50	36	10	33	Chitral .			R. H. Macdonald.
50	36 1	12	36	Chilas .			Capt. W. Hayes-Sadler.
50		115	35 1	Astor .			Martyn Kennard.
-50	•••			Do			Capt. H. V. Oliver.
-50	29	111	31	Haramosh			E. L. Phelps.
-49 3	•••	1112	40	Astor .			H. Lennard.
-49 1	32 1	101	43	Gilgit .			Col. J. Biddulph.
-49½	35	10	371	Kajnag .	•		P. H. G. Powell-Cotton.
49 1	36	111	32	Chilas .			L. W. S. Oldham.
-49	39 1	$II_{\overline{g}}^{1}$		Chitral .	•		C. R. Johnson.
-49		11	35	Haramosh	•	•	Capt. H. W. Codrington, Officers' Mess, Q.O.
49	35	14	44	3			Corps of Guides. A. O. Hume, C.B. (See
-49	•••		42	Astor .	•		illustration on page 334.) Otho Shaw.
48½	•••	1112	45	Do		•	Capt. M. Murphy.

1 Shot by Capt. Harry V. Brooke.

ASTOR and the PIR PANJAL MARKHOR (Capra falconeri typica and C. falconeri cashmiriensis)—continued.



Skull and Horns of Pir Panjal Markhor. From A. O. Hume's specimen.

Leng	Length.						
On outside curve.	Straight line.	Circum- ference.	Tip to Tip.	Localit	у.	Owner.	
481	311/2	113	26 <u>1</u>	Baltistan	•	H. Z. Darrah.	
48	374	123	29	Pir Panjal	•	A. O. Hume, C.B. above illustration.)	(See
-48	311	12	38	Haramosh		J. V. Phelps.	

ASTOR and the PIR PANJAL MARKHOR (Capra falconeri typica and C. falconeri cashmiriensis)—continued.

Len	Length.				
On outside curve.	Straight line.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-47½		11	•••	Kashmir.	 Major-Gen. Alexander A. A. Kinloch.
471	37	121	38	Chilas .	 L. W. S. Oldham.
47	36	118	39	Do	 Sir Victor Brooke's Collec



Horns of Variety of Pir Panjal Markhor. From A. O. Hume's specimen.

47	34 1	12	40	Pir Panjal .	•	Dr. W. P. Y. Bainbrigge.
47	• •	11	32	Rondu, Baltistan	•	F. W. H. Walshe.
-47		1 I ½		Indus Valley .		Captain II. M. Biddulph.
46½		$11\frac{1}{2}$	Odd horn	Astor Valley .		British Museum.

Z



Skull and Horns of (1) Cabul, and (2) Suleman Markhor. From A. O. Hume's specimens.

ASTOR and the PIR PANJAL MARKHOR (Capra falconeri typica and C. falconeri cashmiriensis)—continued.

Len	gth.	Circum-	Circum-		0		
On outside	Straight line.	ference.	Tip to Tip.	Locality.	Owner.		
	34	•••	***	?	Mahataja of Travancore, G.C.S.I.		
46	31½	10	35	Bunji	Catalog and the state of		
46	$33\frac{1}{2}$	10	$34\frac{1}{2}$	Kajnag .	Sir Edward Ion Grogan, Bart.		
451	•••	113	$37\frac{1}{2}$	Chilas	Major C. S. Cumberland.		
-45	•.			Kajnag	Otho Shaw.		
45	35 4	11	36	Haramosh	P. Church.		
434		10½	327	Kashmir	Hon. Charles Ellis.		
43½	32½	12	381	Do	I. Morse.		
43	31	91	28	Baltistan	Major C. H. Hayes.		
-43		9₽	30	Do	Capt. T. A. Salt.		
-43	20½	13	35	Astor	E. L. Phelps.		
42 3	•••	12	321	Kashmir	Sir Robert Harvey, Bart.		
42½	32½	111	321	Astor	Hon. Walter Rothschild.		
42	31	I I ½	32	Pir Panjal	W. R. Lawrence.		
-42	36	113		Chitral	C. R. Johnson.		
414		10	30₹	Gilgit	St. George Littledale.		
41		121	428	Indus Valley	Capt. M. Murphy, British Museum.		
41		10	27 8	Pir Panjal	R. Lydekker, British		
39 3	313	11	37	?	Museum. P. W. Cobbold.		
$39\frac{1}{2}$	29	11	$27\frac{1}{2}$	Baltistan	Major H. C. Morland.		
$-37\frac{1}{2}$	•••	12	33	Kashmir	Dublin Museum.		
$37\frac{1}{2}$	$27\frac{1}{2}$	11	35½	Do	Major A. Nugent.		
\$ 16 ‡	13 8	5#	13	Astor	Martyn Kennard.		



Head of Suleman Markhor. From A. J. Grant's Waziristan specimen.

CABUL and SULEMAN MARKHOR

(Capra falconeri megaceros and C. falconeri jerdoni).

In the Cabul race of the markhor (C. falconeri megaceros), from the trans-Indus districts near Cabul, the horns are nearly straight, but show a slightly open spiral, being intermediate between those of the Pir Panjal and Suleman races. On the other hand, in the latter (C. falconeri jerdoni) the relatively small horns form a perfectly straight cone, upon which the spiral ridges run like the "worm" of a screw. The distributional area of the latter race includes the trans-Indus hill-ranges on the frontier of the Punjab, Afghanistan, and Baluchistan,

extending in the Suleman range as far as Mithankot, and also to the Quetta district.

Length in straight line.	Circum- ference.	Tip to Tip.	Locality.	Owner.
48 <u>1</u>	7 4	Odd horn Suleman race	?	British Museum.
39 8	10 1	241	Afghanistan	The late Col. Grant, British Museum.
39 1	121	38	Mountain range 25 miles N.W. of Wana	A. J. Grant. (See illustration, page 340.)
38 1	101	23 7	Afghanistan	H.R.H. the Duke of Saxe-Coburg and Gotha.
37	10	32	?	Capt. G. Roos Keppel.
1 35½	10	25	Baluchistan	Col. J. Biddulph.
-35	•••	•••	Afghanistan	Major A. E. Ward.
343	103	28	Baluchistan	The late Capt. E. W. Codrington.
33	91	194	Afghanistan	Sir Edmund G. Loder, Bart.
321	10	$22\frac{1}{2}$	Bunu	Officers' Mess, Queen's Own Corps of Guides.
32 1	9 3	28	?	A. O. Hume, C.B.
32	10	271	Cabul	A. O. Hume, C.B. (See illustration on page 338.)
317	91/2	27	Suleman Range	Hon. Walter Rothschild.
30 3	83	211	?	Sir Victor Brooke's Collection.
29 2	81	23 3	Baluchistan	Major R. H. Rattray.
29	9	21	Do	The late Capt. E. W. Codrington.
-28	•••		5	Otho Shaw.
27 1	83	21	Sheik Budin	Sir Edmund G. Loder, Bart.
27	9	18	Do	Col. J. Biddulph.
26.75	11.25	22.5	Suleman Range	A. O. Hume, C.B. (See illustration on page 338.)
26 1	•••	18	Bunu	Capt. H. W. Codrington.
26	83	181	Sheikh Budin	Gen. Sir H. Collett, K.C.B.
26	9	16 1	Bunu	Capt. H. W. Codrington.
24	94	20	3	G. Blois Johnson.
-24		•••	Sheikh Budin	Major-General Alexander A. A. Kinloch.
22½	104	153	East Afghanistan .	J. E. Penton.

¹ Straight line 30%.



Head of Himalayan Ibex. From H. Z. Darrah's specimen.

ASIATIC IBEX (Capra sibirica).

The first of the true ibexes, in all of which the long scimitar-shaped horns carry bold transverse knots on the front surface. In the present species the horns are characterised by their large size and well-developed front surface; the second distinctive feature being the long beard of the males. Height at shoulder from 40 to 42 inches; weight about 206 lbs., when cleaned from 128 to 153 lbs.

Distribution.—The mountains of Central Asia, from the Altai to the Himalaya (exclusive of the Pir Panjal), and from the neighbourhood of Herat to Kumaon. Apparently two races of this species, distinguished by the coloration of the lower part of the legs, may be recognised; namely, the Thian Shan ibex (C. sibirica typica), from the Thian Shan, Altai, Baltistan, etc.; and the Himalayan ibex (C. sibirica sacin), from the mountains to the northward of Kashmir and adjacent districts as far east as the source of the Ganges.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-56	•••		Tagdumbash	Major A. E. Ward.
548	10‡	25	Gilgit	Officers' Mess, Queen's Own
53 1	103	20.	Do	Corps of Guides. ¹ Col. J. Biddulph.
52	I I ½	•••	Do	Major A. E. Ward.
-52	10		Baltistan	Maharaja of Travancore, G.C.S.I.
- 52		***	?	The late E. Blyth, P.Z.S. 1840,
$-51\frac{1}{2}$			Kashmir	p. 80. Martyn Kennard.
514	101	16	?	The late E. Blyth, P.Z.S. 1840,
51	91	$28\frac{1}{2}$?	p. 80. Sir Victor Brooke's Collection.
-50	11	301	5	Marquis of Lansdowne, K.G., K.C.M.G.
-50 ·	10.1	21	?	Capt. II. Newton.
-50			?	Capt. II. Trevor.
-50	•••		?	Officers' Mess, 7th Hussars.
49‡	108	251	?	Sir Edmund G. Loder, Bart.
49½	$10\frac{1}{3}$	42	Kashmir	Martyn Kennard.
-49½	111	$27\frac{1}{2}$	Baltistan	Major James Grant.
49	10}	9½	Skardo	Major George Douglas.
-481	9 1	$34\frac{1}{2}$	Gilgit	Officers' Mess, Queen's Own Corps of Guides.
481	9 §	303	Nubra valley, north of Leh	
-48	•••		Skardo	J. L. Wood.
48	107	241	Tilel Valley, Kashmir	J. Campbell of Kilberry.
-47 1			Thian Shan	F. A. Labouchere.
- 47-5	9.5	20.5	Mushral, Nr. Khoga .	A. O. Hume, C.B.
47	113	•••	?	Carl Hagenbeck.
-47	•••	•••	Baltistan	Major E. Guinness.
47	107	294	Kashmir	Martyn Kennard.
47	10		?	Capt. J. Manners Smith, V.C.
-47			Baltistan	A. R. Oldfield.
46½	11	243	Kashmir	Hon. Charles Ellis.
		1 Picked	up by LieutCol. R. E. Hut	chinson.

¹ Picked up by Lieut.-Col. R. E. Hutchinson.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
-46½		•••	?		Indian Museum.
46	93	25	?		Sir Victor Brooke's Collection.
-46	111	27	Sind Valley .		A. O. Hume, C.B.
-46			Kashmir		Capt. Hon. J. G. Beresford.
-45 1	9§	24 1	Baltistan	•	Capt. H. W. Codrington, Officers' Mess, Queen's Own Corps of Guides.
45	103	29	Ladak	•	P. H. G. Powell-Cotton.
45	103	26	Astor		Capt. F. C. Quicke.
-45	10	20 1	Gilgit		Col. J. Biddulph.
-45	•••		Kashmir		Major R. L. Walter.
45	9	121	Baltistan		Capt. C. B. Vandeleur.
-443			Do		Hon. C. B. Fulke-Greville.
445	103	30	Upper Shyok Valley	٠.	A. O. Hume, C.B.
441/2	113	26	?		Major J. A. Orr-Ewing.
441	11	19	?		J. Benett-Stanford.
-44½	•••		Kashmir	•	Major G. A. L. Carew, D.S.O.
441/2	$9\frac{1}{2}$	27 ½	Astor		The late Capt. E. W. Codrington.
441	10	34	Ladak		LieutCol. H. Wade-Dalton.
-44	•••		Baltistan		E. Ezra.
-44		•••	Do		C. H. Rankin.
43 ²	9½	161	Kashmir		S. V. Occleston.
43	10‡	241	?		Sir. Robert Harvey, Bart.
-43½	97	19	?		Hume Collection, British Museum.
-43½	•••	•••	Astor	•	Otho Shaw.
-43½	1 I 🖟	•••	Himalaya		Major-General Alexander A. A. Kinloch.
-43½	10	311	Altai	•	H. J. Elwes.
431	9 1	183	Kashmir		Capt. M. S. Wellby.
43½	91/2	23	Do		Dr. W. P. Y. Bainbrigge.
43½	•••	•••	Baltistan		J. Vaughan.
-43 ¹ / ₂	11		?		Maharaja of Travancore, G.C.S.I.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
$-43\frac{1}{2}$			Haramosh	A. R. Oldfield.
-43 1	103	$22\frac{1}{2}$	Do	J. V. Phelps.
43 1	9₿		Mustagh Range	Capt. F. E. S. Adair.
43	121	35½	Basha Nalla, Baltistan	T. W. Greenfield.
43	11	10	5	Major J. Harden.
43	10	9	5	E. F. Holden.
43	94	28 <u>1</u>	3	R. Rankin.
-43	•••		5	Lucknow Museum.
42 3	10	$25\frac{1}{2}$	Baltistan	P. S. Allan.
-42 3	103	24	Haramosh	J. V. Phelps.
42 §	83	21	Baltistan	H. Z. Darrah.
42 <u>1</u>	10½	22	Kashmir	Col. R. J. Heber-Percy.
42 1	91/2	21½	?	Arnold Pike.
-42½	10 1	27	Astor	E. L. Phelps.
-42½	9 1	•••	Mongolia	J. D. Cobbold.
42	91	26	Ladak	LieutCol. H. Wade-Dalton.
-42	•••	•••	Kashmir	The late Major W. D. B. Fenton.
-42	•••	•••	Baltistan	F. W. Wormold.
-41 4	10	•••	Do	Capt. H. M. Biddulph.
-41 3	9 1	194	Haramosh	E. L. Phelps.
-41 §	10 <mark>1</mark>	$20\frac{1}{2}$	Skardo	Major G. Douglas.
411	9	21	Chitral	Capt. R. L. Tottenham.
41	101/2	26 3	,	Capt. J. H. Purvis.
-4 I	10‡	14	Baltistan	Major H. R. Kelham.
41	9 1	26½	Shyok Valley	Capt. F. E. S. Adair.
41	114	$25\frac{1}{2}$	Baltistan	V. F. A. Keith-Falconer.
41	93	24	Do	Major A. Nugent.
41	9 1	12	Do	W. J. M'Lachlan.
40 1	$10\frac{1}{2}$	22½	Do	Major C. II. Hayes.
40 ⁶	10‡	334	?	Sir Edmund G. Loder, Bart.

Length on front curve	Circum- ference.	Tip to Tip.	Lo	cality.			Owner.
40½	10	201		?			A. C. Hall.
40½	111	2112		?			Hon. Charles Ellis.
40 <u>}</u>	107	19	Altai .				Prince Demidoff.
40	$10\frac{1}{2}$	29‡	Baltistan				Major C. S. Cumberland.
40	10	30	Chilas				L. W. S. Oldham.
- 40	$9\frac{1}{2}$	20		;			Dublin Museum.
-40	9 1	203	Kandgut				Viscount Edmond de Poncins.
40	91	221		?			Viscount Fincastle, V.C.
40	10	97		?			H. De Prée.
40	94	23	Kashgar				R. P. Cobbold.
40	9	29½	Kashmir				B. Vincent.
39 1	9	211	Mongolia				St. George Littledale.
39½	107	29 ³	Kashmir			•	Capt. N. C. Taylor.
$39\frac{1}{2}$	10	213	Ladak				W. R. Codrington.
39½	$9\frac{1}{2}$	221	Do.				Col. Strachey, British Museum.
39½	9½	143	Thian Sha	ın			Col. J. Biddulph.
$39\frac{1}{2}$	93	23	Baltistan				Capt. W. E. Gordon.
$39\frac{1}{2}$	93	13		?			J. R. Carden.
-39½	12	28	Kashmir			•	J. Benett-Stanford.
39	103	$27\frac{1}{2}$?			Capt. L. I. B. Hulke.
39	93	20	Kashmir				Earl of Dunmore.
39	91	17	Do.				A. Leslie Renton.
38½	10	211	Do.				S. Melville.
38 1	107	19	Do.				E. Langworthy.
38 1	go1	22½	Do.				Hon. Walter Rothschild.
38	$6^{\frac{1}{7}}$	16		?			David T. Hanbury.
38	10	27	Altai .		•		Prince Demidoff.
37 ½	107	21		?			Duke of Portland.
35 1	9	11	Saiar Mou	ntain	s, Al	tai	St. George Littledale, British Museum.
$33\frac{1}{2}$	9	21	Altai .				St. George Littledale.

ASIATIC	IBEX	(Capra	sibirica)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-27 7	83	97	Russian Turkestan	Viscount Edmond de Poncins.
- 9 13 1	5	8	Tilel Valley .	J. Campbell of Kilberry.
♀11 5	4	5	Kashmir	Sir Edmund G. Loder, Bart.
911∄	4	81	Do	Capt. Cecil Levita.

ALPINE IBEX (Capra ibex).

This species, which only survives in a protected state, differs from the Asiatic ibex by the much smaller size of the beard of the male, as well as by a slight variation in the horns; the height at the shoulder reaching to about 40 inches. Weight from 85 to 106 lbs., clean. Formerly distributed throughout the higher Alps of Switzerland, Savoy, and the Tyrol, but now surviving only in a few valleys on the Italian side of Monte Rosa. Most of the few specimens now obtainable are comparatively small, and good horns are very scarce in English collections.

Circum- ference.	Tip to Tip.	Locality.			Owner.
10}	1	Valley of Aosta.	•		H.M. the King of Italy.
		Do			Do.
		Do. .			Do.
9	26 3	Do. .		•	Do.
93		Styria			Sir Edmund G. Loder, Bart.
9	26	Valley of Aosta .		•	H.M. the King of Italy.
9	398	Do			Shot by H. M. the late Victor Emanuel, King of Italy.
9 1	183	Alps of Savoy .		•	Sir Edmund G. Loder, Bart.
9	21	?			Dublin Museum.
98	18 '	Valley of Aosta.			C. H. Wilczek.
83	22 lg	Do.		٠	British Museum.
83	171	?			Do.
8 <u>\$</u>	147	Valley of Aosta.		•	Do.
81	13	Do.			Count E. Hoyos.
	9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	ference. Tip. 10½ 1 9 26¾ 9¾ 9 26¾ 9¾ 9 26 9 39¾ 9½ 18¾ 9 21 9¾ 18 8¾ 22½ 8¾ 17½ 8½ 14½	Tip. Valley of Aosta Do. Do.	Tip. Valley of Aosta Do. Do.	Tip. Valley of Aosta

¹ Height at shoulder, 338; weight, 170 lbs.



Head of Arabian Ibex, from the Sinaitic Peninsula.

ARABIAN IBEX (Capra nubiana).

This species may be easily distinguished from both the Asiatic and the Alpine species by the form of the horns, which are very long, rather slender, and with the outer front angle much bevelled off, so that the proper front surface is very narrow, and its transverse knots proportionately short.

Distribution.—The mountains of Southern Arabia, Palestine, the Sinaitic Peninsula, Upper Egypt, and probably also those of Morocco and the interior of Senegambia. Arabian name, beden. Comparatively few European sportsmen have killed this handsome ibex, but it has been shot by E. N. Buxton.

ARABIAN IBEX (Capra nubiana)—continued.



Head of Arabian Ibex, from Arabia; Sclater, P.Z.S. 1897.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
50	9		Southern Arabia .	. Capt. J. T. Brinkley.
42	71		Arabia	. East India Club.
4 I $\frac{1}{20}$			Hadramut, S.E. Arabia	. J. Menges.
392	9 7		Markat	. Do.

ARABIAN IBEX (Capra nubiana)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.	
$39\frac{1}{2}$	8 <u>7</u>	19½	South-East Arabia	٠		Hon. Walter Rothschild.
$35\frac{1}{2}$	$6\frac{1}{2}$	12	Sinaitic Peninsula			W. E. Pease.
35 1	7	15 8	Do			J. D. Cobbold.
-32 1	61/2	7	Do			Captain John Marriott.
321	7	187	Do			E. H. Pease.
318	8	131	Arabia			Dr. Percy Rendall.
318	6 <u>1</u>	131	Sinaitic Peninsula			W. Moncreiffe.
314	6 1	7	Do			W. E. Pease.
29‡	6§	9 1	Do			E. H. Pease.
29		13	Do			Hon. Walter Rothschild.
$28\frac{1}{2}$	71/2	6	Do		•	E. N. Buxton.
25 1	6	12	Do			British Museum.
Ç 10≸	34	21/3	Do			W. E. Pease.
ç 8≇	318	54	Do			W. Moncreiffe.
Th	e follow	ing spe	ecimens are of A	Afric	an	origin :—
46 1	8		North Africa .			British Museum.
43	71	22	Suakin	•		Col. G. E. Lloyd, D.S.O.
421	91	151	Mountain Range of		ut-	The late Major E. M. Barttel

46 1	8	•••	North Al	rica	•	•	British Museum.
43	71	22	Suakin				Col. G. E. Lloyd, D.S.O.
421	91	15‡	Mountair ribai, I				The late Major E. M. Barttelot.
413	7≸	178	North A				Major W. H. Besant.
38 7	71/2	121	Upper E	gypt			Sir Edmund G. Loder, Bart.
38½	71/2	271	Suakin		•		Col. G. E. Lloyd, D.S.O.
38 1	8	21	Do.				Capt. A. King.
35 1	7 8	12	Do.				W. P. Gore-Graham.
30½	8	171	Do.				Commander A. T. Hunt.

ABYSSINIAN IBEX (Capra vali).

This ibex, which inhabits the mountains of the interior of Abyssinia, is still very imperfectly known. Its most distinctive feature appears to be a prominent boss on the forehead.

	25	7:	1	3 Abyssinia .				Hon. Walter Rothschild.
--	----	----	---	---------------	--	--	--	-------------------------



Skull and Horns of Sind Wild Goat. From A. O. Hume's specimen.

PERSIAN WILD GOAT (Capra hircus ægagrus). SIND WILD GOAT (Capra hircus blythi).

The horns of the Persian wild goat, which appears to be the ancestral form of the domestic goat of Europe (C. hircus), differ from those of the various species of ibex by having no distinct front surface, but merely a sharp notched keel, representing the inner front angle of the ibex horn. In old males the beard is very long. The general colour of the upper parts is brownish gray in winter and reddish brown in summer, with the under parts white, and blackish brown and white markings on the face and limbs. Height at shoulder reaching to 37 inches.

The so-called Sind ibex (C. hircus blythi) appears to be a second local race of the species, inhabiting Sind and parts of Baluchistan, where it probably passes imperceptibly into the Persian race. It is distinguished mainly by a slight difference in the form of the horns.

Distribution.—The islands of South-Eastern Europe, and the mountains of South-Eastern Europe and South-Western Asia from the Caucasus through Persia to the confines of Baluchistan and Sind. Native Persian name, pasang (rock-footed).

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
55½	•••	24	Persia		Carl Hagenbeck.
52 8	7 8	83	Sind		A. O. Hume, C.B. (Shot by Col.
48 1	83	137	Caucasus		F. Marston.) See illustration. British Museum.
-48	8	20½	Sind		J. D. Inverarity.
46 <u>8</u>	78	14	Do		Sir Edmund G. Loder, Bart.
46 1	87	18 7	Asia Minor .		F. C. Selous.
46	75	11	Sind		Major C. S. Cumberland.
-45 2	83	9 3	Daghestan		E. N. Buxton.
45 1	8	114	Sind		Col. J. Biddulph.
441	87	213	Caucasus		British Museum.
441	8§	11	?		Hume Collection, British Museum.
438	88	148	Kurrachi District		A. O. Hume, C.B.
43	91		Asia Minor .		F. C. Selous.
43	9	153	Taurus Mountains		Sir Edmund G. Loder, Bart.
43	8 <u>1</u>	22 ⁸	5		J. Carr Saunders.
-42	9	•••	5		The Maclaine of Lochbuie.
-42	7	12	Sind	•	J. D. Inverarity.
-41½			Do	•	Capt. E. C. Tidswell.
-40 8			?		Indian Museum.
39 7	98	11	Erzerum	•	British Museum.
39 7	77	27	Mekran Coast, near Ormara		A. O. Hume, C.B.
38	9 \$?		Major C. S. Cumberland,
-38	10 1	15	Asia Minor .		Capt. John Marriott.
-38	•••	•••	3		Maharaja of Travancore, G.C.S.I.
36	8 <u>‡</u>	6 <u>a</u>	Baluchistan .	•	Hon. Walter Rothschild.
-35 2	8	1112	Do	•	LieutCol. L. L. Fenton.
-35½	9	16	Daghestan		Ford G. Barclay.

WILD GOAT (Capra hircus ægagrus and blythi)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
35₹	9 1	13 8	Mount Ararat	British Museum.
35	81	6 1	Julfa, Persian Frontier	Prince Demidoff.
34½	91/2	18	Baluchistan	Sir Victor Brooke's Collection. (Shot by Major Hogg.)
321	83	113	Julfa	Prince Demidoff.
$3^{2\frac{1}{2}}$	9	131	North Persia	Hon. C. Hardinge, C.B.
-31	7 1	151	Sind	Dr. Percy Rendall.
30 8	81	152	Julfa	Prince Demidoft.
291	82	12	Asia Minor	H. O. Whittall.
29 🖁	9 1	88	Cilician Taurus, Asia Minor	C. G. Danford, British Museum.
27	•••		Antimilo	H. Toppin.
-26 3			Do	Marquis of Ivrea.
₽ 10 8	4 8	3ቔ	Caucasus	Sir Victor Brooke's Collection.

DOMESTIC GOAT (Capra hircus typicus).

521	101	40∰	Daghestan			Sir Edmund G. Loder, Bart.
44 1	6	29 <u>1</u>	Angora .			} Do.
401	91		Daghestan .		٠	J Do.
-37	71/2	35	Meoble, N.B.			Walter Jones.
-30½	71/2	35	Do.			Do.
25 1	7 3	4	South Wales			Major G. Palmer.
24			Joura			Marquis of Ivrea.
-16	6	131	Loch Ness, N.1	В.		C. V. A. Peel.



Head of Spanish Tur. Drawn from Abel Chapman's specimen.

SPANISH TUR or IBEX (Capra pyrenaica).

Although commonly designated an ibex, the Spanish wild goat has horns more like those of one of the Caucasian tur, and is therefore better designated as a species of that group. The horns, which have a sharp inner edge, are twisted in a very open semi-spiral, with the tips generally turned outwards, and are quite unlike those of the true ibex. In having dark and light markings on the limbs the species is, however, much more like the Persian wild goat than either of the Caucasian tur. The beard of the males varies greatly in size according to age and season. Height at shoulder from about 27 to 32 inches at the shoulder; weight, when clean, about 10 stone.

Distribution.—The Pyrenees and the high ranges of Central Spain, Andalusia, and Portugal. The typical form of the species inhabits the Pyrenees; those inhabiting the more southern parts may be distinguished as a separate race (C. pyrenaica hispanica).

SPANISH TUR or IBEX (Capra pyrenaica)—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
31	83		Pyrenees .		Sir Victor Brooke's Collection.
-30 1	91/2	231/2	Central Spain		Abel Chapman and W. J. Buck.
-29 3	81	23 1	Almeira .		H. Brinsley Brooke.
291	91/2	231	Sierra Nevada		Abel Chapman.
28	91	238	Spain .	•	British Museum.
27%	9	25	Do		Sir Edmund G. Loder, Bart.
27½	108	19½	Pyrenees .		British Museum.
25 §	81	16 §	Spain .		Hon. Walter Rothschild.
24½	10	14	Val d'Arras		E. N. Buxton.
238	8	16	Southern Spain		Pablo Larios.
22 3	91/2	18 3	Val d'Arras		E. N. Buxton.
22	7 1	14	Spain .		Pablo Larios.
16	71/2	11	Do		Col. R. J. Heber-Percy.
- 9 10 <u>3</u>	5ቔ	71/2	Val d'Arras		A. E. Leatham.
Ŷ 9⅓	58	6 1	Do. .	•	E. N. Buxton.



Skull and Horns of West Caucasian Tur. Shot by St. George Littledale.

WEST CAUCASIAN TUR or IBEX (Capra caucasica).

Of the two peculiar kinds of wild goats inhabiting the Caucasus, and locally known as tur, the present species is easily recognised by the approximation in the form of its horns (especially in immature individuals) to those of true ibex, and the uniform bright chestnut-brown colour of the hair of the adult male in the summer coat; the chin, beard, and lower parts of the legs being alone black. Although the horns present a considerable resemblance to those of ibex, they are decidedly thicker, have a different curvature, and show distinct knobs, or knots only in their upper half. In young males the knots extend the whole length of the front of the horns, and the long winter coat is a pale drab colour. Height at shoulder about $37\frac{1}{2}$ inches.

Distribution.—The western half of the main chain of the Caucasus. Certain peculiarly-shaped horns appear to indicate the existence of a hybrid race between this species and the East Caucasian tur in the Central Caucasus.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.				Owner.
40 1	125	151	Caucasus	•	•	٠	St. George Littledale.
36‡	115	275	Do_{ullet}				Do.
34 8	1 I 💆	22 7	Do.				Do.
34½	12	22	Kouban, C	aucas	us .		Prince Demidoff.
33 1	105	26 3	Do.				St. George Littledale.
321	112	25½	Do.				Do.
32 1	1112	25 1	Do.				Prince Demidoff.
311	121	24½	Do.				Do.
30 §	118	16	Do.				Sir Edmund G. Loder, Bart.
30	I I ½	191	Do.		•		Hon. Walter Rothschild.
26 2	I I ½	20½	Do.				St. George Littledale.
1 22 ⁸	107	22 <u>1</u>	Do.				Do.
197	10	16 1	Caucasus				P. A. Holt, British Museum.

¹ Presumed hybrid between C. caucasica and C. cylindricornis.



Head of East Caucasian Tur.

EAST CAUCASIAN TUR (Capra cylindricornis).

A very different-looking creature from the last is the wild goat commonly designated by sportsmen the "Caucasian bharal"; this name being derived from the resemblance of its horns to those of the true Himalayan bharal. In spite of this point of resemblance, its affinities are, however, evidently with the goats, although it not improbably indicates a step from the more typical members of that group in the direction of the sheep. In addition to the peculiar form of its horns, this tur is characterised by the extreme shortness of the beard, which merely forms a curling fringe on each side of the chin, instead of the long central tuft observable at certain seasons in the West Caucasian species. The general colour of the fur is uniform dull brown, except on the chin, the tip of the tail, and portions of the legs, where it is blackish brown. Height at shoulder about 3 feet.

Distribution.—The Eastern Caucasus, from Kasbek to Daghestan.

It has been killed by few English sportsmen.

EAST CAUCASIAN TUR (Capra cylindricornis)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.				Owner.
38 1	121		Eastern Cau	ıcasus			Hon. Walter Rothschild.
36	13		Daghestan				J. D. Cobbold.
34½	107	131	Caucasus				Major Talbot.
338	I2	194	Do				British Museum.
321	101	13½	Kasbek .				Prince Demidoff.
32	13	19	Caucasus				Berthold Smith.
311	12	7 1	Do				Ford G. Barclay.
31	11	28	Northern Ca	aucasus	3		Capt. H. H. P. Deasy.
30	12	17	Daghestan	•	. •	. •	Ford G. Barclay.
291	12	20	Caucasus	•			Sir Edmund G. Loder, Bart.
28½	11	16 1	Do.	•			Sir Victor Brooke's Collection.
26 1	10§	18	Do.				Sir Edmund G. Loder, Bart.
26	11	19	Do.	• •			Major C. S. Cumberland.
-26	131	26	Do.				Clive Phillipps-Wolley.
221	10 3	19½	Northern Caucasus				Arnold Pike.
20½	10	1112	Caucasus	•			H.R.H. le Duc d'Orléans.



Head of Male Bharal.

BHARAL (Ovis nahura).

Having horns not unlike those of the East Caucasian tur, the bharal, or blue sheep of Tibet, differs from the goats by the absence of a beard and a strong odour in the males, and on account of these and other points of difference from the goats is placed among the sheep, of which group it forms a very aberrant member. The most distinctive external features are the comparatively smooth and olive-coloured horns, which curve at first outwards and then backwards from the sides of the head; and the bluish gray colour of the thick fur of the back and sides; the flanks, under parts, and legs being handsomely marked with black and white. Height at shoulder about 33 inches; weight about 130 lbs. In the complete absence of glands on the face the bharal differs from the more typical sheep and resembles the goats.

Distribution.—Tibet, from Shigar, in Baltistan, and near Sanju, southeast of Yarkand, to Moupin in Eastern Tibet, and from the main axis of the Himalaya, or locally some distance south of the same, to the Kuenlun and Altyn Tag; in summer usually met with at elevations between 14,000 and 16,000 feet, and apparently never found below about 10,000 feet.

Length on front curve.	Circum- ference.	Tip to Tip.	L	ocality.	Owner.
-32				;	The late B. H. Hodgson, P.Z.S. 1840, p. 66.
317	131	22 l	Ladak		 J. Campbell of Kilberry.

BHARAL (Ovis nahura)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	L	ocality.	•		Owner.
30 7	121	21 7	Gurhwal				A. O. Hume, C.B.
-30½	11	151	Ladak				P. H. G. Powell-Cotton.
30	11	24	Hanle, S	piti			Capt. B. L. Carew.
29 3	121	22½	Gurhwal				A. O. Hume, C.B.
29½	115	25½		?			H.R.H. the Duke of Saxe-
29½	12	26]	Northern	Sikin	n .		Coburg and Gotha. SurgCapt. A. Pearse.
-29	12			?			Major A. E. Ward.
-29			•	?			Lucknow Museum.
-28½				?			Capt. H. Trevor.
28 <u>1</u>	121	26 1	Hanle				F. W. H. Walshe.
-28½	101			?			Indian Museum.
28	11	20 1		?			Hume Collection, British
28	11	16	Hanle	•			Museum. Arnold Pike.
27 8	101	10	Ladak				Capt. G. Campbell.
271	11	21½	Do.	•			St. George Littledale.
-27		•••	Do.		•		Otho Shaw.
27	111	28	Tibet .		•		Major C. S. Cumberland.
26 3	111	23	Ladak		•		Col. F. C. Lister-Kay.
26½	101	20		?			A. S. Crum.
26 1	11 7	23		?			Sir Victor Brooke's Collection.
26 1	103	22		?			Sir Edmund G. Loder, Bart.
26	103	25½		?			R. Johnstone.
26	101	20½		?			Capt. W. H. Williamson.
25 1	121/2	311/2	East of P	adam,	, Zans	kar	Capt. M. S. Wellby.
25 1	114	18 <u>1</u>		?			Capt. C. B. Vandeleur.
- 25½	117		Ladak			•	Major-General Alexander A. A. Kinloch.
258	111	251	Do.				St. George Littledale.
-251	I (½	$24\frac{1}{2}$	Do.				Capt. H. W. Codrington.
25	10	251		?			Major H. C. Morland.
$-24\frac{1}{2}$	124	26		?			J. Carr Saunders.

BHARAL (Ovis nahura)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.		
- 24 1	•••		Upper Indus Valley	Col. J. Biddulph.		
241	11	26 <u>1</u>	?	J. R. Carden.		
24½	11	18	Shyok Valley .	J. V. Phelps.		
-24½			Do. .	E. L. Phelps.		
248	10 7	27 1	South-east of Leh	Major J. A. Orr-Ewing.		
24	11	22 <mark>8</mark>	Nepal	The late B. H. Hodgson, British		
24	112	25	Ladak	Museum. P. Church.		
24	1112		?	Maharaja of Travancore, G.C.S.I.		
-23%	112		Kumaon	W. H. Lane.		
23 1	10	18	Ladak	Capt. F. E. S. Adair.		
23½	9 <u>ł</u>	171	Hanle	Capt. H. W. Codrington.		
23 1	13	25	?	Dr. W. P. Y. Bainbrigge.		
238	112	24 <u>1</u>	?	Sir Robert Harvey, Bart.		
23	12	26	Ladak	H.R.H. the Duke of Saxe-		
23	101	234	Do	Coburg and Gotha. Reginald Beech.		
₹7	41	82	Northern Sikim	Hon. Walter Rothschild.		



Head of Male Arui.

ARUI or UDAD (Ovis lervia).

The only wild sheep found throughout the continent of Africa is the arui, or fechstal of the Arabs, the udad or Barbary sheep of naturalists; a species with horns not very unlike those of the bharal, and also lacking glands on the face, but readily distinguished by its uniformly tawny colour, the fringe of long hair depending from the throat, chest, and the upper portion of the fore-legs, and the unusual length of the tail, which exceeds that of all other wild sheep. In the length of this appendage the arui approaches domesticated sheep, of which, however, it is not likely to be the ancestor. Height at shoulder about 3 feet 3 inches.

Distribution.—The mountains of Northern Africa, from Egypt to Morocco. This sheep is very difficult to find on its own ground.

ARUI or UDAD (Ovis lervia)—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
29 1	123	19	Algeria .	•		V. Cholmondeley.
28½	1112	18	Do. .	ē		Hon. John Ward, British Museum.
28½	1 1 §	181	Algeria .	•		Sir Edmund G. Loder, Bart.
-28	1112	167	S. Tunisia .			J. J. S. Whitaker.
-27 3	111	1418	Algeria .	٠.	•	Viscount Edmond de Poncins.
26 8	103	154	North Africa			British Museum.
26	11	173	Do	•		Do.
25½	117	171	Algeria .			Hon. John Ward.
25½	112	17	Do. .	•		F. de Murietta.
25½	108	16	Do. .			Capt. G. J. Cuthbert.
-25½	121	20½	Do	•		Capt. John Marriott.
-25½	11	18	Tunisia .			A. E. Pease.
25	114	13½	North Africa	•		W. E. Pease.
24½	103	191	Do			Hon. R. A. Ward.
- ♀ 20 1	10	16	Algeria .	•		A. E. Pease.
- 8 19	7	17	Tunisia .			Capt. John Marriott.
- 9 17	71/2	16 1	Atlas Mount	ains		Viscount Edmond de Poncins.



Head of American Bighorn, from W. F. Sheard's specimen.

AMERICAN BIGHORN (Ovis canadensis).

The bighorn of the American continent, inclusive of its local races (frequently regarded as distinct species), is a large sheep, distinguished from the Asiatic argalis, among other features, by the comparative smoothness of the horns, in which the outer front angle is prominent, and the inner one rounded off, and also by the smaller size of the faceglands. There is a well-marked whitish patch on the rump, but the amount of white on the under parts and legs shows considerable local variation. In the typical Rocky Mountain race (O. canadensis typica) the ears are long and pointed, with short hair, and the horns, which are very heavy, diverge but little outwards, and generally have the tips The Californian O. canadensis nelsoni is a paler southern race. On the other hand, in O. canadensis stonei of the North-West Territories the colour of the back is very dark, and the white on the belly and legs sharply defined. And both in this race and the light-coloured O. canadensis dalli of Alaska the horns are lighter, more divergent, and sharper pointed, while the ears tend to become shorter, blunter, and more hairy. Height at shoulder about 3 feet 2 inches. Weight about 350 lbs.

The horns of the ewes are very small in comparison to those of the rams, seldom measuring more than 15 inches on the curve from base to tip. Large male horns are now difficult to obtain, and of late years it is seldom that those of fresh-killed specimens are seen exceeding 38 inches on the curve from tip to tip. American sportsmen are keen to obtain horns of large basal girth; but these, as will be seen from the

following table, rarely exceed 16 inches. The Maclaine of Lochbuie possesses a specimen whose girth according to his own measurement is 19 inches.

Distribution.—North America, from the Rocky Mountains southwards to Sonora, Northern Mexico, and California, and northwards to Alaska and the shores of Bering Sea. The Alaskan race, for at least some portion of the year, is snow-white.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.		
-52½	181		The Selkirks, B.C., 1885	W. F. Sheard. (See illustra-		
-45			?	tion.) W. Grant Mackay.		
$-42\frac{1}{2}$	16 1	25 3	Lower California	George H. Gould.		
42	16	(tips much	Wyoming	Picked up by T. W. H. Clarke.		
	17‡	worn)	Do	T. W. H. Clarke.		
-41½	15		Kootenay, B.C	Measured by John Fannin,		
-40 3	161		Yellowstone	Provincial Museum, B.C. British Museum.		
40 1	151	201	?	Sir Edmund G. Loder, Bart.		
-40	151		Rocky Mountains	Otho Shaw.		
40	15	2112	British Columbia	J. W. R. Young.		
39#	15#	•••	Colorado	St. George Littledale.		
391	161	244	Montana	British Museum.		
391	151	19	?	Sir Edmund G. Loder, Bart.		
-39	152		?	W. A. Baillie-Grohman.		
388	151	22	?	Gerald Buxton.		
38 1	168	•••	Bighorn Mountains .	H. Seton-Karr.		
38 1	15‡	19 1	Montana	Edmund Littledale.		
38 1	16	19	N.W. Territories	S. Ratcliff.		
38	17	•••	Alberta, N.W.T	Arnold Pike.		
38	15	•••	British Columbia	Captain F. Cookson.		
-38	161	•••	Do	Major C. C. Ellis.		
374	152	23 8	Mexico	J. A. H. Drought.		
-37 3	164	221	British Columbia	J. O. Shields.		
371	15½	16	Do	J. Turner-Turner.		
-37	16	31	Wyoming	T. W. H. Clarke.		

AMERICAN BIGHORN (Ovis canadensis)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
37	16 <u>‡</u>	•••	Montana	Major Maitland Kirwan.
37	16 §	16	British Columbia	R. H. Venables Kyrke.
37	151	181	Wyoming	Lord Rodney.
36 3	19	15	British Columbia	C. H. Kennard.
36 3	154	$22\frac{1}{2}$	Wyoming	Moreton Frewen.
36½	141		Do	Gerald Buxton.
36½	16		?	Thomas Bate.
36 1	14		?	J. D. Cobbold.
36 1	148	181	?	Gerald Buxton.
36	144	16 1	Montana	R. H. Sawyer.
36	151/2		Alberta, N.W.T	Arnold Pike.
36	143	16	Wyoming	Captain G. Dalrymple White.
-35 7	144	$17\frac{1}{2}$	Do	Count E. Hoyos.
35 3	151	18 <u>1</u>	British Columbia	G. Wrey.
354	132	171	Do	Hon. S. Tollemache.
35½	16	21	Do	T. P. Kempson.
351	124	16	California	Sir Victor Brooke's Collection.
351	151	181	British Columbia	Sir Peter Walker, Bart.
35	14	181	Do	Admiral Sir Michael Culme-
-35	15	193	Wyoming	Seymour, Bart. Count Scheibler.
35	14	16	Do	Gerald Hardy.
34½	144	19	S.E. Montana	J. A. Jameson.
34½	141	•••	California	G. P. Fitzgerald.
-34	16	17	N.W. Wyoming	A. Rogers.
34	16 <u>‡</u>	20	British Columbia Border	Barclay Bonthron.
331	15½	•••	British Columbia	Admiral Sir Michael Culme- Seymour, Bart.
33	158	18	Do	Capt. E. G. Verschoyle.
33	142	241	Wyoming	LieutCol. Hon. W. Coke.
33	141	22	?	F. H. B. Ellis.
33	14	23	British Columbia	T. P. Kempson.

AMERICAN BIGHORN (Ovis canadensis)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.		
33	151	22	British Columbia.	A. E. Butter.		
323	15½	171	?	C. G. R. Lee.		
$-32\frac{1}{2}$	145	19½	Fraser River, B.C.	A. E. Leatham.		
$32\frac{1}{2}$	15	171	Lower California .	G. Barnardiston.		
32	151	19½	British Columbia .	J. W. Wood, jun.		
32	144	17‡	Yellowstone River .	British Museum.		
311	141	17½	N.W. Territory .	Major Algernon Heber-Percy.		
31	171	•••	Grand Encampment, Wyoming	Frank Cooper.		
- 31	13	22	British Columbia .	T. E. Buckley.		
30 2	15	23 about	?	Hon. Walter Rothschild.		
30 1	152	17½	Lower California .	Ely Quilter.		
30½	15½	18	Wyoming	J. L. Scarlett.		
-30½	14	15½	Do	Hugh Peel.		
30	151	14	Alberta, N.W.T	F. C. Williamson.		

ALASKAN BIGHORN (Ovis canadensis dalli).

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.			
34	125	18 1	Alaska			Rowland Ward.			
33	123	. 15	Do.			Hon. Walter Rothschild.			
32½	131.	201	Do.			J. T. Studley, British Museum.			
₽ 9 1	47	8	Do.			Do.			



Skull and Horns of Male Kamschatkan Bighorn.

KAMSCHATKAN BIGHORN (Ovis canadensis nivicola).

Although the Kamschatkan wild sheep can be readily distinguished from the Rocky Mountain bighorn (O. canadensis typica) by the slenderness of the horns at the points, and their wider tip-to-tip measurement, as well as by shorter, blunter, and more thickly haired ears, the shorter face, the longer hair of the body, and the inferior size of the white patch on the rump, yet there are several varieties of the bighorn from the more northern districts of America which tend more or less completely to bridge over these points of difference. Among these are the Alaskan bighorn (O. canadensis dalli) and the N.W. bighorn (O. canadensis stonei), in which the horns are of the Kamschatkan type, and the ears are shorter than in the Rocky Mountain race. These transitions indicate that all the bighorns are essentially local modifications of the same animal; the Kamschatkan form being, as might be expected from its isolation, the most aberrant. Like the Alaskan race, the Kamschatkan bighorn appears, at least sometimes, to turn white in winter. at shoulder about 37 or 38 inches; weight about 330 lbs.

Distribution.—Kamschatka, and the Stanovoi Mountains as far south as the sources of the Utschuri River; perhaps also in parts of Siberia.

KAMSCHATKAN BIGHORN (Ovis canadensis nivicola)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.				Owner.		
-38	131	26	Kamschatka		ė	•	Dr. F. H. H. Guillemard's party in the "Marchesa," P.Z.S. 1885, p. 675.		
$-35\frac{1}{2}$	14	26½	Do.				Do.		
341	11 7	172	Do.				British Museum.		
34	13½	238	Do.				Hon. Walter Rothschild.		
324	131	23	Do.				Sir Edmund G. Loder, Bart.		
- 311	14	26	Do.				Dublin Museum.		
3112	138	21	Cape Chepuns	ke			Col. J. Biddulph.		
$-31\frac{1}{2}$	134	24	Kamschatka				Dublin Museum.		
$27\frac{1}{2}$	138	$25\frac{1}{2}$	Do.				Rowland Ward.		
\$ 10 ³	5	141	Do.				Hon. Walter Rothschild.		



Skulls and Horns of Siberian Argali, from specimens shot in the Altai by Major C. S. Cumberland.



Head of Marco Polo's Sheep. From David T. Hanbury's specimen.

MARCO POLO'S SHEEP (Ovis poli).

In common with the following members of the genus Ovis, this splendid sheep has the transverse wrinklings of the horns well developed, and small glands are present on the face. The horns of the adult male are characterised by their comparative slenderness and great length, forming a spiral of more than one complete circle, with the front angles typically well developed. The hind-quarters show a large amount of white, extending over the greater part of the thighs; and in winter the throat is furnished with a voluminous ruff of long white hairs, which disappears in summer. Height at shoulder about 4 feet, or perhaps rather less; weight of adult male about 22 stone.

Distribution.—Typically the plateau of the Pamirs in Central Asia, but represented by a closely allied race (O. poli karelin) in the Thian Shan range of Turkestan, in which the horns are generally somewhat shorter, and may have the front outer angle rounded off. This sheep was first definitely made known in England by

specimens obtained during the Second Yarkand Mission under the late Sir D. Forsyth; since which date it has been killed by Mr. St. George Littledale, Major C. S. Cumberland, Capt. H. Bower, Viscount de Poncins, and many other sportsmen.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
-75	16	54½	Pamir	Field-Marshal Lord Roberts, V.C.
73	15	48	Little Pamir	The late Col. H. C. B. Tanner.
-71 ·	151	534	Great Pamir	Viscount Edmond de Poncins.
70	17	52	Do	Maharaja of Kuch Behar.
69 <u>1</u>	154	56	?	Marquis of Lansdowne, K.G.,
69 <u>1</u>	141	39	Tagdumbash	K.C.M.G. Sir Edmund G. Loder, Bart.
68 <u>1</u>	15	35₹	?	Lewis Flower.
68	17	43	Pamir	Col. R. Pole Carew, C.B.
-68	16	52	?	Indian Museum.
-67 8	16	53≇	?	Col. J. Biddulph, Indian Museum.
67	16 1	42	Pamir	Col. R. Pole Carew, C.B.
-67	16		Thian Shan	Maharaja of Travancore, G.C.S.I.
67	151	42½	?	Duke of Westminster.
66 7	158	46	?	Sir Edmund G. Loder, Bart.
66 <u>7</u>	137	46 1	Valley between Little and Great Pamir	A. O. Hume, C.B.
66	163	47	Do.	Do.
66	151	44	Do.	Hume Collection, British Museum.
66	154	42	?	A. Leslie Renton.
65 <u>1</u>	16	53	Great Pamir (16,000 feet) .	Col. T. E. Gordon, British Museum.
-65	16	41	Central Asia	Dublin Museum.
65	16 1	49½	?	Major C. F. Blane.
-65			?	St. George Littledale (presented to the Empress of Russia).
64 <u>1</u>	16 <u>1</u>	46	Pamir	Major C. C. Ellis.
64 1	161	41	Do	W. Lawrence.
642	151	39	Do	H. C. V. Hunter.
64	15½	50	Little Pamir	R. P. Cobbold.
64	15	49	?	Duke of Portland.

MARCO POLO'S SHEEP (Ovis poli)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
63§	16 <u>1</u>	42½	Pamir		Hon. Walter Rothschild.
63½	154	57½	Valley between Gree	at and	A. O. Hume, C.B.
638	158	46 1	Little Pamir Tagdumbash .		Earl of Dunmore.
63	16	49½	Little Pamir .		R. P. Cobbold (Bachelors' Club).
623	16 ‡	51	Tagdumbash .		T. W. Greenfield.
62½	15	57	?		Sir Edmund G. Loder, Bart.
62	151	40	?		Hon. Charles Ellis.
61 1	151	46]	Tagdumbash .		E. L. Phelps.
60₹	154	46 7	Pamir		St. George Littledale.
60	16	49½	Little Pamir .		R. P. Cobbold.
60	15\$	46 3	Tagdumbash .		T. W. Greenfield.
60	151	46	Do		P. Church.
-60	•••		?		Capt. H. Bower.
- 60	17	•••	Tagdumbash .		H. Dauvergne.
-60	16	52	Do		H. Lennard.
59 3	151	45 8	Pamir		Earl of Dunmore.
59	16]	47	5		Viscount Powerscourt.
59	15\$	41	5		Martyn Kennard.
59	14	423	Tagdumbash .		David T. Hanbury.
58	16	43	Do		Do.
57 ₹	15	463	?		Major-General Sir Arthur Ellis, K.C.V.O.
574	143	50	Pamir		Reginald Beech.
57½	16 1	49 1	?		W. Lawrence.
$57\frac{1}{2}$	144	48	Pamir		Reginald Beech.
57	154	42	Do		Hon. R. A. Ward.
56 8	15 1	$35\frac{1}{2}$	Do		St. George Littledale.
56	171	42	Do		E. L. Phelps.
56	15	44	Do		E. P. Tennant.
554	153	43	?		J. Carr Saunders.

MARCO POLO'S SHEEP (Ovis poli)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.
55€	16 1	43	Pamir			Major C. S. Cumberland.
55½	16 7	44	Do			Sir Edmund G. Loder, Bart.
55½	15 1	39	?			The late W. F. Webb.
55	16	44	?			H.R.H. le Duc d'Orléans.
54 7	15‡	488	Pamir		•	Sir Victor Brooke's Collection.
54 1	16	34\$	West Mongolia .			St. George Littledale.
54	163	39	?			Viscount Fincastle, V.C.
53 8	15§	433	Lake Karakol .			St. George Littledale.
53	141	45	Tagdumbash .			Major H. C. Morland.
53		•••	?			Lucknow Museum.
-52	16	41	Tagdumbash .			H.M. the King of Italy.
52	151	39	Do			Capt. J. Manners-Smith, V.C., C.I.E.
511	15	48	Do			J. G. Millais.
51	15‡	383	Do			A. Leslie Renton.
50½	15	42 ³	Altai Plateau, Pami	r.		Capt. J. Manners-Smith, V.C.,
- ♀ 14·7	5.90	15.75	Great Pamir .			C.I.E., British Museum. Viscount Edmond de Poncins.
\$ 10⅓	5½	131/2	Tagdumbash .	•		David T. Hanbury.



Skull and Horns of Siberian Argali. From a specimen shot by St. George Littledale in the Altai.

SIBERIAN ARGALI or AMMON SHEEP (Ovis ammon).

Closely allied to Marco Polo's sheep, but distinguished by the thicker and less expanded horns, which often have the outer front angle much rounded off, and the wrinklings very fine. A more or less distinct white patch on the rump, not extending on to the thigh, which is dark-coloured like the back; no ruff on the throat, even in the long winter coat. In summer the coat of old males, which is very short, tends to become more or less light-coloured all over. Height at shoulder from about 45 inches to 4 feet; weight from about 250 to 350 lbs.

As in the Tibetan race, considerable individual variation may be noticed in the horns, some having the outer front angle much more developed than usual; it does not appear that these differences can be accounted for by age.

Distribution.—In former times apparently extending from the Baikal Mountains in the south of Eastern Siberia through Northern Mongolia to the Semipalatinsk Altai; now chiefly restricted to the two latter localities.

SIBERIAN ARGALI or AMMON SHEEP (Ovis ammon)—continued.

Length on	Circum-	Tip to Tip.		Locality			Owner.		
front curve.	ference.		A 14 *	-			St. Coome Littledele		
62 <u>‡</u>	193	381	Altai .	•	•	•	St. George Littledale.		
62	19	382	Do.	•	•	•	H. J. Elwes.		
611	19‡	39 1	Do.	•	•	٠	St. George Littledale, British Museum.		
59 1	198	38 1	Do.		•	•	St. George Littledale.		
56 <u>1</u>	18 <u>1</u>	$33\frac{1}{2}$	Do.	•			Major C. S. Cumberland.		
55	18	39	· Do.				Prince Demidoff.		
54 2	183	30	Do.			•	Major C. S. Cumberland.		
54 1	192	334	Do.	•		•	Do.		
541	19	35	Do.				Do.		
53 1	19	22	Do.			•	St. George Littledale.		
52½	194	33	Do.			•	Prince Demidoff.		
52½	18 <u>1</u>	38½	Do.	•			Do.		
52	19	37½	Do.				Do.		
511	18 1	26	Do.				Do.		
512	198	31	Do.				Do.		
51	192	33	Do.				Do.		
51	19‡	33	Do.				St. George Littledale.		
501	18 1	374	Do.				Do.		
50	19½	31 1/2	Do.				Major C. S. Cumberland.		
50	19‡	36	Do.	•			St. George Littledale.		
50	19‡	271	Do.				Şir Edmund G. Loder, Bart.		
49½	20 1	25	Do.	ě			St. George Littledale.		
47 3	19	311	Siberia	٠.			British Museum.		
451	20	271	Altai .				Prince Demidoff.		
40	16	33	Do.				Princess Demidoff.		



Head of Tibetan Argali.

TIBETAN ARGALI (Ovis ammon hodgsoni).

Chiefly distinguished from the true ammon sheep by the development of a distinct white ruff on the throat of the males, at least in the winter coat, and also by the less degree of lateral expansion of the horns, which do not form more than a single complete circle, and are generally broken at the tips. The wrinkles on the horns are perhaps somewhat less prominent, and the outer front angle is frequently well developed.

The height at the shoulder is perhaps rather less than in the true ammon, of which this sheep is best regarded as a local race. A specimen measured by Major Greenway was 76 inches from the nose to the tip of the tail, and the weight about 212 lbs. In another male, whose age was estimated at 10 years, the height at the shoulder was 43 inches, the girth 50 inches, and the weight 205 lbs. (P. H. G. Powell-Cotton).

Distribution.—The plateau of Tibet, from Northern Ladak to the districts north of Sikim, and northwards to the Kuenlun; eastern limits unknown.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.				Owner.			
57	183	29	Ladak				Arnold Pike.			
50½	18 1	19	Pangong	Lake			Sir Edmund G. Loder, Bart.			
-50	17	•••	Tibet .				W. H. Lane.			
$-48\frac{1}{2}$	19	•••	Do				E. Howard Brooke.			
48	16	23	Rudok				Major G. A. L. Carew, D.S.O.			
48	181	20	Ladak				Major H. C. Morland.			
48	18	31	Do.	•			Major Hon. A. Dalzell.			
-47	17		Do.			•	Otho Shaw.			
46½	193	20	Do.				A. O. Hume, C.B.			
-46½	161	21	Ladak				P. H. G. Powell-Cotton.			
46½	162		Do.		•		Capt. H. M. Biddulph.			
-46	19			;			Major C. S. Cumberland.			
46	16	17		?			Hon. R. A. Ward.			
45½	16 1	17		?			Duke of Teck, G.C.B.			
45	17	16	Ladak			•	J. V. Phelps.			
-45				?			Lucknow Museum.			
448	171	22½	Tibet.				The late B. H. Hodgson.			
-44½	17	2178	Ladak	•			A. E. Leatham.			
44	16 <u>1</u>	19‡	Do.			•	Major C. S. Cumberland.			
44	16	171	Do.	•		•	W. R. Lawrence.			
-44	18	•••	Do.				Maharaja of Travancore, G.C.S.I.			
-43	•••	•••	Do.			•	Capt. W. Bailey.			
42½	161	19		;			A. O. Hume, C.B.			
42½	17			;			LieutCol. T. Greenaway.			
$-42\frac{1}{2}$	16	15	Ladak			•	Capt. H. M. Biddulph.			
42½	16	18	Do.	•			Hume Collection, Brit. Museum.			
42 1	151	•••		?			J. Carr Saunders.			
42 8	16 1	20	Tibet .	•	•		H.R.H. the Duke of Saxe-Coburg and Gotha.			

TIBETAN ARGALI (Ovis ammon hodgsoni)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.		Locality.			Owner.
42 1	16 3	14‡	Ladak	•			Hume Collection, British Museum.
42	15 1	17	Near H	lanle			Capt. F. E. S. Adair.
42			Rudok				Major G. A. L. Carew, D.S.O.
41 7	16		Tibet.				Hon. Walter Rothschild.
411	16	15		;			Duke of Westminster.
-41½	•••	•••	Ladak				G. G. Thatcher.
41 lg	163	19	Do.				Sir Victor Brooke's Collection.
-41	•••	181	Do.				Major A. E. Ward.
41	174	•••	Do.	•			E. L. Phelps.
41	16 1	17	Do.	•			R. Johnstone.
40½	141	26 1		?			St. George Littledale, Brit. Museum.
40 8	175	20 3	Ladak	•			Hon. Charles Ellis.
40 1	16]	19	Do.	•			Col. R. J. Heber-Percy.
40 1	152	183	Do.	•			A. C. Bailey.
-40	17		Do.		•	•	Major-General Alexander A. A. Kinloch.
-40				?			Capt. H. Trevor.
40	17	19½	Ladak	•	•	•	Sir Edmund G. Loder, Bart.
-40	17	•••		;			Indian Museum.
-40	···	•••	Ladak	•	•	•	Otho Shaw.
-40 1	174	•••	Hanle	•	•	•	Capt. H. W. Codrington.
39½	17	14	Ladak	•	٠	•	F. W. H. Walshe.
39 1	17	144	Do.	•	•	•	Hon. Charles Ellis.
39 1	171	21½	Do.	•	•	•	Dr. W. P. Y. Bainbrigge.
39	17	19	Do.	•	•	•	Sir Robert Harvey, Bart.
ბ 18 3	81	23	Do.	•	•	•	David T. Hanbury.
\$ 18	7	194	Tibet.		•	٠	The late B. H. Hodgson, British Museum.
♀ 14½	71/2	171	Ladak				Sir Victor Brooke's Collection.

MONGOLIAN ARGALI (Ovis ammon jubata).

Nearly allied to the Tibetan race, having a distinct yellowish-white throat ruff and generally similar horns, but, in some cases at least, the outer front angles of the latter much rounded off. The white on the buttocks and hinder surface more abundant and purer in colour.

Distribution.—Eastern Mongolia to the north of Pekin.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.					
44 8	16 1	23	Manchuria .		•	Henri de Bardi.	Bourbon,	Comte de		



Head of Male Shapu.

SHAPU or URIAL (Ovis vignei).

A much smaller sheep than either of the Asiatic argalis, with comparatively slender and well-wrinkled horns of considerable length, which when fully developed curve forwards along the sides of the face; the males with a more or less strongly developed white ruff on the throat. General colour varying from rufous brown to gray in summer, with the chest, under parts, and portions of the legs white, and sometimes blackish "points." Females with small horns. Height at shoulder about 32 inches; weight about 120 lbs.

Distribution.—From Ladak and Zanskar to Russian Turkestan, Afghanistan, Baluchistan, Southern Persia, the North-West Frontier of India, and the Punjab Salt-Range. Two local races, which probably intergrade in the Indus valley, may be recognised. First, the typical urin of Astor, the sha or shapu of Ladak (O. vignei typica); and secondly, the smaller Punjab urial (O. vignei), in which the colour is redder, the ruff more developed, and the front angles of the horns often show a knotted keel. There is also O. vignei blanfordi of Baluchistan; and it is possible the Persian form, in which the ruff is said to be but little developed, may form a fourth.

(a) SHAPU (Ovis vignei typica).

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
39	112	151	?	Major Sir H. S. Rawlinson, Bart.
38₫	121	114	?	J. Carr Saunders.
38½	111	81	Near Leh	E. L. Phelps.
36 1	$11\frac{1}{2}$	41	Near Ley	A. O. Hume, C.B.
-36 1	113	•••	Near Leh	Major A. E. Ward.
-36 1	107	104	Baltistan	Capt. T. A. Salt.
348	11	•••	?	East India Club.
-334	121	13	Ladak	P. H. G. Powell-Cotton.
33 1	12	12	Do	Arnold Pike.
-33 1	1112	10 §	Do	Sir Edmund G. Loder, Bart.
33	101	15	Do	F. W. H. Walshe.
32 1	101	114	,	Sir Victor Brooke's Collection.
-32	11	9 3	Ladak	Col. J. Biddulph.
314	104	7	Do	Col. F. C. Lister-Kay.
-31	11	•••	Do	J. D. Cobbold.
301	81	19 ·	Do	K. Dingwall.
30	114	16	?	Hon. Walter Rothschild.
-30	•••	•••	Ladak	Otho Shaw.
-30	11	•••	?	Maharaja of Travancore, G.C.S.I.
-30	•••	•••	?	Lucknow Museum.
29 8	108	•••	. 5	H. C. V. Hunter.
29½	101	13	Ladak	J. V. Phelps.
29½	101	151	Tochi Valley	J. Johnston-Stewart.
29	114	•••	?	Sir Robert Harvey, Bart.
29	11	9 1	Ladak, W. bank of the Indus	A. Leslie Renton.
-28	•••	•••	Gilgit	Otho Shaw.
28	103	13	?	B. Vincent.
271	118	18 1	?	Martyn Kennard.
-27½	10	•••	West Ladak	Capt. F. E. S. Adair.

(a) SHAPU (Ovis vignei typica)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.		Loc	ality.			Owner.
$-27\frac{1}{2}$			Skardo .			•		Major George Douglas.
-27½	114	19	Astor .					Capt. H. W. Codrington, Officers' Mess Queen's Own Corps of Guides.
-27 1	101	15	Do.					Capt. F. E. S. Adair.
26 3	103		Shigar .			•		E. L. Phelps.
$26\frac{1}{2}$	$10\frac{1}{2}$	13	Ladak .			•	•	St. George Littledale.
26½	9 1	$19\frac{1}{2}$?			W. J. M'Lachlan.
$25\frac{1}{2}$	111	155			?			Hon. Charles Ellis.

(b) URIAL (Ovis vignei cycloceros).

Strictly speaking, cycloceros has no right to stand for this race, as it was given to the typical form, but at present it seems inadvisable to change it.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
39½	103	181	Punjab	Major F. H. Taylor. (See illustration.)
$-38\frac{1}{2}$		•••	Jouaka Land	Royal Artillery Mess at Attock.
-37	9	•••	?	Major J. C. Shirres, D.S.O.
38 3	9.75	9-5	Chita Oapar Range, near Attock	Royal Artillery Mess, Woolwich, Major C. F. Massey.
36 ₫	9	9	Hills north-west of Peshawur	Mess of 60th Rifles, Lord Walter Fitzgerald.
35½	101	16	Gulran, Afghanistan .	Dr. J. Aitchison, British Museum.
-351	104	20 3	?	Mess, 21st Punjab Infantry.
35 1	$10\frac{1}{2}$	9 1	Punjab	Major F. H. Taylor.
35	11	14	Near Cabul	Major J. W. M. Cotton.
$34\frac{1}{2}$	81	16	Sind	L. Napier.
331/2	91	$12\frac{1}{2}$	Salt-Range	A. O. Hume, C.B.
$33\frac{1}{2}$	9§	83	?	Capt. R. L. Tottenham.
321	10	I I ½	?	Sir Victor Brooke's Collection.
32 3	7 1	12	Punjab	G. Blois Johnson.
321	10	1112	3	Hume Collection, British Museum.
311	91/2	16 3	Salt-Range	Capt. H. W. Codrington.
-31	104		Punjab	Major-General Alexander A. A. Kinloch.

(b) URIAL (Ovis vignei cycloceros)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.		
31	98	142	Punjab			Major R. H. Rattray.		
30 ⁷ 8	98	2018	Salt-Range			Sir Victor Brooke's Collection.		
30∄	114	11‡	Do.	•		Col. J. Biddulph.		
-30	81	61/2	:	?		Dublin Museum.		



Head of Male Urial. Major F. H. Taylor's specimen.

29g	108	·	Salt-Range		H. C. V. Hunter.
293	101	163	Sind Hills .		Major C. S. Cumberland.
29½	9		Sheik Budin, nea Ismail Khan	r Dera	Capt. Harry V. Brooke.
28‡	9	151	Kusan, Afghanis	itan .	Dr. J. Aitchison, British Museum.
$-27\frac{1}{2}$		•••	Afghan Hills		Col. J. Biddulph.
$26\frac{1}{2}$	91	12	Salt-Range		Capt. B. L. Carew.
-26			Sind		J. D. Inverarity.
-26	••		Salt-Range		Major-General Alexander A. A. Kinloch.
-25 7	8 <u>1</u>	141/2	Punjab .		Dr. Percy Rendall.
25 8	93	133	Salt-Range		Sir Edmund G. Loder, Bart.

(b) URIAL (Ovis vignei cycloceros)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
25 1	98	171	Kohrod, Persia .	Sir Victor Brooke's Collection.
-24	81	13	Salt-Range	Col. J. Biddulph.
21½	9₹	15	?	LieutCol. H. Wade-Dalton.

(c) KELAT RACE (Ovis vignei blanfordi).

Possibly inseparable from the last, in which case the name blanfordi should supersede cycloceros, which was really given to specimens of typica.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality	7.	Owner.		
37½	101	11	Haji Khan, 3000 ft.	Kelat,	A. O. Hume, C.B.		
36	9 1	17	Kelat		Do.		
318	98	134	Do		Hume Collection, Brit. Museum.		
28 <u>1</u>	9	10	Baluchistan		Col. J. Biddulph.		
$-25\frac{1}{2}$	9 1	102	Do		LieutCol. L. L. Fenton.		

ARMENIAN MUFLON (Ovis orientalis).

This sheep differs from all the preceding species by the absence of horns in the females; the horns of the males being not unlike those of the urial, but curving backwards, so that their points are situated behind the neck instead of beneath the eyes. General colour of upper parts some shade of yellow or foxy red; the under parts and lower portions of the legs white. Height at shoulder about 2 feet 9 inches.

Distribution.—The mountains of Armenia, Eastern Persia, and Asia Minor.

Length on outside curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
40 1	101	51/2	5	W. Burchart Barker, British Museum.
36 1	108	53	?	British Museum.
36	101	15	Persia	Hon. Walter Rothschild.
30½	108	18	?	Sir Victor Brooke's Collection.
-26 1 0	81	12}	Cilician Taurus	C. G. Danford.
1 24 ¹ / ₂	9‡	I I ½	Persian Frontier	Prince Demidoff.
24	9ቔ	17	Asia Minor	Col. J. Biddulph.
1 23 ³	91/2	131/2	Persian Frontier	Prince Demidoff.
234	91/2	113	Persia	Major C. S. Cumberland.

¹ Determination provisional.



Head of Male Cyprian Muslon. (From Biddulph, Proc. Zool. Soc. 1884.)

CYPRIAN MUFLON (Ovis orientalis ophion).

A small local race of the preceding distinguished by certain differences in coloration, and the complete rounding-off of the front outer angle of the horns of the male. This is the smallest of the wild sheep, and comparatively a rare trophy. Height at shoulder about 28 inches. The horns closely resemble those of the Armenian race in general characters, but are less massive, and curve gradually from the base, instead of diverging nearly straight outwards, as is generally the case in the latter. The type specimen of the species is preserved in the Berlin Museum.

Distribution.—The Tröodos Mountains of Cyprus.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
25	8	152	Tröodos Mountains		H. Williamson.
-24	8	(weight 5 stone) 4	Do.		The late Lord Lilford.
$23\frac{1}{2}$	8-15	I 2·20	Do.		Col. J. Biddulph.
23	7	54	Do.	• .	British Museum.
22 ⁷ / ₈	7 1	6	Do.		Gen. Sir R. Biddulph, British
221	8	124	Do.		Museum. Sir Victor Brooke's Collection.
$-17\frac{1}{2}$	8	11/2	Do.		Dublin Museum.
16	71	I 6½ (weight 4 stone)	Do.		Hon. Walter Rothschild.



Head of Male European Muflon. From a specimen in the British Museum, shot by Ford G. Barclay.

EUROPEAN MUFLON (Ovis musimon).

The large light-coloured saddle on the otherwise dark summer coat of the rams of this handsome and well-known little sheep is so distinctive of the species that nothing in the way of description need be attempted in this place. It may be observed, however, that the ewes are generally hornless, and that the horns of the rams curve forwards so as to have their tips below the eyes, and are comparatively massive, with the wrinkles of a type somewhat different from that obtaining in the Armenian sheep. Height at shoulder about 27 inches.

A good account of muslon-shooting will be found in Mr. E. N. Buxton's Short Stalks (first series).

Distribution.—At the present day the islands of Sardinia and Corsica.

Length on front curve.					7.		Owner.
341/2	83	163	Sardinia				W. Moncreiffe.
30½	8 7	11	Do.			•	C. Sloane Stanley.
30 1	85	14	Do.				Rhys Williams.

EUROPEAN MUFLON (Ovis musimon)—continued.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.			Owner.		
29 3	85	8 3	Sardinia			Rhys Williams.		
29 3	8	12 1	Do.			C. Sloane Stanley.		
29 3	8		Do.			J. D. Cobbold.		
29§	83	11	Do.			Hon. R. A. Ward.		
29	83	11	Do.			W. Moncreiffe.		
28 1	81	21	Do.			E. N. Buxton.		
28 3	9	10	Do.	,		Ford G. Barclay.		
28 1	9₿	98	Do.			Edinburgh Museum.		
27 g	84	5 1	Do.			W. E. Pease.		
27	$9\frac{1}{2}$	101	Do.			Sir Edmund G. Loder, Bart.		
27	81/2	10	Do.		•	British Museum.		
27	83	1112	Do.			Rhys Williams.		
-26 §	81	9	Do.			H. Brinsley Brooke.		
-26	10 8	101	Corsica			Capt. John Marriott.		
$25\frac{1}{2}$	8≩	6	Sardinia		٠	A. Y. Lethbridge.		
25	8 1	10	Do.			Hon. Walter Rothschild.		
$24\frac{1}{2}$	91	9 7	South-We	st Sau	rdinia	Sir Victor Brooke's Collection.		

DOMESTIC SHEEP (Ovis aries).

The history and ancestry of the various breeds of domestic sheep are lost in the mists of antiquity, and naturalists are totally unable to point to the wild stock from which any or all of them are derived. This is the more to be regretted, seeing that the Swedish breed is the type of the genus *Ovis*. Most domesticated breeds differ from wild sheep by the woolly nature of their coat; but since hairy tame sheep are met with in several uncivilised countries, this point of difference is of comparatively little importance. More weight has been attached to the great length of the tail, which is much longer than even that of the arui; and, as mentioned above, that species is almost certainly not the father of the domesticated sheep. There is, however, some degree of probability that the long tails of the domestic breeds are due to a kind of degeneration. And if this be really the case, their ancestry

might be looked for among the musions or urial or some allied extinct form, since the horns of most breeds approximate to the musion type. The presence of horns in the females of many breeds, Dorsetshire, for example, may be either an inherited character, or a redundancy of a nature similar to that which has produced four, or even five, horns in the males of certain oriental breeds. Some eastern sheep, like the Wallachian, have further departed from the musion type by the development of upright corkscrew horns comparable with those of the markhor or kudu.

Length on outside curve.	Circum- ference.	Tip to Tip.	Locality.				Owner.		
39 1	8 <u>a</u>	21	?				H. E. Surtees.		
37	81	20	Loch Awe, N.B.			•	H. Murray.		
351	8	16 7		?			Sir Victor Brooke's Collection.		
33	11	221	Yarkand				Hume Collection, Brit. Museum.		
32	85	27 8	Do.				Do.		
28 3	8	211	Scotland				Rowland Ward.		
2 2 §	91	20 <u>1</u>	Yarkand				Hume Collection, Brit. Museum.		
22 8	9 1	221	Do.				Do.		
18	83	16 1	Fezzan				British Museum.		

Abnormally-horned Specimens.

			•	
Length on outside curve.	Circum- ference.	Tip to Tip.	Number of Horns.	Owner.
20 1 + 16	6+42	151	Four	Sir H. B. Meux, Bart.
194+14	$6\frac{1}{2} + 4\frac{1}{2}$	21 + 16	Do.	P. C. Millbank.
178+148	78+4 1	$4\frac{1}{2}+6\frac{8}{4}$	Do.	Hume Collection, British Museum.
17+112	$6\frac{3}{4} + 5$	$6 + 8\frac{1}{2}$	Do.	British Museum.
$13\frac{1}{4} + 10\frac{7}{8}$	$5\frac{3}{4} + 4\frac{1}{2}$	•••	Five	Hume Collection, British Museum.
13+98	7§+5	142	Four	A. O. Hume, C.B.
12+92	6 1 + 4	171	Do.	Do.

The following specimens belong to the Wallachian breed:—

Length on the curve.	Length in a straight line	Girth.	Tip to Tip.	Owner.
33	171	$7\frac{1}{2}$	19½	H.R.H. le Duc d'Orléans.
32	24	7 1	38	Do.

MUSK - OX



Head of Bull Musk-Ox.

MUSK - OX (Ovibos moschatus).

In spite of its name, this Arctic ruminant has no near affinity with the members of the ox tribe, the molar teeth being more like those of the sheep and goats, the muzzle, except for a small strip between the nostrils, hairy, and the tail reduced to a mere stump concealed among the long hair of the hind-quarters. On the other hand, the resemblance to the sheep is not very close, the horns, which in old males nearly meet in the middle line of the forehead, being of a totally different form and structure, and the skull likewise very distinct. In the males the horns are much flattened and expanded at the bases, after which they are bent suddenly down behind the eyes, to curve upwards again at the tips. In the females they are much smaller, less expanded, and not approximated at their bases. In both sexes their texture is coarse and fibrous, and their colour yellow. The long coat of dark brown hair depending from the back and sides like a mantle affords an adequate protection against the rigors of an Arctic winter; and the broad spreading hoofs, with hair on their under surface, give a firm

foothold on snow and ice. Height at shoulder about 4 feet; weight about 8 to 9 cwt.

Distribution.—Arctic America, approximately north and east of a line drawn from the mouth of the Mackenzie River to Fort Churchill on Hudson Bay, Greenland, and Grinnell-land, in lat. 32° 27'; approximate southern limit lat. 40° N.

Length on outside curve.	Breadth of Paint	Tip to Tip.	Locality.		Owner.
-29%	13		?		W. W. Hart.
27 }	10	27 1	Barren Grounds		David T. Hanbury.
27 }	121	27	Northern Cana Do.	da	Earl of Lonsdale.
27	91/2	251	Do.		David T. Hanbury.
267	11	27	Do.		Warburton Pike.
26 3	I 2 8		North America	-	J. Rae, British Museum.
26]	131	27 §	Do.		British Museum.
244	11	$25\frac{1}{2}$	Barren Grounds		Warburton Pike.
241	71/2	19	Northern Cana Do.	aa	J. Talbot Clifton.
241	101	26	Do.		Hon. Walter Rothschild.
24	20 (circum-	30	North America		Earl of Lonsdale.
24	ference) 9 2	231	Do.		Sir Edmund G. Loder, Bart.
223	91	19‡	Grinnell-land		Col. W. H. Fielden, British
211	9	27	?		Museum. Dublin Museum.
♀ 18g	41	•••	North America		A. G. Dallas, British Museum.
5 18 1	41	•••	Do.		Do.



Horns of Cape Buffalo (F. H. Barber's specimen).

CAPE BUFFALO (Bos caffer).

Inyati of the Swazis and Zulus.

Mbogo and Nyati of the Swahilis.

Nari of the Basutos.

Nadi in the Barotse and Lake N'gami countries.Mboa and Nyati in the Chilala and Chibisa countries.

Among the distinctive features of this splendid bovine may be noted the enormous helmet-like mass formed by the closely approximated bases of the horns in old bulls, the backward inclination and comparatively slight angulation of the horns themselves, the shortness of the face, and the great width and size of the heavily fringed and flapping ears. In colour both the skin and the sparse hairs with which it is clothed are for the most part jetty black; the hairs themselves being directed uniformly backwards from the nape to the rump. Height at shoulder about five feet.

Distribution.—Southern Africa from the Cape to the southern bank of the Congo on the west side, and approximately to the neighbourhood of the Victoria Nyanza on the east side of the continent. Northwards of this it not improbably gradually passes into the Abyssinian buffalo. Except on the Zambesi, Chobe, and some neighbouring rivers, buffaloes have now become very scarce in South Africa. But between Umtali and the east coast at Beira, and also from the latter station to the mouth of the Zambesi, they are to be met with in vast herds; and a few years ago existed in countless numbers. Here they are much protected by the unhealthy nature of the country, which is deadly to Europeans, except between the end of May and November. Save for a few protected

herds in the Addo bush, the Knysna and Zitzikamma forests, and thickets of the Fish and Sunday rivers, the species has long since been exterminated in the Cape. It is even rare and local in British East Africa, where, as in other districts on the eastern side of the continent, the rinderpest has of late years played sad havoc with the herds.

Greatest Outside.	Width. Inside.	Tip to Tip.	Width of Palm measured on face of horn.	Locality.	Owner.
-49½	43 8	30 7	12	Sabi River	F. H. Barber. (See illustra-
49	441	40]	111	Limpopo	tion.) Sir Richard Glyn, Bart.
-48 7	44½	36	10	Chiromo, B.C.A	H. C. Macdonald.
47	41	$28\frac{1}{2}$	12	Limpopo	The late W. F. Webb.
47	40 7		121	East Africa	F. J. Jackson, C.B.
•••	40½	26	•••	Do	Prince Boris Czetwertynski.
46 <u>1</u>	441	371	61/2	Pungwe	C. M. Swire.
45 3	411	37 1	•••	South Africa	British Museum.
45 3	40	283	12	Nyasaland	F. Vaughan Kirby.
451	41	30	•••	East Africa	W. Astor Chanler.
45½	•••	•••	111	Kilimanjaro	H. C. V. Hunter.
-45	394	36½	14	East Africa	Count Scheibler.
-45	•••	•••		Nyasaland	S. Pulley.
44 2	39 8	271	114	East Africa	Sir Robert Harvey, Bart.
441	391	29	12	Pungwe	F. S. Staples.
44½	39 1	29	14	Near River Ramok- webani, S. Africa	F. C. Selous.
44½	•••	$37\frac{1}{2}$	12	East Africa	F. Charrington.
44	40 1	374	12	Do	Capt. R. A. J. Montgomerie, C.B., R.N.
-44	40	•••	111	South Africa	J. Lamont.
-44	37½	14	•••	Chiringoma, P.E.A.	F. Vaughan Kirby.
43€	37 3	26½	123	East Africa	Hon. Walter Rothschild.
431	$38\frac{1}{2}$	25½	9	Pungwe	H. R. Holden.
43 1	38 1	$24\frac{1}{2}$	13‡	South Africa	Sir John Willoughby, Bart.
43	37ቔ	293	14	Do	C. D. Rudd.
42 3	38	338	9\$	East Africa	E. J. L. Berkeley.

CAPE BUFFALO (Bos caffer)—continued.

Greatest Outside.	Width. Inside.	Tip to Tip.	Width of Palm measured on face of horn.	Locality.		Owner.
-42½		•••	13	Manicaland .		F. Lean.
-42 1 2	36½	29	141	E. C. Africa .		James J. Harrison.
428	36 3	29½	12½	?		Sir Victor Brooke's Collec-
42	38	35 1	9	East Africa .		tion. Lord Delamere.
42	37 8	26 <u>‡</u>	11	Pungwe		Capt. F. H. Lehmann.
42	37	30 1	10}	Barotseland .		E. D. Scott.
-42	36	19	16	?		Mr. Justice Hopley.
41 ½	37 3	34½	10	Pungwe		R. H. Venables-Kyrke.
$41\frac{1}{2}$	36 7	28	•••	South Africa .	•	British Museum.
$-41\frac{1}{2}$	39	39	144	South-east Africa	•	James J. Harrison.
41	38	35	87	East Africa .		Lord Delamere.
41	36 3	37	11	Zambesia .		Comdr. A. T. Hunt.
41	36	30	111	?		Rowland Ward.
41	35 8	28 3	•••	Mashonaland .		F. C. Selous, British Museum.
41	36 <u>1</u>	26½	$9\frac{1}{2}$	Pungwe		Capt. Lord Douglas Compton.
-40 §	35	19§	131	Do		Count E. Hoyos.
40½	36 3	$25\frac{1}{2}$	$9\frac{1}{2}$	Do		Marquis of Hamilton.
-40½	36	35	15	South Africa .	•	A. Ohlsson.
-40 1	36	20 3	12	Lower Shiré .		Staff-Surgeon J. Dowson, R.N.
	36	21		East Africa .		Sir John Willoughby, Bart.
40	37	35		Do		Gen. Sir Lloyd William Mathews.
40	35₹	•••	145	Do		F. J. Jackson.
9 4 0	34 8	21	8 1	Pungwe		J. W. Allen.
40	35	214	10	Matabeleland .		Major James Grant.
40	34 ³	214	8	Pungwe		C. C. Gouldsmith.
394	$34\frac{1}{2}$	91/2	14	Do		G. L. Bonham.
39 1	35	$34\frac{1}{2}$	9	Do		R. Hughes.
39.12	37	36.11	12	British East Africa	•	Prince Henry of Liechtenstein.
39	36	35 1	12	Zululand .	•	A. J. Brandon.

CAPE BUFFALO (Bos caffer)—continued.

Greatest Outside.	Width. Inside.	Tip to Tip.	Width of Palm measured on face of horn.	Locality.	Owner.
39	$34\frac{1}{2}$	$22\frac{1}{2}$	10 (Mashonaland	A. Neilson.
39	34 1	27	117	Benguela	G. W. Penrice.
39	34	19	10]	Shiré Valley	H. H. Williams.
$38\frac{1}{2}$	334	27	91	Pungwe	R. K. Micklethwait.
38 1	$32\frac{1}{2}$	193	13	3	Sir Edmund G. Loder, Bart.
381	$33\frac{1}{2}$	231	91/2	Shiré Valley	C. C. Bowring.
38	33 1	26½	8	Pungwe	A. Cameron.
38	$33\frac{1}{2}$	311	144	?	Julius Jeppe.
38	321	$20\frac{3}{4}$	11	5	Do.
38	341	314	10 7	?	F. Watkins.
37½	33 3	30	10	British Central Africa	J. E. Gough.
$-37\frac{1}{2}$	313	25	113	Shiré Valley	Alex. R. Alston.
37 1	$33\frac{1}{2}$	32 3	121	East Africa	Sir John Kirk, K.C.B.
37 1 8	317	238	13	?	H.R.H. the Duke of Saxe- Coburg and Gotha.
37	32	16 1	9½ (about)	Pungwe	Viscount Ennismore.
37	33	30 1	103	Goronza Plain, P.E.A.	Earl of Dunmore.
37	311/2	21	I I ½	?	L. Rawstorne.
37	30½	19	10	South Africa	R. A. Cooper.
♀ 36½	33	26 <u>1</u>	5½	Mozambique	F. Vaughan Kirby.




Skull and Horns of Abyssinian Buffalo.

ABYSSINIAN BUFFALO (Bos caffer æquinoctialis).

A smaller animal than the last (height at shoulder about 4 feet), with the general colour blackish or tawny brown, tinged locally with rufous, and tending to grayish on the legs. Horns smaller, much more flattened at the bases, where they are more widely separated, and in some, although not all, cases retreating less markedly behind the plane of the eyes. The fact that in East African specimens of the Cape buffalo the horns are usually less rugged than in more southern examples, tends to confirm the view that the present animal should be regarded as a variety rather than as a species.

Distribution.—From Abyssinia and southern Somaliland through the Egyptian Sudan for a considerable distance up the White Nile; but the southern limits, if definable, not yet determined.

Greatest Outside.	Width. Inside.	Tip to Tip.	Width of Palm.	Locality.	Owner.
40 -37 8 36	37 34 1 29	32 34 1 24	114 84 	Abyssinia Upper Basaland	E. Lort-Phillips. J. Menges. E. Gedge.
32 314 314 315 31 -304 302 - 2 238	28 28 1 26 1 26 2 28 1 26 2 23 20 1	22 25	6# 8½ 9½ 10 10 8½ 5½	Basé Country	The late F. L. James. Col. Ralph Vivian. G. P. V. Aylmer and W. D. James. British Museum. Sir Edmund G. Loder, Bart. Count T. Palffy. LieutCol. Hon. W. Coke. J. Menges.



Horns of old Male Senegambian Buffalo, from the type specimen in the British Museum.

SMALLER AFRICAN BUFFALOES.

Lake Tchad Buffalo (Bos caffer brachyceros). Senegambian Buffalo (Bos caffer planiceros). Dwarf Congo Buffalo (Bos caffer nanus).

Despite the enormous differences, alike in size, colour, and the form of the horns, between the little red buffalo of the Congo and its gigantic black cousin of the Cape, such a gradual and apparently complete transition can be traced from the one type to the other, that both seem local modifications of one very variable animal. In the dwarf buffalo of the Congo the colour of the hair is red or yellow, the ears are fringed with very long hair, and the horns are much flattened at the base, with long smooth tips directed upwards. This type may be traced, with some modifications, to the Gaboon and Gambia; but in the interior of Senegambia it apparently passes into the larger brown buffalo, with more laterally expanded and recurved horns, known as B. caffer planiceros. And this latter is, in its turn, very close to the Abyssinian buffalo (B. caffer æquinoctialis) of East Central Africa, which is nearly allied to the typical Cape form. Another type is indicated by the Lake Tchad buffalo (B. caffer brachyceros), first discovered by Denham and Clapperton. In height the Congo dwarf buffalo only reaches some 42 inches at the shoulder.

SMALLER AFRICAN BUFFALOES—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.
23½	15	10	Gold Coast	Major G. S. C. Jenkinson.
23	16 1	81	Nigeria	Julius Jeppe.
-21	154	4	Do	Capt. C. A. Wilding.
21 g	123	21	West Africa	British Museum.
21	15	9 1	Nigeria	A. Ohlsson.
184	104	5½	Central Africa	Capt. Denham and Col. Clapperton, Brit. Museum.
18	16 1	17½	Victoria Island, Lower Niger	R. H. Monck-Mason.
17	181	191	Nigeria	A. W. M. Brodie.
9 17	113	6 3	Central Africa	British Museum.
17	12	141	Nigeria .	P. A. Clive.
16 <u>4</u>	10}	6 1	Jebba, Nigeria	Capt. A. H. Festing.
152	11	9 1	Nigeria	Sir Edmund G. Loder, Bart.
15	6 <u>1</u> 1	14	Do	Major A. J. Arnold, D.S.O.
-14 2	111	88	Gaboon	Dr. Percy Rendall.
14 §	8 1	7 1 8	Sette Cama., W. Africa	British Museum.
141	I 2 1	108	?	Sir Victor Brooke's Collection.
13 7	10]	28 <u>1</u>	Gambia	Late Earl of Derby, Brit. Museum.
\$ 13	10 <u>1</u>	101	Nigeria	Julius Jeppe.
♀11 <u>1</u>	91		Lokoja	Major A. J. Arnold, D.S.O., Brit. Museum.
119	8 <u>1</u>	10	West Africa	Sir Edmund G. Loder, Bart.
♀9 1	74	81	Do	Hon. Walter Rothschild.

1 Width of palm.



Skull and Horns of Congo Buffalo, from Major A. J. Arnold's specimen.



Head of Indian Buffalo. Shot by the Maharaja of Kuch Behar.

INDIAN BUFFALO or ARNA (Bos bubalis).

No one is the least likely to confuse this animal with the Cape Both belong, indeed, to the same group of the genus Bos. and have the rounded upper portion of the head and angulated horns. In the Indian species, however, the head is much longer, the ears are narrower and less heavily haired, and the horns of the male are widely separated on the forehead, and totally different in form. Two types of horns may be recognised, one very massive, and curving regularly up from each side of the head in a subcircular manner; the other much slenderer, though quite as long often, directed for the greater part of their length almost straight out from the head, and always with a wider spread; these latter horns being those of females. Height at shoulder about 6 feet 2 inches; girth behind shoulder, 10 feet 8 inches. In a bull shot by the Maharaja of Kuch Behar the length from the nose to the tip of the tail was 14 feet 2 inches, and to the base of the tail II feet; the maximum girth being 10 feet 8 inches, and the weight of the head when cut off, 158 lbs.

Distribution.—Typically India, where the range includes the plains of the Bramaputra and Ganges from the eastern end of Assam to Tirhut, and the Terai as far west as Rohilcund, the plains near the coast in Midnapore and Orissa, and also the plains in the Eastern Central Provinces as far south as the Godaveri and Pranhita rivers. A fawn-coloured race occurs in Assam, and smaller varieties in Ceylon and some of the Malayan islands. In a domesticated state, South Europe, Egypt, etc. Native name of male, arna; of female, arni.

Length on outside curve.	Circum- ference.	Tip to Tip.		Widest outside.	1	Locality.	•		Owner.
778	17 5			•••		?			British Museum.
-7 I	21½	343	60		Assam			•	Measured by A. O. Hume, C.B.; shot by A. Forbes.
- 9 70 <u>1</u>	18		64	78	Do.	•			The late Sir A. Campbell-Orde.
65₹	201				Do.				Col. J. Mathie, British Museum.
- 9 64 <u>1</u>	18	42½	60	96	Do.	•			The late Sir A. Campbell-Orde.
-63	18			60	Do.				Do.
62	17	101		•••	Do.	•			Hon. Walter Rothschild.
961 3	153	22 }	45 1		Kuch B	ehar			H. H. the Maharaja of Kuch Behar.
9 61 1	16	22	48	•••	Assam	•			Hon. Walter Rothschild.
60	22		64	72	Central	Provinc	ces		B. Vincent.
60	20	.40	52			3			Sir Edmund G. Loder, Bart.
58§	123	•••				?			British Museum.
58 <u>‡</u>	123					?			Do.
♀ 58	15	46 1	59		Assam				Rowland Ward.
-57	18	•••		•••	Central	Provinc	ces		J. D. Inverarity.
57	181	4112	6 0		D	0.			LieutCol. G. D. F. Sulivan.
-57	15	49	52	•••	Assam		•		A. H. Straker.
- 9 57	141	53	61			?			Sir Edmund G. Loder, Bart.
-56	18	29½		57½		?			J. Whitaker.
56 1	211	40	52 4	•••	Assam		:		A. O. Hume, C.B.
♀ 56	19‡	334	50‡		Kuch B	ehar		•	H.H. the Maharaja of Kuch Behar.
56	15½	55½	58		Do	•			Eyre Coote.
55½	181	29	44	•••		?			J. Carr Saunders.
- 2 55	131		•••	•••	Bhutan	Duars	•		Major-Gen. Alexander A. A. Kinloch.
55	22	55 <u>1</u>	62	66	Central	Provin	ces		L. T. Harris.
54½	18 1	38 <u>‡</u>	487	•••		?			Hume Collection, British Museum.

INDIAN BUFFALO or ARNA (Bos bubalis)—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Widest outside.	Locality.		Owner.
♀ 54½	I 2 ⁸	69 3	70		Assam	•	A. O. Hume, C.B.
54½	19	50	59	64	?		Viscount Powerscourt.
54	20	40	47	55	?		Do.
-54	23	39		59 1	Burma		Pegu Club, Rangoon.
53 ‡	121	23 1	40 §		?		Hume Collection, British Museum.
531	121	•••			Nepal	•	The late B. H. Hodgson, British Museum.
1 53 1	23 (6 ft. 2)	30 ins. at si	42½ houlder)	•••	Kuch Behar .	•	H.H. the Maharaja of Kuch Behar.
-53		•••			?		Lucknow Museum.
514	18 <u>1</u>	22	38 1		Assam	•	Sir Peter Walker, Bart.
-51 8	198	45 8	50		Central Provinces	•	P. II. G. Powell-Cotton.
-50 3	183	55 1	•••	65	Patna State .	٠	LieutCol. F. H. Whitby.
♀ 50⅓	144	40 1	59		Central Provinces	•	Col. M. M. Bowie.
50	18 <u>1</u>	48 1	60		Do.	•	Capt. C. F. Pinney.
50	13 1	33	38 1		Assam	•	Noel Fenwick.
50	17	313	50		Do	•	D. M. Lumsden.
-49 8	171	16 <u>‡</u>	37		Kuch Behar .	•	Count Scheibler.
48 <u>1</u>	18 1	33	51		Central Provinces		Col. F. C. Lister-Kay.
48 1	19	27	47		Assam	•	Major E. T. Paul.
-48 <u>1</u>	134	48 2	57		Central Provinces	٠	P. H. G. Powell-Cotton.
48	19	33	50		Do.		Major H. C. Morland.
48	113	29 2	411		Assam	•	SurgCapt. A. Pearse.
-48	20				Do	•	H. Lennard.
♀ 47 <u>1</u>	134	23	47		?		Major C. B. Wood.
47	19	25	50	•••	?		H. De Prée.
-46 <u>1</u>	20	42 1	55½		Kuch Behar .	•	Prince Hans Henry of Pless.
46 <u>1</u>	20	421	54		Assam		Col. R. Pole-Carew, C.B.
461	134	32	41		Kuch Behar .	•	J. C. O'Donnell.
46 <u>1</u>	172	32	41	•••	?		Dublin Museum.

¹ Measured round outside curves across skull, 9 ft. 11% in. Another pair, 10 ft. 5 ins., circumference of horn at bone, 21 inches.

INDIAN BUFFALO or ARNA (Bos bubalis)—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.		Widest outside.	Locality.	Owner.
46	171	32	41		?	Dublin Museum.
45‡	13	39	49		Kuch Behar	Lord Wolverton.
441	141/2	151	38		Assam	L. Truninger.
44	19	271	46 <u>1</u>		Do	Hon. S. Tollemache.
44	20	28 <u>1</u>	44		Do	A. J. Walter.
♀ 43 2	121	381			Durbangah	Duke of Portland.
Th	e follo	wing	speci	mens	are mostly from	Ceylon :
₽ 35	91	314	37 1	•••	Ceylon	R. Wahrmann.
34 1	151	25	35	•••	Do	Earl Cairns.
♀ 33₺	9	311	37		Do	Do.
317	16	28		38 1	Do	Marquis Camden.
314	16		471	•••	South Australia (introduced)	H. L. Heber Percy.
9 31 1	113	34½		381	Ceylon	SurgMajor G. E. Hale, D.S.O.
302	14	28	334	•••	Do	Count J. Potocki.
30 3	14	22	32	•••	Do	H. E. Lindsay.
30 1	15	22	•••	35 1	Do	A. R. Hay.
29 1	101	26	32		Do	Capt. John Fuller.
26 8	143	28 1	25	•••	Formosa (introduced) The late R. Swinhoe, British Museum.




Head of Bull Anoa, from life.

ANOA (Bos depressicornis).

The smallest member of the ox tribe, characterised by its straight, upright horns, and the frequent presence of white spots on the sides of the head or elsewhere. Although so different in the form of the head and horns from the adult Indian buffalo, the anoa is in these respects much more like the young of the latter; and they are connected to a considerable extent by the tamarau, or Philippine buffalo (Bos mindorensis) of the island of Mindoro. As in all the Oriental bovines, the hair of the fore part of the back is directed forwards in the anoa. Height at shoulder about 3 feet 3 inches.

Distribution.—The island of Celebes.

Length on front curve.	Circum- ference.	Tip to Tip.	Locality.				Owner.		
- 15 8	7 1	$7\frac{1}{2}$	Celebes		•		Paris Museum.		
I 2 3	6	$6\frac{1}{2}$	Do.				British Museum.		
121	6 1	8 <u>8</u>	Do.				Do.		
111	54	7 1	Do.				The late Gen. Hardwicke, British Museum.		
10	5 7	58	Do.	•			Sir Victor Brooke's Collection.		
-9 1	81	81/2	Do.				Dublin Museum.		
8 7	48	44	Do.				Sir Edmund G. Loder, Bart.		



Skull and Horns of Bull and Cow Bison, shot by St. George Littledale.

EUROPEAN BISON or ZUBR (Bos bonasus).

The great elevation of the fore-quarters, the mass of long hair clothing the head, shoulders, and fore-part of the body, together with the peculiar form of the head and horns, the latter of which are cylindrical, serve at once to distinguish the bison from the other members of the ox tribe. There is also a difference in the number of ribs between the bison and the more typical oxen, the number in the former varying from 14 to 15 pairs. In the European species the mass of hair on the fore-quarters is not so long as in its American cousin, the form of the skull is different, and there are marked points of difference in the general appearance which render it easy to distinguish between the two species. Some difference may be noted between Caucasian and Lithuanian specimens, but it is doubtful whether these are sufficient to indicate a racial distinction. Height at shoulder 6 feet 1 or 2 inches. In a bull killed by Mr. St. George Littledale, the length from the nose to the root of the tail measured 10 feet 1 inch, the height at the shoulder 5 feet II inches, and the approximate girth of the body 8 feet 4 inches.

Distribution.—At the present day restricted to the Caucasus and the forest of Bielowitzka in Lithuania; the herds in the latter district existing in a protected state. The name aurochs, so commonly applied to the bison, properly belongs to the extinct wild ox of Europe.

EUROPEAN BISON or ZUBR (Bos bonasus)-continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Locality.			Owner.		
181	121	134	191	Lithuania .			British Museum (presented by the Czar of Russia).		
-18.2	10.8	17.5	20.14	Bielowitzka	•	•	Prince Henry of Liechten- stein.		
-18	121	16	20	Caucasus .	•	•	St. George Littledale.		
171	10	181	•••	Bielowitzka			Major Algernon Heber-Percy.		
17.4	13.3	13.4	20.10	Do.		•	Prince Henry of Liechten- stein.		
16 2	121	217	241	?			Sir Edmund G. Loder, Bart.		
₽ 15‡	8	6		?			Major Algernon Heber-Percy.		
9 13	81	92	14	Caucasus .	•		St. George Littledale.		



Skull and Horns of American Bison.

AMERICAN BISON (Bos bison).

Some of the points distinguishing this species from the European bison have been mentioned under the head of the latter, but it may be added that in the typical race of the former the horns are shorter, thicker, blunter, and more sharply curved. In the skull of the American animal the sockets of the eyes have a more tubular form. Height at shoulder about 6 feet; weight from 15 to 20 cwt.; an adult bull weighed by W. T. Hornaday scaled 1727 lbs.

Distribution.—The greater portion of Western North America, ascending to the Great Slave Lake, and descending to New Mexico and Texas; now nearly exterminated. American writers recognise two races (or species), the prairie bison (B. bison typicus) and the larger wood-bison (B. bison athabascæ) of the forest highlands of the North-West.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside spread.	Locality.		. Owner.
-21½	15‡		35 outside	Northern Monta	na	W. F. Sheard.
20 7	15		30½	Wyoming .		Hon. F. Thellusson.
-201	16 1	$33\frac{1}{2}$?		W. H. Root.
-19	121/2		•••	W. Montana		P. Liebinger.
18 7	144		16 7	Do.		The late J. S. Jameson.
18 <u>‡</u>	14	26]	29	Sioux Country		Sir Greville Smyth, Bart.
-18	14		•••	Montana .		F. Sauter.

AMERICAN BISON (Bos bison)—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Locality.	Owner.
174	128	15 1	•••	}	H.R.H. the Duke of Saxe-
-171	I 2 ½			SW. Montana	Coburg and Gotha. Theodore Roosevelt.
171	12	•••	25½	Wyoming	H.R.H. le Duc d'Orléans.
171	131/2	21	•••	?	Viscount Powerscourt.
178	11 8	108	171	?	British Museum.
-17	14	171		Yellowstone, Montana	Count E. Hoyos.
16 §	141	24		Bighorn Mts., Wyoming	Moreton Frewen.
1 16 1	121	198	•••	Colorado	Sir Edmund G. Loder, Bart.
16 1	131/2	14‡		?	Duke of Portland.
16 1	15 7	25 3		Colorado	Sir Edmund G. Loder, Bart.
151	148	•••	194	Wyoming	St. George Littledale.
-15.8	12.14	15		Indian Territory near Texas	Prince Henry of Liechten-
14	•••	121		North Park, Colorado	stein. Col. Ralph Vivian.
131	131	171		?	G. Wrey.
138	12	•••		?	Hon. Walter Rothschild.

¹ Wood Bison.

YAK 407



Skull of Male Yak. From a specimen in the British Museum, presented by A. O. Hume, C.B.

YAK (Bos grunniens).

The plateau of Tibet is remarkable for the number of its peculiar mammals, among which is the yak. Apparently its nearest relatives are the bisons, but the yak has not the great elevation of the withers in comparison with the hind-quarters so distinctive of the latter, and the long hair forms a fringe on each side of the flanks, shoulders, and thighs, as well as a tuft on the chest, while the tail is clothed with a huge mass of similar long hair, forming, when cut off and mounted, the well-known "chowries." Yak horns are much larger than those of living bison, and have a totally different curvature; while there are also important differences in the skull. Height at shoulder from about 4 feet 10 inches to at least $5\frac{1}{2}$ feet; girth behind shoulder, 9 feet $1\frac{1}{2}$ inches; length from between horns to base of tail, 8 feet $\frac{1}{2}$ inch;

tal. 3 feet 23 inches; from between horns to nose I hot 32 hiches (Cago H. M. Bidon ph). Weight about 1140 lbs. Whit yak are neclearly blackish brown in colour, any trace of white indicating dementication, and probably cross-breeding.

Dutroution.—The plateau of Tiber, part of the Kansu province of China, and Northern Ladak, at elevations between about 14,000 and 20000 feet. The peculiar grunting cry from which the animal takes its name is developed only in the domestic breed.

dergrain meuro inemi	Coresto.	Tyn Ty.	Localty.		Iwas.
39			<i>:</i>		Lacines X sem.
194	123	#i	Kaenisen Mac.		A. O. Hime, C.E., shot by late
<i>i</i> 4	:7	19	in .		A. Drigersa. Hume Coleman, Brit. Museum
353	15	16	•		pickers up by Mr. Carey. E. L. Fheips.
35 i	142	17	Tibet.		et. George Littleinie.
34	12	¥4	Ladak		Hon. Walter Rochschild.
7,2 }	164	18}	Dr.		Cape. H. M. Billish.
32 [14	171	Do.		P. H. G. Powell-Cotton.
32	131	151	Do.		H. C. V. Hunter.
32	141	9	Do.		L Morse.
3,2	16	17	Do.	-	P. Church.
314	141	18‡	Chang Cherano .		P. H. G. Powell-Cotton.
314	16	163	Do	•	Rowland Ward.
31	14		Do		Col. F. C. Lister-Kay.
31	131	14	Ladak		Arnold Pike.
y 4	13½	101	Do		Sir Robert Harvey, Bart.
y,	13	152	Chang Chenmo .		H. Z. Darrah.
29	14	12	?		Hume Collection, British Museum.
21) 1	131	11	Chang Chenmo .		Lieut-Col. G. D. F. Sulivan.
29	15	9	Do. .		Sir Edmund G. Loder, Bart.
29	131	1,3 1	Ladak		Arnold Pike.
289	15	13	Do.		G. B. Milne.
281	158	141	Do.		P. Church.
28	16	17 7	Tibet		Major-General Alexander A. A. Kinloch.

YAK

YAK (Bos grunniens)—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Locality.		Owner.
28	13	28	Tibet		J. Benett Stanford.
28	131	291	Near Manasarow Lake	rar	Capt. B. H. Boucher.
27 8	113	18	Lake ?		The late B. H. Hodgson, British Museum.
27½	131	16 <u>1</u>	Tibet		Capt. C. B. Vandeleur.
27	13	151	Ladak		Major C. S. Cumberland.
26½	131	19	Chang Chenmo .		Col. J. Biddulph.
254	12	163	Ladak		Major J. A. Orr-Ewing.
251/2	13	17	Do		Capt. H. H. P. Deasy.
$24\frac{1}{2}$	111	14	Tibet		T. W. Greenfield.
♀ 24 <u>1</u>	81	54	Ladak		G. B. Milne.



Head of Bull Yak.



Head of Male Gaur.
(After Forsyth.)

GAUR (Bos gaurus).

This splendid bovine, the miscalled bison of Anglo-Indian sportsmen, is the typical representative of a group of three oriental species nearly related to the domestic ox, but presenting certain well-marked points of difference. Among these may be noted the shorter head and tail, the frequently elliptical section of the horns, and, above all, the presence of a more or less distinct ridge running from the withers to the middle of the back, where it terminates in a sudden step. In the gaur this ridge is very strongly developed, but the most distinctive feature of the animal is the great arch on the crown of the head between the horns, which bends forward to communicate a concave profile to the forehead. With the exception of the white "stockings"

GAUR 411

common to all the members of the group, the colour of the gaur is uniform; but the much flattened horns are of a peculiar yellowish green tint at the base. The height of adult bulls at the shoulder varies from about 6 feet to 6 feet 4 inches, though specimens of not more than 5 feet 5 or 6 inches are killed.

Distribution.—The forest hilf-tracts of Peninsular India, Assam, Burma, and the Malay Peninsula, as well as the forests along the outer Himalaya as far west as Nepal. South of the Ganges, where it has not been exterminated, the gaur inhabits suitable districts in Chutia Nagpur, Orissa, the Northern Circars, Central Provinces, Hyderabad territories, and all the Western Ghats.

Length on outside curve of longest horn.	Circum- ference at base.	Tip to Tip.	Widest inside.	Widest outside.	Locality.	Owner.	
-391	201	183	•••	43	Salwin, Burma	Bombay Natural History Society's Museum.	
-39	19‡	18 1	32		N. Travancore.	Arthur W. Turner.	
-35	18	•••	•••		S. India	The late General Douglas Hamilton.	
34 8	20	25		40]	Vardi Mullay .	Baron von Massow.	
-34	19‡	33 1			?	Measured by Colonel J. Biddulph.	
33 8	18	24		41	Pulny Hills, Madura District	J. D. Goldingham, Bethnal Green Museum.	
	177	28 3	•••	43½	Travancore .	H.H. the Maharaja of Travancore, G.C.S.I.	
	174	25	•••		Do	Do.	
33½	18 1	25	33 1	421	Do	T. W. Greenfield.	
-33½	21	•••	321	382	Mysore	Surgeon-Captain C. W. H. Whitestone.	
33 1	18	23½	341		Kuch Behar .	Maharaja of Kuch Behar.	
33	171	201	32 1	•••	?	Capt. P. Z. Cox.	
-33	14			34	Western Ghats	J. D. Inverarity.	
$32\frac{1}{2}$		27			Do.	LieutCol. L. L. Fenton.	
-32	19	27			Assam	Major James Grant.	
-32	18	•••		• • •	Burma	Indian Museum.	
317	171	218	321		Travancore Hills	A. O. Hume, C.B.	
-311	18	29		43	?	Bombay Natural History Society (<i>Proceedings</i>).	
-31½	17	21	32 1	•••	?	Dublin Museum.	
314	171	225	27½	401	Pulny Hills, Madura District	J. D. Goldingham, Bethnal Green Museum.	
31	191	241	30 3	$37\frac{1}{2}$	Mysore	Viscount Powerscourt.	

GAUR (Bos gaurus)—continued.

Length on outside curve of longest horn.	Circum- ference at base.	Tip to Tip.	Widest inside.	Widest outside.	Locality.	Owner.
30 1	191	16		35	Travancore .	Lord Douglas Compton.
30 1	18 1	$22\frac{1}{2}$	37‡	•••	Do	A. T. Mackenzie.
30	19‡	11	32 1		Kuch Behar	H. H. the Maharaja of Kuch Behar.
-30	20	15	26	•••	?	G. Beck.
-30	16		38	41	Central Provinces	LieutCol. J. W. H. Flana- gan.
-30	19	16	32		Travancore .	J. D. Rees.
29 8	18 1	30	34	•••	?	Sir Edmund G. Loder, Bart. (Sir Victor Brooke's Collec-
29 3	18 5	251	30 3	•••	Hassanoor Hills, S. India	tion. (Shot by the late Gen. Douglas Hamilton.)
-291	19	22 1 8	31 1	36 7	Mysore	Captain M. M'Neill.
-29½	19‡	16‡	26½	•••	Narbada Valley	Major-General Alexander A. A. Kinloch.
-29½	18		•••	33	Central Provinces	
$29\frac{1}{2}$	174	281	39		Siam	A. Waley.
29½	17	32½	43		Coimbatore, S. India	P. Church.
29 1	127		18		Nepal	The late General Hardwicke, British Museum.
-29	22				?	Otho Shaw.
-29	171/2	•••		35	North Kanara .	LieutCol. L. L. Fenton.
29	16	30		41½	Assam	A. J. Walter.
283	18 1	141	30		Travancore .	Lord Wenlock, G.C.I.E.
283	174	23‡	31	•••	South India .	Captain H. M'Micking.
28½	154	18 §	28 §		?	Major-General Sir Arthur Ellis, K.C.V.O.
28½	18	15%	33½		Travancore .	H. L. Cottingham.
28 <u>1</u>	15‡	17	27		?	Hume Collection, British
-281	181	28 1		38₫	Raipur, C.P	Museum. Captain M. M'Neill.
28	18	$22\frac{1}{2}$	34 3		Assam	Hon. S. Tollemache.
28	16	101	24		Central Provinces	Martyn Kennard.
28	193	34½	43		Upper Burma .	C. W. A. Bruce.
28	16 <u>1</u>	152		3112	?	H. W. Keys.
28	151	16‡		31	Khandesh .	A. Cumine.
-28	17½	17		36 <u>1</u>	Mysore	Veterinary-Captain G. H. Evans.

GAUR (Bos gaurus)-continued.

Length on outside curve of longest horn.	Circum- ference at base.	Tip to Tip.	Widest inside.	Widest outside.	Locality.	Owner.
278	178	3 2 1	32 3		Central Provinces	Major C. S. Cumberland.
278	194	37	41		Kalkerray, S. India	Captain W. E. Fairholme.
274	18 <u>8</u>	24 ⁸	30½	•••	?	Major G. A. L. Carew, D.S.O.
-27 1	20½	26 <u>1</u>		36½	?	Major-General W. Rice.
27½	17	26 1	36	•••	Assam	L. Truninger.
27½	19	15	314	•••	Travancore .	Captain Hon. E. Baring.
27½	171	201	35	•••	Tezpore, Assam	A. Y. Thomson.
271	18‡	20 1	34	•••	Travancore .	Captain Hon. E. Baring.
-27	16 3	22	30½		South India .	H. W. Murray.
-27	15		17		Panarel, Central India	Captain A. Hicks-Beach.
26 7	15 7	16 1	27		?	Hon. Walter Rothschild.
26 2	147	138	241	<i>.</i> :.	Sahaydri Moun- tains, S. India	Captain W. Tompson, British Museum.
26½	17	171	•••	321	Central Provinces	
26]	18 1	26 3	•••	36 1	?	Major R. H. Fraser.
26	181	238	324	•••	Central Provinces	G. de H. Smith.
26	19	16 1	321/2	•••	Do.	Captain C. F. Pinney.
26	172	198			?	Lord Wolverton.
26	15	26 1	36	•••	Central Provinces	Captain John Fuller.
26	15	33 1	•••	40	Do.	C. F. Egerton.
26	174	27	36		Do.	Colonel M. M. Bowie.
26	18	23	35	•••	Burma	Captain S. L. Robinson.
· 25½	17	21½	28 1	331	S. India	Captain C. S. Timins.
25	184	181	29	•••	?	Lord Elphinstone.
25	18	234	334	•••	Kanara	Captain G. J. Fitzgerald.
25	172	18 1	34	•••	Central Provinces	Dr. W. P. Y. Bainbrigge.
241	154	24§	34	•••	?	H. De Prée.
-24	18	34	•••	38 2	?	J. D. Inverarity.
-24	19	27	34	•••	Burma	W. F. Loftus-Tottenham.
23	15	22	$33\frac{1}{2}$	•••	Assam	Major E. T. Paul.

GAUR (Bos gaurus)—continued.

Length on outside curve of longest horn.	Circum- ference at base.	Tip to Tip.	Widest inside.	Widest outside.	Locality.	Owner.
♀ 22	101	14§	27		?	H. De Prée.
♀ 20½	101	I 2			Madras	Major J. W. M. Cotton.
- 9 19 <u>1</u>	117		•••		N. Kanara .	Lieutenant-Colonel L. L. Fenton.
♀24	131	13	22	27	N. Travancore Hills	A. O. Hume, C.B. (See illustration.)



Skull and Horns of Cow Gaur, from A. O. Hume's specimen

GAYAL 415



Skull and Horns of Bull Gayal, from A. O. Hume's specimen.

GAYAL (Bos frontalis).

Chiefly known in a half-domesticated condition, the gayal is a rather smaller animal than the gaur, of a blacker colour, with a large dewlap, and with a straight line between the bases of the massive horns, which are more divergent, less curved upwards, more nearly cylindrical in section, and of a darker colour. The head, too, is much shorter, with the forehead very broad and flat.

Distribution.—Apparently the Tenasserim district in a wild state, but kept in a semi-domesticated condition by many of the hill-tribes of Assam and Chittagong. Gayal have been crossed with the American bison, the resulting progeny being fertile.

Length on outside curve.	Circum- ference.	Tip to Tip.	Locality.	Owner.	
15	1112	26 §	?	British Museum.	
141	131	28	Tenasserim Hills	A. O. Hume, C.B. tration.)	(See illus-
-14	14	•••	?	Indian Museum.	
12§	13‡	27 ³	?	British Museum.	



Head of Male Burmese Banting. (From the Proc. Zool. Soc., 1898.)

BANTING (Bos sondaicus).

The third and last member of the gaur group departs less widely from the type of the common ox than do the two others, the ridge on the withers being less developed, and the horns almost cylindrical. The cows, too, are always reddish coloured, although the bulls may be black, and in the latter sex at least there is always a large white patch on the rump. Very distinctive of the species is the presence of a horny shield on the crown of the head between the bases of the horns. Height at shoulder about 5 feet 9 inches.

Distribution.—Burma, the Malay Peninsula, Borneo, Java, Bali, and perhaps Sumatra. At least two distinct races of the banting are distinguishable. First, the true banting, or Java ox (Bos sondaicus typicus), from Java, and perhaps some of the other Malayan islands and the Peninsula. In this race the old bulls become of a deep blackish brown colour. The same tint is characteristic of the old male banting in Borneo, but the horns are directed more uprightly.

The second well-defined race (B. sondaicus birmanicus) inhabits Burma, and may extend northwards to Manipur. Old bulls retain the reddish tint of the cows throughout life, showing more or less of gray on the head.



Skulls of Gaur and Banting. Shot by Vet.-Capt. G. H. Evans.

Length on outside curve.	Circum- ference.	Tip to Tip,	Widest inside.	Locality.				Owner.
$-33\frac{1}{2}$	17	26 7	35	Tammu, Burma		Upper		Surgeon-Capt. H. S. Wood.
-30	17		•••	Java	٠.			Indian Museum.
-28 1	171	30½	36	Burma				VetCapt. G. II. Evans.
-28½	15	26½	36 3	Java				H. Van-Son.
26 1	16]	211	28	Do.				Sir Edmund G. Loder, Bart.
-26	18	172	27	Burma				C. W. A. Bruce.
25 8	16 §	20 ⁷ 8	26 3	Siam				H. C. V. Hunter.
-25½	131	29	32	Burma				W. H. Prendergast.

2 E

BANTING (Bos sondaicus)—continue	BANTING	(Bos	sondaicus)-	-continue
----------------------------------	---------	------	-------------	-----------

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Locality.	Owner.
$25\frac{1}{2}$	191	24½	30	Burma	C. W. A. Bruce.
24 <mark>8</mark>	121	154	241	Java	British Museum.
-24 ¹ / ₂	141	30	35	Pegu	VetCapt. G. H. Evans. (See illustration.)
218	12‡	13 1	19]	Borneo	H. B. Low, British Museum.
20 5	124	18 1	22 1	Do	Do.
20	16	22 3	· 24½	Kudah, B. N. Borneo	H. Ll. Davies.
♀ 20	12	$21\frac{1}{2}$	28	Upper Burma	C. W. A. Bruce.
-19‡	114	17	20	Borneo	Sir Edmund G. Loder, Bart.
19	115	18	218	Java	The late H. J. H. Platt.
18	131	121		Borneo	Sir Edmund G. Loder, Bart.
175	10 8	7 1 8	141	British North Borneo	W. B. Pryer, British Museum.
171	10 1	171	208	Siam	H. C. V. Hunter.
16 1	103	9 1	138	Sarawak	British Museum.
16 1	128	171	18 <u>1</u>	N. Borneo	A. D. Boden.
♀ 15½	71/2	71/2	16 1	Tammu, Burma .	Surgeon-Capt. H. S. Wood.
♀7 1	7‡	6 <mark>2</mark>		Java	The late H. J. H. Platt.

DOMESTIC OXEN (Bos taurus and B. indicus).

The domesticated cattle of Europe are the descendants of the primitive wild ox, or aurochs (a name frequently misapplied to the bison), of Europe and North Africa (Bos taurus primigenius), now completely extinct in the wild state, although it survived on the continent till the Middle Ages. On the other hand, the humped oxen (zebu) of India and the Galla cattle of Africa (Bos indicus) appear to trace their origin to a totally distinct species, also extinct in the wild state.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Locality.	Owner.
81 <u>‡</u>	18 <u>‡</u>	1031		N'gamiland .	The late W. C. Oswell.
-7	17	100	124	Do	A. Ohlsson.

DOMESTIC OXEN (Bos taurus and B. indicus)—continued.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest inside.	Locality.	Owner.
57	191	82	•••	Bechuanaland	Sir Edmund G. Loder, Bart.
56 1	171	76]		South Africa.	R. A. Cooper.
47	15 §	218	• •	Abyssinia (Galla Ox)	The late H. Salt, British Museum.
44	127	70½ (about)		(Galla Ox)	British Museum.
42 lg	235	228		Central Africa	The late Col. Denham, British Museum.
413	14	524		Madagascar .	Bethnal Green Museum.
40 1	124	60 ⁷		Vienna (Polish Bull)	British Museum.
$38\frac{1}{2}$	101	543		Italy	Bethnal Green Museum.
354	122	59½	•••	Cape of Good Hope	Do.
30 1	128	35		Spain	Do.
298	I I 🛔	28 7		Gambia .	A late Earl of Derby, British Museum.
171	108	30 1		Buenos Aires (Niata Cattle)	G. Claraz, British Museum.
171	10 1	25 1 8		Gambia .	A late Earl of Derby, British
17 1	103	254		India	British Museum.
12	8	16 1		Angola .	Sir Victor Brooke's Collection.
111	71/2	single horn	•••	Nepal	The late B. H. Hodgson, British Museum.

The following specimens belong to the so-called Wild Cattle of certain British parks, which are, however, certainly the descendants of at least partially domesticated breeds.

Length on outside curve.	Circum- ference.	Tip to Tip.	Widest Inside.	Locality.	Owner.
18 1	91	36½	•••	Chartley Park	Hon. Walter Rothschild.
181	7	34 8		Do.	Major James Grant.
181	108	20	$21\frac{1}{8}$	Chillingham Park	Earl of Tankerville, British Museum.
\$ 18 1	7	10 <mark>1</mark>	15 8	Do.	Do.
151	98	178	18 1	Northumber- land	Duke of Hamilton, British Museum.
15	71/2	201		Chartley Park	Captain G. W. Hill, R.N.

HIPPOPOTAMUS (Hippopotamus amphibius).

Swahili name Kiboko. Abyssinian name Gumare.

Danakil name Dul. Boer name Zee-koe.

Galla name Robi. Swazi name Imvubu.

Basuto name Ikubu.

Such a familiar animal as the uncouth and unwieldy hippopotamus—the largest member of the swine group—requires but little in the way of description here. It is widely distinguished from the pigs and warthogs by the broad and rounded muzzle, so unlike the disc-shaped snout of the latter; and consequently forms a family by itself. The tusks and molars are likewise of a totally different and distinctive type; while the feet have four sub-equal toes with symmetrical, rounded nails. In all its organisation the hippopotamus is beautifully adapted for a sub-aquatic life; the eyes and nostrils forming the highest points of the head, and thus allowing the creature to come up and breathe with the least possible exposure of its body. The weight of a full-grown bull hippo is at least three tons.

Distribution.—Formerly this animal frequented most of the rivers of Africa south of the Sahara, but it has long since been exterminated from the lower reaches of the Nile, and is daily becoming scarcer in the South African rivers. In the Zambesi, where it is less easily attacked than in the smaller rivers of Mashonaland, it is still abundant, as it also is in the Chobe and neighbouring rivers, as well as in Lake Ngami; and even in the lower reaches of the Orange River a few are still to be met with. North of the Zambesi these animals occur in great numbers. Hippos are chiefly hunted for the sake of their hides, which are manufactured into sjamboks, or raw-hide whips. Their tusks have also a certain commercial value, although not so great as formerly, when they were employed for artificial teeth.

In Liberia and some other parts of the West Coast there occurs the much smaller pigmy hippopotamus (*H. liberiensis*), an animal measuring only some six feet in length, and possessing more the habits of a pig. It also differs from the common species by having, as a rule, only one, instead of two, pairs of incisor teeth between the tusks.

A hippopotamus, shot on the Shiré River, British Central Africa, by F. Vaughan Kirby, measured as follows:

	ft.	in.		ft.	in.
Total length, nose to tail	Ι4	2	Length of tusks .	2	5
Tail	I	8	Circumference of tusks		81
Vertical standing height	3	$10\frac{1}{2}$	Weight per pair "	$14\frac{1}{2}$	lbs.

These tusks measured 31 and $30\frac{1}{4}$ inches, with circumference $8\frac{3}{4}$ inches, after they were extracted from the skull.

Length round out- side curve of tusk.	Circumference.	Locality.	Owner.
1-51	9 (malformed)	S. E. Africa	J. Lamont.
141	8 Do.	Tana River, East Africa	J. Benett Stanford.
1-38 protruding from jaw		Nyasaland	Major P. W. Forbes.
¹ 37½	81	Shiré River	Comdr. A. T. Hunt.
32	9 <u>‡</u>	Lualaba, Central Africa	S. L. Hinde.
314	108	?	Sir Clement Hill, Bart.
31§	9 1	?	Sir Edmund G. Loder, Bart.
² 31 ¹ / ₄		S.E. Africa	F. Vaughan Kirby.
30½	91	St. Lucia Bay	Hon. Charles Ellis.
30	9 (weight 15 lbs.	Do	Do.
30	pair) 9	Shiré River	Staff-Surgeon J. Dowson, R.N.
30	9	Zambesia	E. W. Tompson.
-30	(weight 11 lbs.)	Katungas, Shiré River	E. B. Vertue.
29 ³	83	Do.	F. C. Selous.
29½	8	Do.	Dr. W. P. Y. Bainbrigge.
29	8	Do.	Do.
29	8 1	Shiré River	F. Vaughan Kirby.
29	91/2	?	Rowland Ward.
27 l		Atbara River, North- East Africa	W. D. James.
27	8	East Africa	E. Gedge.
26 ½	8 <u>1</u>	Pungwe River	Earl of Dunmore.
$-26\frac{1}{2}$	81	Zambesia	James J. Harrison.
25½	8	Nyasaland	Alex. R. Alston.
193	•••	Do	Sir John Kirk, K.C.B.
-II 1 protruding from jaw	81	Shiré River	Dr. Percy Rendall.

¹ Malformed.

² Straight tusk 19 ins. long.



WILD BOAR (Sus scrofa and S. cristatus).

Most of the European and Asiatic species of wild swine, forming the typical group of the genus Sus, are so like each other, that it is often a matter of difficulty to discriminate between them, and naturalists are not yet in accord as to the number of species which should be recognised. The European wild boar is a large coarsely-haired species, with an undercoat of woolly fur, no warts on the face, and standing about 33 inches at the shoulder. It lacks the crest or mane of long black bristles running from the nape down the back in its Indian cousin; and the last tooth in the lower jaw is of a rather less complex structure than in the latter. The wild boar of India (S. cristatus) is dear to the heart of the Indian sportsman, though he does contemptuously call its chase "pig-sticking."

"Pig-sticking" is considered quite as dangerous, if not more so than tiger-shooting. It is not a game of long bowls, but a close personal attack on a very fierce and pugnacious animal, endowed with strength, swiftness, and much tenacity of life. It takes a good horse and a good man to try conclusions satisfactorily with an old gray boar over nasty country, and the weapon used is only a spear. There are parts of the hilly country in India where it is impossible to ride, and here the rifle may take the place of the spear; but then there is not much glory in shooting a pig. A solitary boar is frequently a morose and dangerous animal.

A Spanish boar killed by H.R.H. le Duc d'Orléans weighed 302 lbs. Distribution.—Europe, Asia Minor, and North-East Africa.

Length outsic of tusk Right.		e Length of tusk out of gum.		Local	lity.			Owner.
-1 1 ½			Caucasus					Col. Veernhof.
-111			Do.					Capt. Robert Finnie.
9 3			Asia Mino	or			•	Admiral Sir Michael Culme Sey- mour, Bart.
² 9 §		•••	Caucasus	•	•		•	Prince Demidoff.
9 1			Do.	•	•	•	•	Do.
-9			Do.			•		Capt. Robert Finnie.
9			Albania	•				Sir Reginald Cathcart, Bart.
-8§			Algeria				•	Sir Edmund G. Loder, Bart.
1_81	81	3	Near Bon	a, A	lgeria			Viscount Edmond de Poncins.
-81/2	81	1 §	D	ю.				Do.
-8			Albania					Capt. Robert Finnie.
8	2 ½		Russia					Count J. Potocki.
		1 Wei	ght 275 lbs. (clean.			2 V	Veight 372 lbs.
Length outsi		1	Locality.					Owner.
-14 ³ (malformed -14 ⁸	P d)	urneah, Low	_				Janu	P. Carter, recorded in <i>Field</i> , 19th ary 1895. (See illustration, p. 422.) ir Neville Chamberlain.
-12 <mark>3</mark>	Ja	va				I	I. Vai	n Son.
-10 §	N	orth Kanara	ι			В	omba	y Natural History Society's Museum.
-10½			?			N	feeru	t Tent Club.
-10	Iı	ndia				Т	he la	te Sir Samuel Baker.
-9 3			;			N	l eeru	t Tent Club.
9 8			?			D	r. Tr	avers.
91	В	urma .				L	angfo	ord Whitehouse.
98	N	orth Kanara	ι			L	.ieut.	Col. L. L. Fenton.
98	С	entral Provi	nces .			N	I ajor	J. S. Ashby.

WILD BOAR (Sus scrofa and S. cristatus)—continued.

Length outside curve of tusks.		L	ocality		Owner.	
91	Central Pa	rovin	ces			The late J. Moray-Brown.
-84	Do					Col. F. D. Lugard, C.B., D.S.O.
84	Jumna Va	lley				Capt. F. C. Quicke.
81	Bengal					J. M'Kie.
-8. 30	Assam					Viscount Edmond de Poncins.
-8 1	Bikanir					Maharaja of Bikanir.
-7.85	Assam					Viscount Edmond de Poncins.
8			?			Capt. P. A. Bainbridge.
71	Mhou .					H. Dibble.
7 1			;			Maharaja of Bikanir.



Head of Wild Boar.

CAPE BUSH-PIG (Sus [Potamochærus] chæropotamus).

Bosch-vark of the Boers. Ingulubi of the Swazis and Zulus.

The bush-pigs, or river-hogs, of Africa and Madagascar form a peculiar group of swine characterised by having only 42, in place of 44, teeth, small tusks, and a large ridge-like prominence on each side of the face, due to the presence of a ridge of bone on the sheath of the tusk. The ears may be surmounted with tufts of long hair. various species are best distinguished by the character of their skulls, colour forming a very uncertain guide. The Cape bush-pig, or boschvark, is very generally gray, but the late Sir Andrew Smith stated that "scarcely any two specimens of this species exhibit the same colours; some are a brownish black variegated with white, and others are almost entirely of a light reddish brown or rufous tint, without any white markings; indeed, such are the varieties that it is scarcely possible to say what are the prevailing colours." In British Central Africa, where they have been wrongly identified with the West African species, they are invariably reddish. Height at shoulder about 31 inches; weight, 35 lbs. Lower tusks average 6 to 7 inches long, and a good specimen shot by F. Vaughan Kirby has tusks protruding out of jaw 48 inches.

Distribution.—South and South-East Africa.

WEST AFRICAN BUSH-PIG or RED RIVER-HOG

(Sus [Potamochærus] porcus).

In this species the colour is always some shade of rufous, either shining brownish red with a tinge of yellow, or dark reddish yellow with black on the forehead, ears, and limbs, the mane of the back, part of the margins of the ears, tips of the long tufts of hairs with which they are surmounted, and streaks above and below the eyes white. Distribution.—West Africa, from Angola to Senegambia, and eastwards

Height at shoulder.	Weight.	Locality.	Owner.
-23 1	35 lbs.	Shiré River, British Central	Dr. Percy Rendall.

to Monbuttu.


BABIRUSA (Babirusa alfurus).

The pig-deer (to translate its Malay name) is not the least notable of several remarkable animals restricted to Celebes; the peculiar form and position of the upper tusks of the boars rendering them almost comparable to horns. Unlike other pigs, in which they curve upwards from the sides of the lips, the tusks grow from the centre of the muzzle, piercing through the skin, and as they are not worn by the lower pair, attain extraordinary dimensions. Both pairs are quite devoid of enamel; the lower ones growing from the sides of the jaw in the ordinary manner. The other teeth are somewhat less numerous than in ordinary pigs. In other parts of its organisation the babirusa is, however, very like the latter, although its nearly naked skin is remarkable for its coarse and rugged nature, being almost comparable to the bark of a tree. The height at the middle of the back, the highest point of the animal, is about 42 inches. Unlike the rest of its tribe, the female babirusa produces only a pair of young at a birth, which are of the same uniformly slaty hue as their parent. Babirusa are confined to the island of Celebes, where they afford good sport to the natives, who drive them into nets and then spear them.

Dr. F. H. H. Guillemard recorded in the Cruise of the "Marchesa":

Weight, male, 128 lbs.; female, 85 lbs. Height at shoulder, $27\frac{1}{2}$ in.; female, $25\frac{1}{2}$ in.

Len	gth.							
Upper tusks.	Lower tusks.		Loc	ality.		Owner.		
-17		Celebes				H. Van Son.		
-14½	•••	Do.			<i>'</i> .	Dr. F. H. H. Guillemard.		
-91 ¹ 8	7.%	Do.				Dr. Percy Rendall.		



Tusks of Male Wart-Hog, F. H. Barber's specimen.

WART-HOG (Phacocherus æthiopicus).

Bango or Nguruwe of the Swahilis.

Ikulubi of the Basutos.

Indaigazana of the Swazis and
Zulus.

Karkary of the Somalis.

Ngolobwi in the Barotse country.

Ngolobwi in the Lake Ngami country.

Ngolobwi in the Chilala and Chibisa countries.

Vlak-vark of the Boers.

For downright ugliness the African wart-hog is hard to beat, and as it is well armed and possesses a bad temper, it is in all respects an undesirable acquaintance. The name is derived from the presence of three pairs of wart-like protuberances on the sides of the face between the eyes and the tusks; the head itself being characterised by the disproportionate length and flatness of the face. Unlike the true pigs, the stout upper tusks are longer than the lower pair; the inferior surfaces of their basal halves being worn to smooth facets by the points of the latter. Another peculiarity of the upper tusks is that they have no enamel, except at the extreme tips, which are soon worn off by use. But a further peculiarity is presented by the last molar teeth of each jaw, which, together with the tusks, are often the only teeth remaining in very old animals. They are both long and tall,

consisting of a number of closely-packed cylindrical columns of enamel, which, when worn, present a characteristic pattern. Except along the neck and back, where it carries a mane of bristly hair, the skin is nearly naked; and the young differ from those of ordinary pigs in being neither striped nor spotted. Height at shoulder, 30 inches.

Distribution.—The wart-hog is typically an inhabitant of South and South-East Africa. Animals of the same genus extend, however, right through East and Central Africa to Abyssinia; those from the latter country having been described as a distinct species, under the name of *P. africanus*. Not improbably, however, the northern animal is only a local race of the southern form. Warthogs, possibly from their habit of going to ground when pursued, are but seldom hunted with the spear; "pig-sticking" being, in fact, a sport practically unknown in Africa. The lower tushes seldom exceed 6 inches on the front curve.

Length on outside curve.	Length exposed from gum.	Locality.	Owner.
⁻²⁷ }		Annesley Bay.	Capt. Ralph Berners, R.N.
20	173	South Africa	F. H. Barber.
-16 <u>‡</u>	131	?	J. Whitaker.
$\frac{-15\frac{1}{2}}{15}$	131	Matanari Bush	F. Vaughan Kirby.
-15	•••	South-East Africa	W. Russell Bowker.
142		British Central Africa .	James Yule.
	134	Zambesia	R. J. Cuninghame.
1311 ₀		Somaliland	B. R. M. Glossop.
-13		Timbabti River, S.E.	W. Russell Bowker.
	13	South-East Africa	F. C. Selous.
	114	Somaliland	Viscount Edmond de Poncins.
-12	•••	South-East Africa	James Meldrum.
-II ½	•••	Do	Julius Jeppe.
-II 1 ½	•••	Do	James J. Harrison.
-111	•••	Somaliland	J. D. Inverarity.
•••	103	Mashonaland	S. Chillingworth.
-11	•••	East Africa	Count Scheibler.

WART-HOG (Phacocherus æthiopicus)—continued.

Length on outside curve.	Length exposed from gum.	Locality.	Owner.
11	•••	British Central Africa .	R. Skeffington Smyth.
·	108	Sabi Flats, Transvaal .	Dr. Percy Rendall.
	101	Near Ruo River, South- east Africa	C. C. Bowring.
10}	•••		W. R. Bindloss.
10g	•••	South-East Africa	F. Vaughan Kirby.
•••	9#	•••	W. W. Ashley.
	9	Pungwe	Count E. Hoyos.
	9	Somaliland	Prince Boris Czetwertynski.
•••	9	Do	Capt. F. C. Quicke.
		Lower Tusks.	
-9 1		Somaliland	J. D. Inverarity.
$-9\frac{1}{2}$		Do	Viscount Edmond de Poncins.



Head of Male Wart-Hog.



GREAT INDIAN RHINOCEROS (Rhinoceros unicornis).

In addition to being the giant among its Asiatic kindred and possessing but a single horn, this huge rhino is specially characterised by the form of the folds in its hide, and the large tubercles on the foreand hind-quarters, which look as though the skin had been fastened to the body by means of rivets. A fold before and behind the shoulder marks off one large triangular shield on each side, while another fold before each thigh separates a large rump-shield; the saddle-shaped body-shield being defined by the fold behind the shoulder and the one in front of the thigh, both of which extend across the back. Very characteristic, too, are the great folds which form heavy rings of skin round the neck. Although the tubercles are largest on the fore- and hind-quarters, they also occur on other parts of the body. Height at shoulder from 5 feet 8 inches to at least 6 feet; girth, 105 inches; length, 12 feet 10 inches, of body 11 feet.

MAHARAJA OF KUCH BEHAR'S SPECIMENS

Height at shoulder	6 ft. $4\frac{1}{4}$ ins.	6 ft. 1 in.	6 ft. $\frac{1}{2}$ in.
Total length	14 ft. 1 in.	13 ft. 2 ins.	13 ft. 10 ins.
Body	11 ft. 11 ins.	11 ft. 2 ins.	11 ft. 8 ins.
Girth behind shoulder .	1 19 ins.	1 1 2 ins.	
Biggest girth	144 ins.	142 ins.	
Round neck nearest body	90 ins.	84 ins.	
" " head	74 ins.	75 ins.	
Horn	16 <u>‡</u> ins.	1 3 7 ins.	

At one time this animal was found over the greater part of the Indian peninsula, as attested by fossil remains, but now it is restricted to Assam; its place farther east in the Sanderbans, Chittagong, and Burma being taken by the smaller *R. sondaicus*.

Distribution.—Chiefly the Assam plain at the present day.

Length on front curve.	Circum- ference.	Weight.	Locality.		Owner.
-24	•••	•••	?		The late Dr. T. C. Jerdon.
19 1	228		India		British Museum.
-19	18	•••	Singpo, Burma .		Sir C. A. Elliot, Bart.
16 2			Belsire, Assam .		W. C. Sherwill.
♀ 16 1	•••	•••	Kuch Behar .		Maharaja of Kuch Behar.
9 16	•••	3½ lbs.	Nowgong, Assam		L. Fabre Tonnerre.
141	21	•••	Assam		Dr. W. P. Y. Bainbrigge.
9 14	22 7	41 lbs.	Nowgong, Assam		L. Fabre Tonnerre.
-13 7	•••	•••	Kuch Behar .		Maharaja of Kuch Behar.
¹ 13	•••	•••	Do		Do.
13	20½	•••	Assam		G. A. Dolby.
123	23	•••	Do	•	H. B. Firman.
121	215	•••	Do		H. C. Holland.
-12½	15½		Kuch Behar .		James J. Harrison.
I 2 ½	21	•••	?		J. W. Grieve.
-12	•••	•••	Bhutan Duars .	•	Major-General Alexander A. A. Kinloch.
11	20	•••	Kuch Behar .	٠	Capt. Hon. W. Lambton.
-103	212	•••	Foot of Garro Hills	•	A. O. Hume, C.B.
101/2	17	•••	,		Earl of Dunmore.
10	•••	•••	Kuch Behar .	•	Duke of Portland.
ç −8 <u>₹</u>	8	• • •	Do. .		Countess Scheibler.
8 <u>1</u>	15	•••	Assam		Sir Peter Walker, Bart.
8 1	147	•••	Do	•	Hugh G. Barclay.
8	17		Do	•	Major E. T. Paul.
6 1	15		Tezpore, Assam .		A. Y. Thomson.
5 1	141	•••	Kuch Behar .	•	Major Henry Streatfeild.
			¹ Height at shoulder, 6 ft.	6 i	ns.

JAVAN or LESSER ONE-HORNED RHINOCEROS (Rhinoceros sondaicus).

A less gigantic and smaller-headed species than the last, with the skin divided up into a kind of mosaic pattern, and the fold in front of the shoulder continued right across the body like the two hinder folds. The neck also lacks the large ring-like masses of folded skin. Horn never very large, and generally almost or completely wanting in the female. A female has been measured which stood $5\frac{1}{2}$ feet at the shoulder, and it is probable that the male stands little less than the great Indian species, although it is of lighter build.

Distribution.—The Sanderbans and other parts of Eastern Bengal, to the Terai, Sikim, Assam, and thence through Burma and the Malay Peninsula to Sumatra, Java, and Borneo.

Length on front curve.	Circumference.	Locality.					Owner.
108	19§	Java					British Museum.
10	191	Do.					H. Van Son.
81	20	Do.					A. S. Campbell,

SUMATRAN RHINOCEROS (Rhinoceros sumatrensis).

The smallest of the Asiatic rhinos, and the only one with two horns; differing, however, from the African members of the genus by the folds in the skin and the presence of teeth in the front of the jaws. Only the fold behind the shoulders is continued across the back, and the brown or black skin is rough, granular, and more or less hairy. Height at shoulder from about 4 feet to $4\frac{1}{2}$ feet; weight about 2000 lbs. Distribution.—From Assam (where the species is very rare) to Siam, the Malay Peninsula, Sumatra, and Borneo. Specimens from Chittagong are remarkable for the excessive development of the hair, which is long and very thick; they may indicate that the

Length on front curve.	Circumference.	Owner.	
321g	178	British Museum.	
27 1	177	Do.	

Assamese form is a distinct local race (R. sumatrensis lasiotis).



Common African Rhinoceros Skulls and Horns, from A. H. Neumann's specimens. The biggest horn of the three measures 40 inches on front curve.

COMMON AFRICAN RHINOCEROS (Rhinoceros bicornis).

Aurarisse of the Abyssinians. Chipamberi of the Lower Zambesi Upelepe of the Basutos. natives. Gurhu of the Danakil tribes. Favu of the Swahilis.

Sipejana of the Swazis and Matonga. Upeiana of the Matabeles and Zulus. Weel of the Somalis. Zwart Rhinoster of the Boers.

The African rhinoceroses are two-horned animals, readily distinguished from their Asiatic relatives by the absence of the folds of skin on the body which form such a characteristic feature of the latter; as they also are by the lack of front teeth in both the upper and lower jaws. The hide, too, is almost completely naked, although there are some bristly hairs on the margins of the ears and the tip of the tail. Average height at shoulder, 5 feet.

Perhaps the most distinctive external feature of the black rhinoceros, as the present species is commonly called, is to be found in the prehensile tip to the upper lip, which is rounded and not very wide in front. Other points of distinction are, however, shown by the form of the horns and ears and the position of the eyes; while in bodily size this animal is also considerably inferior to the next. If anatomical characters be taken into account, the black rhinoceros is also well characterised by the comparative shortness of its skull, and the form and structure of the molar teeth, which are adapted for a diet of twigs and leaves.

Captain Edgar G. Harrison, writing in the Field, December 25, 1897, records the measurements of a five-horned Rhinoceros shot by him in East Africa as follows:—



Head of Common African Rhinoceros.

First Horn.—Counting from snout, measurement $14\frac{3}{4}$ in.; is a normal front horn curving backwards.

Second Horn.—15 \(\frac{1}{4}\) in., curving forwards instead of backwards or being straight, as is usual, and leaning considerably over to the off-side.

Third Horn.—II $\frac{1}{4}$ in. long and $5\frac{1}{2}$ in. wide half-way between base and apex; is flat and perpendicular, and, though a distinct horn, grows out of the off-side of the base of No. 2.

Fourth Horn.—Is a stumpy, abortive horn, 9 in. long, growing partly from the base of No. 3, but quite separate and inclining slightly backwards.

Fifth Horn.—A distinct horn, $9\frac{3}{4}$ in. long, with its own base separated about 3 in. from No. 4.

Distribution.—From Abyssinia and Somaliland through East and Central Africa, in suitable localities, to the Cape. Now rare to the south of the Zambesi, and probably more abundant in the districts between the interior of Somaliland and Lake Rudolph than anywhere else. Although more alert and active than Burchell's rhinoceros, and thus a more dangerous animal, this species is by no means difficult to kill with modern weapons; and in the old days hunters frequently shot half-a-dozen in a single evening as they came to drink at a pool. In spite of its wide distribution, it has not been found possible to split up the species into local races; although this may perhaps be due to the want of sufficient specimens for comparison.


Lengt outside		Circum	erence.	Locality.	Owner.	
Front Horn.	Rear Horn.	Front Horn.	Rear Horn.	Locality.	Owner.	
44	•••			?	The late F. Holmwood.	
43		$21\frac{1}{2}$?	A. Beit.	
41½	10	201	16 <u>1</u>	Zululand	LieutCol. Hon. W. Coke.	
-411	•••	22½		Orange River	Major-Gen. Sir William Crossman, K.C.M.G.	
-4I			•••	?	Carl Hagenbeck.	
41	•••		•••	East Africa	The late F. Holmwood.	
-40	•••	22		Do	Berlin Museum.	
40	144	181	20]	Mt. Kenia, East Central Africa	A. H. Neumann.	
-39 1		211	•••	South Africa	J. Lamont.	
38%		21		?	Hon. Walter Rothschild.	
381	•••	19		Masailand	Sir John Kirk, K.C.B.	
-38	11	24	214	South Africa	Earl of Dartmouth.	
-36				East Africa	Sir Bartle Frere, Bart.	
-35 1		18		?	H. Murray.	
♀33 1	•••	171	•••	Matabeleland	W. Van Ness.	
-32½	21	19	15 1	Portuguese Northern	F. Vaughan Kirby.	
-32	112	197	191	Zambesia Athi Plains, East Africa	Count Scheibler.	
31	•••	18		East Africa	R. P. Carroll.	
31	191	16	16 1	Do	H. C. V. Hunter.	
301	•••	211		Zambesi Valley	Sir John Kirk, K.C.B.	

COMMON AFRICAN RHINOCEROS (Rhinoceros bicornis)—continued.

Leng outside		Circum	erence.				
Front Horn.	Rear Horn.	Front Horn.	Rear Horn.	Locality	7.	Owner.	
-29½	141	•••		Somaliland		A. H. Straker.	
29		19	•••	East Africa		Prince Boris Czetwertynski.	
28 <u>1</u>	•••	18	•••	South Africa		C. D. Rudd.	
28 <u>1</u>	88	18 <u>‡</u>		Do.		F. C. Selous.	
-27 ¹	12	•••	•••	Lake Njire		T. E. Buckley.	
27 1	10	21	17	Masailand .		Capt. R. A. J. Montgomerie,	
27	•••	20½	•••	East Africa		C.B., R.N. E. Gedge.	
27	16 1	178	171	Do.		Sir Robert Harvey, Bart.	
27	12	•••	•••	Do.		Sir John Willoughby, Bart.	
26 8	134	22	•••	?		Sir Edmund G. Loder, Bart.	
26 1	101	191	15	East Africa		Henry Charrington.	
♀ 25♣	17	16	15	East shore of Rudolph	of Lake	A. H. Neumann.	
25	9 1	178	•••	East Africa		F. J. Jackson, C.B.	
♀24 1	12	16	171	Masailand .		Capt. R. A. J. Montgomerie, C.B., R.N.	
♀ 24		101		Kilimanjaro		T. E. Buckley.	
22 <u>8</u>		19 1		East Africa		W. Astor Chanler.	
22½	141	17	16 1	Somaliland		Julius Jeppe.	
$22\frac{1}{4}$	10	22 3	19	Do.		Sir H. D. Tichborne, Bart.	
-22	12.3	20-8	19.2	East Africa		Prince Henry of Liechtenstein.	
212	118	19	19	Somaliland		J. Kenneth Foster.	
218	17	14	16	Abyssinia .		British Museum.	
-20½	91	214	198	Somaliland		Count E. Hoyos.	
20	$II_{\frac{1}{2}}$	21½	18	Do.		J. Byng Paget.	
20	68	201	191	'Do.		Major H. G. C. Swayne.	
-20	8 <u>1</u>	23½	21 7	Do.		Count E. Hoyos.	
-191	8		•••	Do.		Capt. M. M'Neill.	
-19‡	18 1	8 <u>1</u>	171	Sabi Flats .		Dr. Percy Rendall.	
-191	6 <u>1</u>	151	14	E. C. Africa		James J. Harrison.	
-19	22	18 1	20	B. E. Africa		Count Scheibler.	

COMMON AFRICAN RHINOCEROS (Rhinoceros bicornis)—continued.

Length on outside curve.		Circumfe	rence.				
Front Horn.	Rear Horn.	Front Horn.	Rear Horn.	Locality.	•	Owner.	
-19	8	14		Somaliland		Count Grudzinski.	
182	101	18 <u>1</u>	16 <u>1</u>	Do.		A. S. Trevor.	
-18 1	7	22	201	Do.		P. H. G. Powell-Cotton.	
-181		211	•••	Do.		Count T. Zamoyski.	
18	I 2 ½	20	22	East Africa		Lord Delamere.	
18	12	20	18	South Africa		Durban Museum.	
18	71	21	17	Somaliland		R. Wahrmann.	
171	72	19‡	198	Abyssinia .		British Museum.	
171	9 8	20	171	Somaliland		Prince Nicolas Ghika.	
-171	124	17	181	Do.		W. W. Ashley.	
17	6 <u>1</u>	21	17	Do.		Prince Demeter Ghika.	
17	13	16 1	181	Do.		J. Benett-Stanford.	
164	•••	15 1		Sudan .		Col. Ralph Vivian.	
-16 1	71	171	151	Somaliland		C. V. A. Peel.	
16]	8	20	18 1	Do		Count J. de Bylands.	
- 9 16	•••			Do		Capt. M. M'Neill.	
16	91	142	16	Do		Major V. M. Stockley.	
-152	6	18	15	Do		A. Leslie Renton.	
-154	8	20	19	Do		Count J. Potocki.	
-15	111	151	16 <u>1</u>	Athi Plains, E	ast Africa	Countess Scheibler.	
141	6 <u>1</u>	16		Somaliland		P. B. Vander-Byl.	
111	5 1	16 <u>1</u>	152	Do.		Col. Arthur Paget.	
10	6+3	(3 horns)		East Africa		Sir John Willoughby, Bart.	



Front Horns of Burchell's Rhinoceros.

From specimens in the collection of the late Roualeyn Gordon Cumming, in the possession of Col. W. Gordon Cumming.

BURCHELL'S RHINOCEROS (Rhinoceros simus).

Next to the Indian elephant this is the largest mammal that has trodden the earth in modern times; and its practical extermination cannot fail to be a matter of lasting regret. Its huge bulk, bluntly truncate muzzle, which has no prehensile tip, the great length of the skull, and the enormous front horn, with its expanded base, form the most striking external characteristics of this species. And on looking at the skull it will be found that the molar teeth are of quite a different type of structure from those of the preceding species; being, in fact, adapted for chewing grass. In walking, the animal carried its head low, so that in examples in which the front horn bends forward, its tip became worn by being pushed along the ground. It is difficult to surmise the reason for the application of the name "white rhinoceros" to this species, unless, indeed, it be that Cape specimens were lighter coloured than any seen in the districts to the north.

Distribution.—South and South-East Africa, in suitable localities, as far north as the Zambesi. Exterminated early in the century to the south of the Orange River; and now represented at the most by a few survivors in North-East Mashonaland, and possibly by others in the reedy swamps at the junction of the Black and White Umvolosy rivers. Between the Zambesi and Orange rivers the species was abundant less than half a century ago; Andersson alone having killed sixty in the course of a few months. Till a few years ago this rhinoceros was unrepresented by adult skins or skeletons in any British Museum; and it is to the credit of Messrs. Coryndon, Eyre and Varndell that complete examples were secured before it became too late.

North of the Zambesi there exists a rhinoceros (R. holmwoodi) at present known only by single horns, which appears to be a near ally of this species, although it was at first regarded as a variety of the common rhinoceros.

Length on outside curve.		Circum	ference.				_		
Front Horn.	Rear Horn.	Front Horn.	Rear Horn.	Locality	•		Owner.		
-62½		221	•••	South Africa		•	Col. W. Gordon Cumming.		
56 <u>1</u>	•••	23½ about		Do.			British Museum.		
-52½	•••	21 ½		Do.			Col. W. Gordon Cumming.		
44		20		?			British Museum.		
43 2		234		South Africa			A. Beit.		

BURCHELL'S RHINOCEROS (Rhinoceros simus)—continued.

	ength on circumference.			0		
Front Horn.	Rear Horn.	Front Horn.	Rear Horn.	Locality.	Owner.	
42 3	•••		25 8	Limpopo, S. Africa.	The late W. C. Oswell.	
-4I	•••			South Africa	J. W. Fitzherbert.	
40 3		29 8		Do	Sir Edmund G. Loder, Bart.	
-401		$22\frac{1}{2}$		Do	Do.	
40	•••	154	•••	Do	British Museum.	
-39				?	A. Ohlsson.	
♀ 38 §		241		?	The late W. C. Oswell.	
38 1		22 <u>‡</u>		?	J. B. Taylor.	
378	172	27 l		Mashonaland	F. C. Selous.	
37 1	·	24		South Africa	British Museum.	
-36 2		20½		. 3	Berlin Museum.	
36		28 <u>1</u>	•••	Mashonaland	C. D. Rudd.	
36		25 lbs.	•••	?	Rev. V. R. Carter.	
35½ (mounted specimen)	78	26	21	Mount Domo, Ma- shonaland	Cecil Rhodes, Cape Town Museum.	
-33½		23		?	Julius Jeppe.	
33		251		?	Duke of Westminster.	
33	13 1	23 1		Mashonaland	F. C. Selous.	
33	•••	25½	•••	?	Duke of Westminster.	
32 3		271		?	The late J. S. Jameson.	
♀ 32 <u>8</u>		27		Lake Ngami	The late W. C. Oswell.	
31 l		19½		?	Mr. Justice Hopley.	
31		24	•••	Zululand	LieutCol. Hon. W. Coke.	
-30 2		25 1		South Africa	Col. H. B. H. Blundell, C.B.	
♀ 29 8	5₺	23	201	Zululand	Julius Jeppe, Pretoria Museum.	
275	12	222	181	Do	Julius Jeppe.	
27		201	•••	Do	Dr. W. P. Y. Bainbrigge.	
27	113	30 8	25	Do	Sir Edmund G. Loder, Bart.	
23 1	•••	26		?	Mr. Justice Hopley.	
22½ (mounted specimen)	7 1	26 <u>‡</u>		Mashonaland	Hon. Walter Rothschild.	
	13	•••	191	?	Mr. Justice Hopley.	
20 2 (mounted specimen)	7	28 <u>3</u>		Mashonaland	British Museum.	
20	6	251/2	181	Zululand	H.R.H. le Duc d'Orléans.	



From a Photograph by Mr. Eaks, Reading.
Skull and Horns of Burchell's Rhinoceros, Mashonaland, 1880.
Shot by F. C. Selous.



INDIAN ELEPHANT (Elephas indicus).

In general a decidedly smaller animal, the Indian elephant differs from its African relative not only in external form, but also in the structure of its molar teeth, which are composed of a greater number of much thinner vertical plates. The females, as a rule, have only very small tusks, not projecting beyond the jaw; and in some cases those of the males are equally poorly developed. Five nails are usually present on the fore-feet, and four on the hinder ones. Externally, the most characteristic distinction is the comparatively small size of the ears; next to which comes the presence of a finger-like process on the front edge only of the tip of the trunk; the African species having one in front and a second behind. The skin is nearly smooth; and the

bristles on the tail are confined to the front and back edges for some distance above the tip. Other noticeable points are the comparative flatness of the forehead, and the regular convex form of the back. Although males do not generally exceed 9, and females 8 feet in height, specimens have been killed measuring 9 feet 10 inches, 10 feet 1 inch, and 10 feet $7\frac{1}{2}$ inches, while one is stated to have reached 11 feet, and there is evidence of still larger individuals, perhaps of 12 feet.

Distribution.—The forest districts of India, Ceylon, Assam, Burma, Siam, Cochin China, Sumatra, and Borneo. The Sumatran elephant has been regarded as a distinct species, and it may possibly form a peculiar local race.

Heigh Shoul	lder.			Locali	ity.		Owner.
ft. –11	in. O	Mysore					Viscount Powerscourt.
-10	4	Gurhwal,	N.W	7.P.			Col. J. E. Campbell.
-10	1	Ceylon					James J. Harrison.
-10	1			?			Major-General Alexander A. A. Kinloch.
-9	7	Mysore				•	The late Col. G. P. Sanderson.
-9	4	Ceylon					Count Scheibler.

Tusks.

side curve.	Greatest circumference.	Weight.	Locality.				Owner.			
ft. ins. -8 9 R	ins. 17 ‡	lbs. 81	Assam	•			Presented to the late Lord Lytton by Sir Steuart Bayley, K.C.S.I.			
-8 2 L		80-2	Do.				Measured by A. O. Hume, C.B. Do.			
		100	Do.				The late Charles Redde, P.Z.S. 1886, p. 176.			
8 o	16 7	90	South In	ndia			Sir Victor Brooke's Collection.			
¹ 7 3 ³	171	102	Burma	•		Į	Marquis of Waterford.			
7 31	178	971	Do.			ſ	marquis or wateriord.			
6 10	171	65∰	India				J. D. Goldingham, Bethnal Green Museum.			
-6 8		77 1	Madura South				Col. G. M. Payne.			
6 7½	148	521	Mysore				Viscount Powerscourt.			
-6 7	121	46 <u>1</u>	Yala, S Ceylo		ovinc	es,	Capt. R. J. Marker.			
-6 3½	•••	73 1	Madura		rict		Col. G. M. Payne.			

¹ The tusks of the sacred white elephant which came out of King Thebaw's Palace, Burma.

INDIAN ELEPHANT (Elephas indicus). Tusks-continued.

Length out- side curve.	Greatest cir- cumference.	Weight.	Locality	·.	Owner.
ft. ins.	ins.	lbs.			
6 o		50	Madras .		J. Fortune.
-5 10			Borneo .		C. M. D. Stewart.
-5 3	15	42	Ceylon .		W. S. Murray.
-5 I⅓	15	35	Travancore		Hon. E. Stonor.
-5 o	•••	•••	Burma .		T. Mumford.
4 10	151	32½	Travancore		Hon. E. Stonor.
-4 7	13		Do.		J. D. Rees.
4 6	15		Do.		Lord Wenlock, G.C.I.E.
4 41	15		Do.		Do
13 112	121	29 ³	Yala, S. Pro Ceylon	ovinces,	Capt. R. J. Marker.
		MEASUREM	ENTS PROTRU	DING FR	om Skull.
3 11			India .		British Museum.
3 6 7			Do		Do.
3 3	•••		Do		Do.
2 5∰	15	•••	Do		Do.
2 01			Mysore .		Viscount Powerscourt.
1 6	48		Sumatra .		British Museum.

¹ Broken end, pair to 6 ft. 7 ins.


Feet.

Circumference at base.	Width at bottom, back to front.	Locality	y.		Owner.			
-67 1	•••	Measureme			from living elephant under charge of			
62 <u>1</u>		South Arcot I						
61	20	Travancore			D. M. Lumsden.			
60	18	Mysore .			Viscount Powerscourt.			
-60	•••	Gurhwal .			B. B. Osmaston.			
-60½	•••	India			Major-Gen. Alexander A. A. Kinloch.			
-591	•••	Gurhwal .		•	B. B. Osmaston.			
58 <u>1</u>	18	Travancore			H. L. Cottingham.			

INDIAN ELEPHANT (Elephas indicus). Feet—continued.

Circumference at base.	Width at bottom, back to front.	Locali	ty.	•	Owner.
58	18	Travancore			Hon. E. Stonor.
574	19	Assam .			H. C. Holland.
1-571	17	Ceylon .			R. Gordon Smith.
571	18 1	Assam .	•		D. D. F. Hosack.
-56	•••	Ceylon .	•		E. J. Brooke.
56	17	Do			G. M. Norrie.
-56	18	Do			Sir Peter Walker, Bart.
-56	171	Do		•	T. H. Mann.
55 1	18 <u>1</u>	Travancore			Lord Wenlock, G.C.I.E.
55	178	Ceylon .		•	Rowland Ward.
² 55	17	Gurhwal .	•	•	B. B. Osmaston.
55	17	N.W. Provin	ces .	•	Capt. G. O. Bigge.
54 2	18	Pyinmana Ta	i.	•	H. Partridge.
54½	171	Ceylon .	•	•	H. Storey.
53 8	18 <u>1</u>	Do	•	•	W. H. Walker.
521	17	Do	•		H. E. Lindsay.
-50	16 1	Do	•	•	James J. Harrison.
49	15	Do	•	•	Capt. R. J. Marker.
48 <u>1</u>	16	Sumatra .	•	•	R. P. Carroll.
46 <u>1</u>	15	?			G. A. Dolby.
45 1	15 1	Mysore .		•	W. G. Deeds.

¹ Measurements 54 ins. × 17 ins. when dried. 2 Measurements, 59\frac{1}{2} when shot.



Head of African Elephant.

AFRICAN ELEPHANT (Elephas africanus).

Njovu in the Chilala and Chibisa countries.

Arba of the Gallas.

Dakana of the Abyssinians (Danakil).

Marodi of the Somalis.

Tho in the Barotse country.

Thio in the Lake Ngami country.

Zahon of the Abyssinians.

Zemba of the Swahilis.

The African elephant, although still abundant in many regions of Central Africa, in the southern part of the Continent is rapidly approaching extinction. A very remarkable exception to this melancholy process of extermination is, however, to be found in the south and east of Cape Colony, where, since the year 1830, wild elephants have been systematically preserved by Government. Strong troops of these protected elephants still roam the dense and impenetrable jungles of the Addo Bush and the Knysna and Zitzikamma forests. A permit to shoot one of these elephants—costing £20—is to be obtained at Cape Town. Farther inland the ivory-hunters have for a generation past been so actively employed, that, despite the vast numbers of these

great mammals which forty or fifty years ago thronged the interior from the Orange River northwards, but a few troops are now left south of the Zambesi. In all Khama's country of Bamangwato, for instance, where Gordon Cumming, Oswell, and others shot most of their elephants, only one solitary troop remains!

ft.	in				Owner.
10	7	" Jumbo"	•		Barnum and Bailey.

Male shot by James J. Harrison in the Chiperoni Forest, Mozambique Province.

			ft.	in.	1	ft.	in.
					Girth of forearm	4	6
Crown to tail-tip .		•	17	3	,, fore-foot	4	11
Vertical standing height	•	•	II	I	The tusks were not very large.		

Estimated height of a huge tuskless male shot by Mr. F. Vaughan Kirby in the Chiringoma Forest, 10 feet 8 or 9 inches, circumference of forefoot 60 inches.

Male shot in Portuguese East Africa by F. Vaughan Kirby.

	ſt.	in.	t	ft.	in.
Length from trunk-tip to crown	9	7	Greatest height above ground as it		
crown to tail-tip.			lay dead		4
Vertical shoulder height	10	6 <u>1</u>	Right tusk on the curve		10
Perpendicular diameter of left	_	_	Left ,, ,,	5	111
over the flap Girth of left fore-foot	,	9 10	Greatest circumference		19

Mr. A. H. Neumann, in his *Elephant Hunting in East Equatorial* Africa, records the following among other dimensions:—

Male.	Height at shoulder in straight line.	Length from root of tail to eye, in straight line.	Girth of forearm below elbow.	Circum- ference of fore-foot.	Long diameter of hind-foot.	Girth of thickest tusk just outside lip.
Shot at El Bogoi (foot of Lorogi Mountains), 30th Sept. 1894.	ft. in. 10 8 or 9 (at least)	ft. in. 12 6	ft. in. 4 6	ft. in. 4 8½	ft. in. I 9	ft. in. I 6½
Shot at Bumi (Lake Rudolph), 29th Dec. 1895.	IO 5 (at least)	12 6		4 8½		1 б
Do. do.	IO 9 (at least)	12 8 or 9	•••	5 O	1 10	1 9 1
Shot at Janjai (near Kenia), 13th May 1894.	10 6 (at least)				•••	•••

The three first were old bulls of the type found, as a rule, in separate herds. The last was a "herd bull" or breeding male.

APRICAN ELEPHANT (Elephas africanus)—continued.

	lder zight	Total leagth (trunk to end of tail).	Girth of belly.	Girth of fore-foot.	Locality.		Owner.
ft.	in.	ft. in.	ft. in.	ft. in.			
-10	6	•••	•••	•••	S.E. Africa .	•	F. C. Selous.
-10	31				E. C. Africa .	•	James J. Harrison.
-10	3	21 0	18 0	5 1	Nyasaland .		Sir John Kirk, K.C.B.
-10	0	23 5	16 6		South Africa .	•	H.R.H. the Duke of Saxe-Coburg and Gotha.
-9	7ŧ		•••	4 6 1 (dried)	Somaliland .	•	Viscount Edmond de Poncins.
-9	61			•	Do	•	Capt. M. M'Neill.
-9	3	24 6			?		Count E. Hoyos.
-9	2				Ruwenzori, E. C.	Africa	Capt. C. Ashburnham.
-8	o			3 81	?		Count Scheibler.

Tusks.

Length (outside curve).		Greatest circumference.	Weight.	1	ocali	ty.	Owner.
ft. 9	in. 5	in. 22	lbs. 184	Africa		•	Sir Edmund G. Loder, Bart.
•••		•	172	Do.			The late Sir Samuel Baker.



Specimens of A. H. Neumann's large Elephant Tusks.

9	4	201	160	E. Africa	Sir John Kirk, K.C.B.
, 9 ,	.4	18	110	Do	Duke of Westminster.
(tip bi −9	roken) 4		151	Elgayu, E. Africa	F. J. Jackson, C.B.
-9	3		129	?	Do.
. 9	0	181	116	N. of Lake Rudolph, E. C. Africa	A. H. Neumann.
9	0	21	150	Africa	Sir Edmund G. Loder, Bart.
9	0	181	116	N. of Lake Rudolph	A. H. Neumann.

AFRICAN ELEPHANT (Elephas africanus). Tusks-continued.

Let (outside	ngth curve).	Greatest circumference.	Weight. Locality.			Owner.
ft.	in.	in.	lbs.			
8	11	18 §	•••	Galla Country.	٠	Measured by Viscount Edmond de Poncins.
8	II	171	90	3		H.M. the Queen, Bethnal Green Museum.
8	10	18	105	East Africa .	•	Lord Delamere.
-8	8	18 1		?		Sir H. B. Meux, Bart.



From a photograph by Mr. J. R. W. Pigott.

The Big Tusk (8 st. 7½ in., 165 lbs.) presented by the Officers serving in British East Africa to H.R.H. the Duke of York on the Occasion of his Marriage.

8	71	223	165	East Africa		H.R.H. the Duke of York.
8	6	18 1	100	Do.		Lord Delamere.
8	3	183	811/2	Do.		T. W. Greenfield.
8	2	18 <u>3</u>	80	Do.		Do.
8	1	20 1	114	N. of Lake R E. C. Afri	ph,	A. H. Neumann.

AFRICAN ELEPHANT (Elephas africanus). Tusks—continued.

	gth curve).	Greatest circumference.	Weight.	Locality.	Owner.
ft. 7	in. IO	in. 19 §	lbs. 107	East Africa	A. H. Neumann.
7	81	231	160	Africa	Chalmers, Guthrie, and Co.
-7	8		108	Galla Country	Dr. Donaldson Smith.
7	6	•••	100	Do	Do.
7	6	17		?	Duke of Westminster.
7	6 <u>1</u>	211	•••	South Africa	The late W. C. Oswell.
7	3 1	191	102	East Africa	Col. F. D. Lugard, C.B., D.S.O.
7	O.	171	60	Kilimanjaro, East	F. J. Jackson, C.B.
7	o	18	(about)	Africa South Africa	Cetywayo's peace-offering in 1879
6	9	(about) 17 1		Kilimanjaro	to Lord Chelmsford. F. J. Jackson, C.B.
6	8	17	74	?	Julius Jeppe.
6	7	16 1		South-East Africa .	F. C. Selous.
₽6	51	121		N. of Lake Rudolph,	A. H. Neumann.
6	41	171		E. C. Africa East Central Africa.	H. S. H. Cavendish.
6	I Laurele)	71/2		Masailand	Sir John Kirk, K.C.B.
-6	ıl tusk) I	18		Somaliland	Sir H. B. Meux, Bart.
5	10	16 1		White Nile	The late Sir Samuel Baker, British Museum.
-5	9	18	70	Nyasaland	S. Pulley.
5	4	141		East Central Africa.	G. E. Smith.
5	4	16 1	78	Ruwenzori, E. C. Africa	Capt. C. Ashburnham.
4	11	15	26	Somaliland	A. H. Straker.
4	11	16		Ruwenzori, E. C. Africa	Capt. C. Ashburnham.
4	91	148	331	Somaliland	E. W. S. Brooke.
4	81	131	•••	Do	Sir Edmund G. Loder, Bart.
4	5	121/2	•••	Do	Lord Delamere.
4	5	101	20	Do	P. R. Denny.
4	5	10		East Africa	W. Astor Chanler.
4	4	13 3	26	Somaliland	Capt. J. M'Call Maxwell.
4	I	141	60	Do	Digby Davies.

AFRICAN ELEPHANT (Elephas africanus). Tusks—continued.

	ngth curve).	Greatest circumference.	Weight.	Locality	γ.		Owner.
ft. -4	in. I	in. 13½	lbs. 15 2	Somaliland			Capt. M. M'Neill.
4	8 <u>1</u>	11	194	Do.			P. R. Denny.
-3	81	145	16 1	Danakil .			Viscount Edmond de Poncins.
-3	41	145	16 }	Do			Do.
-2	10‡	72	$6\frac{5}{16}$	Do		•	Do.
-2	10	7 3		Somaliland		•	Dr. Donaldson Smith.
-2	10			Do.			James J. Harrison.
		Me.	ASUREME	NTS PROTRUD	ING	FROM	A SKULL.
-4	8			Somaliland			Capt. M. S. Wellby.
-4	1	16 1		5			Julius Jeppe.
-3	11	16		?			Do.
3	5 7	15 1		South Africa			The late J. S. Jameson.

Feet.

Circumference at base.	Width at bottom, back to front.	Locality.		Owner.				
-6o 	•••	Somaliland			Capt. M. S. Wellby.			
58 1	19	E. C. Africa			H. S. H. Cavendish.			
56	19	Somaliland		•	The late Col. E. Carrington.			
54	18 1	Do.		•	Major J. S. Ashby.			
54	•••	Kilimanjaro			F. J. Jackson, C.B.			
-54	17	Somaliland			Count E. Hoyos.			
53 8	18 1	South-East Afri	ica		The late J. S. Jameson.			
53	17	Somaliland			Capt. J. M'Call Maxwell.			
52 <u>1</u>	19	Do.		•	Major J. S. Ashby.			
52½	17	South Africa		•	Rowland Ward.			
-51	•••	Galla Country		•	Viscount Edmond de Poncins.			
50	17	South Africa			F. C. Selous.			
49 1	171	Do.			Do.			

AFRICAN ELEPHANT (Elephas africanus). Feet-continued.

Circumference at base.	Width at bottom, back to front.	Locality.	Owner.
-49½	•••	East Central Africa	James J. Harrison.
491	17	Rhodesia	W. W. Ashley.
491	17	Masailand .	Col. F. D. Lugard, C.B., D.S.O.
49 1	16 <u>1</u>	East Africa .	W. Astor Chanler.
48 7	16 1	East Africa .	H. C. V. Hunter.
-48 3	16 1	Somaliland .	Capt. M. M'Neill.
48	•••	Uganda	Col. F. D. Lugard, C.B., D.S.O.
48	16	B. E. Africa .	Col. Trevor Ternan.
44 2	144	Sudan	Col. Ralph Vivian.
44 1	151	Somaliland .	Prince Demeter Ghika.

MAMMOTH or SIBERIAN ELEPHANT (Elephas primigenius).

Nearly allied to the Indian elephant, but the plates of the teeth still narrower and more numerous, the tusks spirally twisted, and the skin clothed in woolly fur with long bristles intermixed.

Remains of this species occur in the superficial deposits of Europe and Northern Asia. In the frozen soil of Siberia the skin, flesh, and hair are frequently preserved, and the ivory is often suitable to the purposes of the turner.

Length (out- side curve).	Greatest Circum- ference.	Weight.	L	ocality.		Owner.		
11 ft.	20% ins.	173 lbs.	Siberia			Sir Edmund G. Loder, Bart.		

LION 453

LION (Felis leo).

Ambassa of the Abyssinians.

Asced of the Arabs.

Imbubi of the Swazis and Zulus.

Lendja of the Gallas.

Libbah of the Somalis.

Libbaka of the Abyssinians (Danaki).

Mkango or Nkalamo in the Chilala and Chibisa countries.
Simba of the Swahilis.
Tau of the Basutos.
Tauw in the Barotse country.
Tauw in the Lake Ngami countries.

Any description of such a familiar animal as the lion (the only cat in which the male is furnished with a mane on the head and shoulders, and a tuft of long hair to the tip of the tail) would obviously be superfluous here. One of the great points of interest attaching to the species is its wide geographical distribution; and it has not yet been determined that the Indian form can be separated as a distinct race, the alleged absence of the mane having been shown to be inconstant. Neither can black-maned and yellow-maned lions be regarded as distinct forms, since examples of both may be met with in one and the same litter. Somali lions seem, however, to run smaller than those from either the Cape or Algeria, although their manes are Heights of 3 feet 41 and 3 feet 8 inches at the often very fine. shoulder have been recorded in African specimens (the larger measurement by F. C. Selous), and 3 feet 6 inches in an Indian example (by Gen. W. Rice). Colonel Paget estimated the weight of a Somali lion at about 550 lbs. Wild lions never develop the enormous manes frequently seen in menagerie examples.

Distribution.—At the present day Africa from Algeria to the Cape, Mesopotamia on the west flanks of the Zagros range, Persia south of Shiraz, and India in the districts of Kathiawar, Sind, the Central Provinces, and Bundelcund. Now very rare in the latter country.

LION (Felis leo).

Owner.	F. Vaughan Kirby.	Do.	Count T. Zamoyski.	W. Yellowby.	F. C. Selous.	Capt. R. A. J. Montgomerie,	Sir John Willoughby, Bart.	T. E. Buckley.	Major A. St. H. Gibbons.	J. Johnston Stewart.	A. E. Leatham.	R. G. Beswick.	Prince Henry of Liechten-	stein. James J. Harrison.	Norman B. Smith.	Kathiawar, India Lord Harris, C.C.S.I.
Locality.	S. E. Africa	۸.	Somaliland .	Edmund's	Hartley Hills,	British East Africa	Mashonaland .	Limpopo	Mushukulumbwe	Somaliland .	Do	Zomba, B. C.	East Africa .	Oliphant River .	North Somaliland	Kathiawar, India
Description.	Full black mane	÷	i	Fine mane .	Do.	Do:	Fair mane .	Do	Full black mane	Yellow and black	Do.	Full yellow mane	i	Yellow mane .	Fine mane	Do.
Weight.	. \$:	:	434	410	916	:	:	:	:	:	:	:	:	:	:
Girth behind Weight.	. . :	÷	:	22	:	\$1 \$:	፥	49 \$	፥	÷	፥	:	:	:	:
Girth of forearm	. s i :	70	÷	22	:	₹61	:	:	:	:	:	15‡	:	:	:	÷
Height at shoulder.	ft. in. 3 7	:	፧	3 6	3	3 6	:	:	3 7	÷	:	3 44	:	:	:	:
Skin measure- ment from tip of nose to tip of tail.	ft. in. 10 10s	(14m) 11 4	:	:	6 11	(raw) 10 8 (descend)	(nacearn)	:	12 1	10 10	9 11	:	:	:	11 3	:
Length from nose to root of tail.	بو. : ii	7 2	:	01 9	:	8 9	÷	:	:	7 o <u>4</u>	:	§ 9 9	01 9	÷	:	÷
Total length fore skinning.	ft. in. -10 5 (A)	-10 4	4 01-	-10 0	-9 11 (c)	(a) for 6-	01 6-	-9 10 (E)	(н) от 6-	01 6-	-9 IO(K)	8 6-	-9 8.13	8 6-	7 6-	(1) 4 6-

Prince Henry of Liechten-	Major A. E. Sandbach.	LieutCol. L. L. Fenton.	James J. Harrison.	A. E. Pease.	Count J. Potocki.	Count Scheibler.	C. V. A. Peel.	Major T. R. Harkness.	Capt. M. M'Neill.	Lord Delamere.	F. C. Selous.	Count Scheibler.	Capt. G. Campbell.	Lord Delamere.	J. Johnston Stewart.	Capt. Smee.	Count E. Hoyos.	F. Vaughan Kirby.	Capt. R. A. J. Montgomerie, C.B., R.N.
East Africa	Somaliland .	Kathiawar .	Oliphant River .	Somaliland .	Do.	Do.	Do.	Do.	Do.	Do.	Pungwe, S. E.	Somaliland .	Do.	Do	Do.	Central India .	Somaliland .	~	British East Africa
:	Fine lioness	Do.	Do.	:	÷	Wellmaned, par-	tially black	Black mane .	:	Lioness .	Fair mane .	Light coloured	mane Fine black mane	Good mane .	Lioness	:	:	Lioness	Old lioness .
:	;	:	:	:	:	÷	፥	:	:	:	:	:	:	:	:	490	:	:	:
:	÷	÷	:	:	:	45	:	84	;	:	484	4	:	÷	:	:	÷	:	÷
i	:	÷	:	:	:	91	÷	:	:	÷	17	15	:	:	:	:	:	:	:
;	÷	÷	÷	3 11	÷	3 11	:	:	;	:	3 4	3 64	÷	i	:	:	÷	3 3	:
į	:	:	:	11 01	9 6	10 3	0 11	0.	(dressed)	:	0 11	(raw) 9 74	4 01	(dressed)	(dressed)	:	9 54	6	(war) 9 7
1 8 9	÷	9 9	:	:	:	9 9	:	:		i	;	1 9	:	į	7 0 9	÷	÷	÷	÷
7	9 6	9 5	9 44	4	4	-9 3	-9 3	9 2 (F)	6	6 2	1 6	-6 1	(c) 1 (c)	9 1	-8 10	₹6 8-	<u>:</u>	-8 74	9 8-

In the accompanying list of skull measurements many will be found belonging to the skins on p. 454.

I have not yet seen a Somali skull measuring over 15 inches. A lion skull may be easily recognised when placed beside that of a tiger by observing that in the lion all the terminations of the sutures of the skull on the frontal are almost level; in a tiger the nasal bones extend much further back—besides this a lion skull will stand much flatter on a table than that of a tiger.

Skulls.

Basil length from back to front.	Width across the zygomatic arches.	Weight cleaned.	Locality.	Owner.
16½ (A)	10		South-East Africa	F. Vaughan Kirby.
16 (end broken)	101	5 lbs.	East Africa	E. Gedge.
-15 ⁷ / ₈	94	6 lbs.	South Africa .	J. Lamont.
15½ (D)	101	5 lbs. 3 oz.	East Africa	Capt. R. A. J. Montgomerie, C.B., R.N.
151	91		Pungwe	Hon. T. Thynne.
-15 ≟ (H)	9 3	•••	Mushukulumbwe .	Major A. St. H. Gibbons.
15	10		Mashonaland .	Basil H. Woodd.
-15	10	•••	Somaliland	J. D. Inverarity.
15	10	•••	Benguela	G. W. Penrice.
15	91/2	•••	Beira	H. T. and A. H. Glynn.
15	91	•••	Athi Plains	C. F. S. Vandeleur, D.S.O.
15	10	•••	Somaliland	Col. Arthur Paget.
15 (C)	10	5½ lbs.	South-East Africa	F. C. Selous.
142	101	•••	East Africa	Lord Delamere.
144	9		Matabeleland .	Capt. Sir K. Fraser, Bart.
141	93	4 lbs.	South Africa .	Sir Edmund G. Loder, Bart.
-14 8 (E)	9 1	•••	Limpopo	T. E. Buckley.
1416	9 1	••	Matabeleland .	Capt. Sir K. Fraser, Bart.
14½ (G)	94	•••	Somaliland	Capt. G. Campbell.
-141	81	3½ lbs.	Nyasaland	Dr. Percy Rendall.
141	9 8	•••	Somaliland	LieutCol. J. W. H. Flanagan.
141	98	4½ lbs.	Do.	Rowland Ward.


LION (Felis leo). Skulls-continued.

Basil length from back to front.	Width across the zygomatic arches.	Weight cleaned.	Locality.	Owner.
−14 <u>‡</u> (K)	91	•••	Somaliland	A. E. Leatham.
148	9		Matabeleland .	Col. R. S. S. Baden-Powell.
♀ 14 <mark>1</mark>	82	•••	South-East Africa	F. Vaughan Kirby.
14	10		Somaliland	Capt. C. J. Melliss.
-14	9₿	•••	Do	H. W. Seton-Karr.
14	91	•••	Do	Lord Delamere.
-14 (1)	91/2	4 lbs.	British East Africa	Count Scheibler.
134	91	•••	Somaliland	Lord Wolverton.
132	91	•••	Do	E. W. S. Brooke.
-134	91	•••	Do	C. V. A. Peel.
-13½	8	•••	South Africa .	Dr. W. P. Y. Bainbrigge.
138	8 §	•••	Zomba	A. H. Sharp.
131	81	•••	Mashonaland .	Sir John Willoughby, Bart.
131	9		Kathiawar, India.	Lord Harris, G.C.S.I.

TIGER (Felis tigris).

A much less noisy animal than the lion, the great striped cat of Asia is also a more variable species, of which at least three local races may be recognised. First, we have the typical Bengal tiger, a large, long-limbed, lithe, and short-haired creature. A much smaller and rougher-haired race (F. tigris virgata) inhabits the Caspian provinces of Persia, and it is probably this form that extends into the Caucasus. Lastly, we have the Manchurian tiger (F. tigris longipilis), characterised by its large size, heavy build, short limbs, and the great length and thickness of the fur, which is often much less fully striped than in Indian specimens. The dimensions attained by tigers are given in the subjoined table.

Distribution.—From the Caucasus through Northern Persia, Afghanistan, India, Assam, Burma, the Malay Peninsula, Sumatra, Java, and China, to Manchuria and Amurland. In India ranging from an elevation of some 7000 feet in the Himalaya to Cape Comorin, but unknown in Ceylon.



before sximing 10 th 12 mm dried ekin 11-7-

	Owner.	Col. Evans Gordon.	H.I.M. the Sultan of Turkey.	(Shot by late paron de Nolde.) Noel Fenwick.	A. M. Markham.	W. Holland.	Maharaja of Kuch Behar.	A. M. Markham.	Maharaja of Kuch Behar.	Sir E. L. Durand, Bart, C.B.	Otho Shaw.	H.H. the Maharaja of Kuch Behar.	Do.	Major J. W. M. Cotton.	Capt. Hugh Fraser.	James J. Harrison.	H. de Barreto.	E. R. Henry.	Maharaja of Kuch Behar.	R. Nolan.	Capt. S. II. Pollen.
inned.	Locality.	Ramshai Hâb,	Duars, Dengal Assam Frontier .	Central Provinces	Bignor, N.W.P.	Kanara, B.P.	Kuch Behar .	Balti Dun, N.W.P.	Kuch Behar .	Nepal Terai	Do	Kuch Behar	О.	:	Kuch Behar .	Central Provinces	Rewa, C.P.	~	Kuch Behar .	Terai Forests,	Duars
)—cont	Weight.	lbs. 491	÷	:	:	:	487	:	530	:	•	462	8	÷	÷	÷	÷	:	540	:	÷
TIGER (Felis tigris)—continued.	Height at shoulder.	A. in.	:	i	3 6	:	3 4½	3 4	3 34	:	:	3 44	3 8	:	÷	:	÷	÷	3 4	:	÷
ER (F	Head.	36 ft	:	:	39	:	364	321	383	:	:	38	40 }	:	;	:	:	÷	36	:	÷
TIG	Girth. Fore	٦. 19	:	÷	21	:	\$ 61	₹61	8	:	:	13‡	21	÷	÷	:	:	:	21	÷	:
	Upper arm.	2 5 %	:	:	:	:	29	:	56 }	÷	:	283	29	:	:	:	:	:	5 0	÷	:
	Girth of body.	inches. 56½	÷	÷	25	:	52	49	484	:	:	Şış	7.	:	:	÷	8	:	22	:	;
	Length dressed.	ft. in. 12 1	8 11	:	12 2 1	:	12 I	12 2	÷	:	÷	12 4	11 7	10 44	01 11	፥	12 6	{ 11 11 }	:	12 2	:
	Length of body.	7. in.	:	:	7 14	:	7 14	1 1	0 2	:	;	01 9	7 14	7 3	፧	:	:	:	11 9	:	:
	Length before skinned.	ft. in. -10 7(A)	9 01-	-10 58	-10 4	-10 4 (D)	-10 3	-10 3(c)	-10 21	-10 21	-10 2	-10 14	£1 01-	1 01-	1 01-	1 01-	-10 O	-10 04	01-	·10 (R)	10 4(F)

Sir Edmund G. Loder, Bart.	Count E. Hoyos.	Do.	A. E. Leatham.	Maharaja of Kuch Behar.	Capt. S. H. Pollen.	Capt. Hon. R. H. Marsham and	Major B. B. Russell.	A. M. Rotheram.	James J. Harrison.	Capt. C. E. G. Norton and Capt.	Count E. Hoyos.	LieutCol. B. D. Möller.	Capt. J. H. Gwynne.	Lieut-Col. L. L. Fenton.	Lieut-Col. F. H. Whitby.	O. V. Bosanquet.	The late Sir Samuel Baker.	Count E. Hoyos.	Hyderabad, Deccan Major G. A. L. Carew, D.S.O., and	Capt. Frank Lee.	J. D. Inverarity.	Count Scheibler.	
Nilgiri Hills	Kuch Behar .	Rewa, C.P.	Nepal	Kuch Behar .	Do	Hyderabad .	Nepal Terai	Hyderahad, Deccan A. M. Rotheram.	Central Provinces	Hyderabad, Deccan	Kuch Behar .	Terai	Philipil	North Kanara .	~	Rewa, C.P.	Do	Do	Hyderabad, Deccan	Deccan	Do	Kuch Behar .	-
÷	÷	:	;	508	20	:	:	:	:	:	:	:	:	:	:	437	:	:	:	:	:	420	
:	3 3½	:	÷	3 4	:	3 9	÷	:	:	3 34	÷	:	:	:	:	:	:	÷	:	:	÷	3 24	
:	37	÷	÷	39 ‡	÷	:	:	:	:	34%	÷	÷	:	:	:	:	:	:	:	:	:	324	
÷	707	:	÷	194	÷	:	:	:	:	17	:	:	:	:	:	:	:	:	:	:	:	81	
:	284	፧	:	29	:	:	:	÷	:	24§	:	:	:	:	:	:	:	:	:	:	:	92	
:	₹ 0 5	2 2	:	72	:	:	፧	:	:	8	:	45	:	:	:	:	:	:	÷	:	÷	49	
÷	:	2 11	11 1 4	:	12 10	:	:	8 11	:	:	:	:	:	:	÷	:	4 11	:	:	6 01	÷	10 4 <u>4</u>	
:	6 9	9 9	:	11 9	6 10 4	:	9	:	:	:	8	÷	:	:	:	:	:	6 3	:	:	:	4	
-10	11 6	11 6	11 6	4 01 €	f or 6−	-9 10	9 10	01 6-	6 68-	€8 6−	8 6-	8 6	8 6	-9 8 (B)	8	-9 7	7 6	9 6-	9 6	9 6-	9 5	6 5	

	Owner.	C. C. Branch.	Capt. Hugh Fraser.	Capt. H. B. Dalgety.	Capt. H. P. Lane.	Н. de Barreto.	Maharaja of Kuch Behar.	O. V. Bosanquet.	Capt. W. H. Hunter.	Bombay Natural History Society's	Capt. M. McNeill.	A. E. Leatham.	Capt. B. H. Boucher.	LieutCol. L. L. Fenton.	Capt. Frank Lee.	Major G. A. L. Carew, D.S.O, and	Capt. C. L. 110105Worth. Viscount Edmond de Poncins.	C. C. Branch.	O. V. Bosanquet.	Surgeon-Capt. E. McK. Williams.
nued.	Locality.	Rewa, C. P.	Kuch Behar	Mount Abu.	Near Rewa, C.P.	Ď.	Kuch Behar .	Eastern Bengal .	Central Provinces	Do.	Chanda, C.P.	Nepal	Ram Naydr,	North Kanara	Deccan	Hyderabad, Deccan	Gwalior	Rewa, C.P	Do	Bignor District
)—conti	Weight.	喜 :	418	:	፥	:	360	:	over 563	448	:	:	:	:	:	:	:	:	:	345
TIGER (Felis tigris)—continued.	Height at shoulder.	.i :	÷	;	÷	:	2 10	÷	:	÷	:	:	:	:	:	:	:	:	:	÷
ER (Fe	Head.	æ :	:	:	፥	:	36	:	39	:	፥	÷	:	:	:	:	:	:	:	÷
TIG	Girth. Fore	∉ :	:	:	:	:	91	:	201	:	:	፧	:	:	:	÷	:	÷	÷	:
	Upper arm.	: ئە	:	:	:	:	22	:	÷	:	፧	÷	:	:	:	:	፥	÷	:	÷
	Girth of body.	inches.	:	:	4	:	41	፥	2 6	:	፥	÷	÷	÷	:	:	÷	÷	÷	:
	Length dressed.	ب ر : ښ	9 11	:	:	:	:	፥	:	:	9 34	:	÷	:	:	:	÷	÷	:	÷
	Length of body.	ن <u>د</u> : به	÷	:	0 9	:	6 3	:	6 4	6 4½	፧	÷	፧	:	:	፥	÷	፧	÷	:
	Length before skinned.	ft. in. -9 44	4	4 6-	-9 3 1	-9 3 4	-99 3	6 3	-6 3	-9 3	-9 3	z 65-	1 66-	66-	0 66-	oi 8	80 80 -	4 88-	-98 7	-98 5

TIGER (Felis tigris)—continued.

Skulls.

Basal length from back across the from back archests in the from back				Martin.	
-15½ Kuch Behar . Maharaja of Kuch Behar. -15½ 10½ N.E. Bengal . A. M. Murdoch. -15½ 10½ Purneah . R. A. Sterndale. -15 (A) 10½ Purneah . R. A. Sterndale. -14½ (F) 10 Duars . Capt. S. H. Pollen. 14½ 9½ 4 10 Do Sir Edmund G. Loder, Bart. 14½ 10½ Deccan . W. J. R. Wingfield. -14½ 10 Terai . Bombay Natural History Society's Museum. 14½ 10½ Central Provinces . Capt. W. H. Hunter. -14½ (C) 10½ Bignor District, C.P. A. M. Markham. 14½ D) 9½ Kanara Jungles . W. Holland. 14½ (E) 9½ Terai . R. Nolan. -14 10 ? J. D. Inverarity. -14 10 ?? Bignor District . A. M. Markham. 13½ 9½ Pilon . Pilon . Pilon . Pilip Wood. 13½ 9½ Indhaorèe, Nimar . LieutCol. H. Wade-Dalton. 13½ 9½ North Kanara . LieutCol. L. L. Fenton. 13.70 9 Duars . Mrs. Lawrie-Johnstone. 13½ 9½ North Kanara . LieutCol. L. L. Fenton. 13½ 9½ North Kanara . LieutCol. L. L. Fenton. 13½ 9½ North Kanara . LieutCol. L. L. Fenton. 13½ 9½ North Kanara . LieutCol. L. L. Fenton. 13½ 9½ North Kanara . LieutCol. L. L. Fenton. 13½ 9½ North Kanara . LieutCol. L. L. Fenton.	from back	zygomatic	cleaned.	Locality.	Owner.
Purneah R. A. Sterndale. -15(A) 10\frac{1}{2}	-152	•••		Kuch Behar	Maharaja of Kuch Behar.
Total Tota	-151	107		N.E. Bengal	A. M. Murdoch.
10	-151	101		Purneah	R. A. Sterndale.
14½ (F) 10 Duars Capt. S. H. Pollen. 14½ 9½ 4 10 Do. Sir Edmund G. Loder, Bart. 14½ 10½ Deccan W. J. R. Wingfield. -14½ 10 Central Provinces Capt. W. H. Hunter. -14½ (C) 10½ Bignor District, C.P. A. M. Markham. 14½ 10½ Central Provinces Capt. E. A. D'Arcy Thomas. 14½ D) 9½ Kanara Jungles W. Holland. 14½ (E) 9½ Terai R. Nolan. -14 10 ? J. D. Inverarity. -14 10 ? J. D. Inverarity. -14 10 ? Bignor District A. M. Markham. 13½ 9½ ? Capt. S. H. Pollen. 13½ 9½ ? Capt. S. H. Pollen. 13½ 9½ Philip Wood. 13½ 9½ North Kanara LieutCol. H. Wade-Dalton. 13½ 9½ North Kanara LieutCol. L. L. Fenton. 13½ 0½	-15 (A)	10]		Bengal	Col. Evans Gordon.
14½ 9½ 4 10 Do. Sir Edmund G. Loder, Bart. 14½ 10½ Deccan W. J. R. Wingfield. -14½ 10 Terai Bombay Natural History Society's Museum. 14½ 10½ Central Provinces Capt. W. H. Hunter. -14½ (C) 10½ Bignor District, C.P. A. M. Markham. 14½ 10½ Central Provinces Capt. E. A. D'Arcy Thomas. 14½ (E) 9½ Kanara Jungles W. Holland. 14½ (E) 9½ Terai R. Nolan. -14 10 ? In Inverarity. -14 10 ? D. Inverarity. -14 10 ? Capt. S. H. Pollen. 13½ 9½ Indhaorèe, Nimar LieutCol. H. Wade-Dalton. 13½ 9½ Madras Col. J. Hinde. -13½ 9½ North Kanara <th>15</th> <th>101</th> <th></th> <th>?</th> <th>Rowland Ward.</th>	15	101		?	Rowland Ward.
14½ 10½ Deccan	14 7 (F)	10	•••	Duars	Capt. S. H. Pollen.
Terai	141	9 3	4 10	Do	Sir Edmund G. Loder, Bart.
Museum. Capt. W. H. Hunter.	141	108	•••	Deccan	W. J. R. Wingfield.
14\frac{1}{2} 10\frac{1}{3} Central Provinces . Capt. W. H. Hunter. -14\frac{1}{2}(C) 10\frac{1}{4} Bignor District, C.P. A. M. Markham. 14\frac{1}{4} 10\frac{1}{3} Central Provinces . Capt. E. A. D'Arcy Thomas. 14\frac{1}{4}(D) 9\frac{1}{2} Kanara Jungles . W. Holland. 14\frac{1}{4}(E) 9\frac{1}{2} R. Nolan. -14 10 ? I. D. Inverarity. -14 10\frac{1}{10} Bignor District . A. M. Markham. 13\frac{1}{8} 9\frac{1}{4} ? Capt. S. H. Pollen. 13\frac{1}{8} 9\frac{1}{4} ? Philip Wood. 13\frac{1}{4} 9\frac{1}{2} North Kanara . LieutCol. L. L. Fenton. 13\frac{1}{4} 9\frac{1}{2} Mrs. Lawrie-Johnstone. -13\frac{1}{2} 9\frac{1}{2} Major J. W. M. Cotton. 13\frac{1}{6} 9\frac{1}{6} Kumaun Terai . H. J. Boas.	$-14\frac{1}{2}$	10	•••	Terai	
141 108 Central Provinces Capt. E. A. D'Arcy Thomas. 141 D) 9½ Kanara Jungles W. Holland. 141 (E) 9½ Terai R. Nolan. -14 10 ? J. D. Inverarity. -14 10½ Bignor District A. M. Markham. 13½ 9½ ? Capt. S. H. Pollen. 13½ 9½ Indhaorèe, Nimar LieutCol. H. Wade-Dalton. 13½ 9½ ? Philip Wood. 13¾ 9½ North Kanara LieutCol. L. L. Fenton. 13·70 9 Duars Major J. W. M. Cotton. 13½ 9½ 3 14 ? Major J. W. M. Cotton. 13½ 9½ Kumaun Terai H. J. Boas.	141	101	•••	Central Provinces .	
14½ D) 9½ Kanara Jungles W. Holland. 14½ (E) 9½ Terai R. Nolan. -14 10 ? J. D. Inverarity. -14 10 ½ Bignor District A. M. Markham. 13½ 9½ ? Capt. S. H. Pollen. 13½ 9½ Indhaorèe, Nimar LieutCol. H. Wade-Dalton. 13½ 9½ Madras Col. J. Hinde. -13½ 9½ North Kanara LieutCol. L. L. Fenton. 13·70 9 Duars Mrs. Lawrie-Johnstone. -13½ 9½ 3 14 ? Major J. W. M. Cotton. 13½ 9½ Kumaun Terai H. J. Boas.	-14½ (C)	101		Bignor District, C.P.	A. M. Markham.
14½ (E) 9½ Terai R. Nolan. -14 10 ? J. D. Inverarity. -14 10½ Bignor District A. M. Markham. 13½ 9½ ? Capt. S. H. Pollen. 13½ 9½ Indhaorèe, Nimar LieutCol. H. Wade-Dalton. 13½ 9½ Madras Col. J. Hinde. -13½ 9½ North Kanara LieutCol. L. L. Fenton. 13.70 9 Duars Mrs. Lawrie-Johnstone. -13½ 9½ Xumaun Terai H. J. Boas.	142	108	•••	Central Provinces .	Capt. E. A. D'Arcy Thomas.
-14 10 ? J. D. Inverarity. -14 10 10 10 Bignor District . A. M. Markham. 13 10	14½ D)	91	•••	Kanara Jungles .	W. Holland.
-14 10 16 Bignor District . A. M. Markham. 13 18 9 1 ? Capt. S. H. Pollen. 13 19 9 1 Indhaorèe, Nimar . LieutCol. H. Wade-Dalton. 13 19 9 1 Madras Col. J. Hinde. -13 18 9 1 North Kanara . LieutCol. L. L. Fenton. 13 70 9 Duars Mrs. Lawrie-Johnstone. -13 19 9 1	14 1 (E)	9 1	•••	Terai	R. Nolan.
13\frac{3}{6} 9\frac{3}{4} ? Capt. S. H. Pollen. 13\frac{3}{4} 9\frac{1}{4} Indhaorèe, Nimar LieutCol. H. Wade-Dalton. 13\frac{3}{4} 9\frac{1}{2} ? Philip Wood. 13\frac{3}{4} 9\frac{1}{2} Madras Col. J. Hinde. -13\frac{3}{4} 9\frac{1}{2} North Kanara LieutCol. L. L. Fenton. 13\cdot 70 9 Duars Mrs. Lawrie-Johnstone. -13\frac{1}{2} 9\frac{1}{2} 3 14 ? Major J. W. M. Cotton. 13\frac{3}{6} 9\frac{7}{6} Kumaun Terai H. J. Boas.	-14	10		?	J. D. Inverarity.
138 9½ Indhaorèe, Nimar LieutCol. H. Wade-Dalton. 13½ 9½ ? Philip Wood. 13¾ 9½ Madras Col. J. Hinde. -13¾ 9½ North Kanara . LieutCol. L. L. Fenton. 13·70 9 Duars Mrs. Lawrie-Johnstone. -13½ 9½ 3 14 ?	-14	10 <mark>.9</mark>	•••	Bignor District .	A. M. Markham.
13½ 9½ ? Philip Wood. 13½ 9½ Madras Col. J. Hinde. -13½ (B) 9½ North Kanara . LieutCol. L. L. Fenton. 13·70 9 Duars Mrs. Lawrie-Johnstone. -13½ 9½ 3 14 ? Major J. W. M. Cotton. 13½ 9½ Kumaun Terai . H. J. Boas.	13 7	9 3	•••	?	Capt. S. H. Pollen.
13½ 9½ Madras Col. J. Hinde. -13½ (B) 9½ North Kanara . LieutCol. L. L. Fenton. 13·70 9 Duars Mrs. Lawrie-Johnstone. -13½ 9½ 3 14 ? Major J. W. M. Cotton. 13½ 9½ Kumaun Terai . H. J. Boas.	138	9 1		Indhaorèe, Nimar .	LieutCol. H. Wade-Dalton.
-13½ (B) 9½ North Kanara . LieutCol. L. L. Fenton. 13·70 9 Duars Mrs. Lawrie-Johnstone. -13½ 9½ 3 14 ? Major J. W. M. Cotton. 13½ 9½ Kumaun Terai . H. J. Boas.	138	98		5	Philip Wood.
-13\frac{3}{4}(B) 9\frac{1}{4} \qquad \text{North Kanara} \qquad \text{LieutCol. L. L. Fenton.} \\ 13.70 \qquad 9 \qquad \text{Duars} \qquad \text{Mrs. Lawrie-Johnstone.} \\ -13\frac{1}{4} \qquad 9\frac{1}{4} \qquad 3 \qquad 14 \qquad ? \qquad \text{Major J. W. M. Cotton.} \\ 13\frac{3}{6} \qquad 9\frac{7}{6} \qquad \text{Kumaun Terai} \qquad \qquad \text{H. J. Boas.} \end{aligned}	132	91	•••	Madras	Col. J. Hinde.
13.70 9 Duars Mrs. Lawrie-Johnstone. -13½ 9½ 3 14 ? Major J. W. M. Cotton. 13¾ 9½ Kumaun Terai . H. J. Boas.	-13 3 (B)	91		North Kanara .	LieutCol. L. L. Fenton.
138 95 Kumaun Terai . H. J. Boas.		9		Duars	Mrs. Lawrie-Johnstone.
-38	-131	91	3 14	?	Major J. W. M. Cotton.
		97		Kumaun Terai .	H. J. Boas.
	-I2 1 8	8,5	•••	Bignor District, C.P.	A. M. Markham.

Skins.

Length of ski	in dressed.	Locality.		Owner.	
ft. I 3	in. 6	Chinese Mongolia			A. Bignold.
12	6	Do.			Rowland Ward.
11	31/2	Do.			W. H. Walker.
12	0	Do.			H.R.H. le Duc d'Orléans.

A, B, C, D, E, F. For skins see p. 460.

LEOPARD (Felis pardus).

The leopard has an even larger range than the lion, and is probably the most widely distributed of all the cats. Sportsmen in India recognise a larger "panther" and a smaller "leopard," although naturalists fail to distinguish them as separate forms. The African and Indian leopards are also generally regarded as identical, but since the former has the spots on the body of smaller size than in the latter, and many of them without light centres, like those on the head, it is quite probable that it constitutes a different race. In Persia, Baluchistan, and the mountains of Sind is found a variety of the leopard (F. pardus tulliana) characterised by its pale colour, long fur, and thick tail; it is to some extent intermediate between the typical race and the ounce. Lastly, there is the Chinese leopard (F. pardus fontanieri), a very distinct animal, of heavy build, with a pale groundcolour to the fur, which is very long and thick, and the spots in the form of large complete rings, without any smaller spots in the centre. In the moist forest districts of Asia black leopards are by no means uncommon; and there is a tendency to darkening of a different type in some South African specimens. Height at shoulder from about 2 feet (India) to 2 feet 4 inches (Africa). A large leopard killed by Lieut.-Col. L. L. Fenton in the Gir forest, Kathiawar, measured 7 feet 8½ inches in length; the length of the tail being 35½ inches, the girth of the neck 211 inches, of the forearm 111 inches, and of the body behind the shoulder 35\frac{1}{2} inches; weight, 160 lbs.

Distribution.—Africa, the Caucasus, Asia Minor, and Asia generally, with the exception of Northern Siberia, the Tibetan plateau, and perhaps Japan.

a.-Indian Specimens.

							M	ale	Fer	male
							ft.	in.	ft.	in.
Length from tip	of no	se to e	end o	of tail			7	0	6	4
,,	"	r	oot o	of tail			4	$3\frac{1}{2}$	3	7
Height at shoul	der						2	$2\frac{1}{2}$	1	9
Girth	•						2	7	2	1
Circumference	of uppe	er arm		•			0	13	0	10
,,	forea	ırm	•		•		0	108	0	81

Shot in Ceylon by Count Scheibler.

LEOPARD (Felis pardus). Indian Specimens—continued.

Len bef skin	ore	Total length dressed.	Weight.	Locality.		Owner.
ft. -8	in. 4	ft. in.	lbs.	Kuch Behar .		Maharaja of Kuch Behar.
-8	218	•••		Do		Do.
-8	112	•••		Do. .		Do.
-8	0	•••	154	Do. .		Do.
-8	0	8 81		India		Count J. Potocki.
-7	10	8 4 3	•••	Ganges Kadir .		See below.1
-7	91	8 6	•••	Central Provinces		James J. Harrison.
-7	9	•••	•••	Nepal		C. H. H. B. Caldwell.
-7	9	•••	•••	Oudh		Mrs. Innes.
-7	8	•••	•••	Hyderabad, Deccan		Capt. C. E. G. Norton and Capt.
-7	8	•••	•••	Ganges Kadir .		Sir K. Fraser, Bart. Muttra Tent Club. ²
-7	7	8 7 1	•••	Siwaliks	•	E. H. E. Green.
		7 10		?		Col. W. Gordon Cumming.
-7	6	•••	•••	Chaubattia		Capt. R. B. Fell.
-7	1	7 10 1	•••	Rewa, C.P.		Count E. Hoyos.
-7	41	•••		Gujerat		Capt. H. J. Morphy.
- 9 7	4	8 I	145	Bignor District .		SurgCapt. E. McK. Williams.
-7	4			Kathiawar		LieutCol. L. L. Fenton.
-7	3		•••	?		LieutCol. F. H. Whitby.
-7	1	7 II I		Kotah		Maharaja of Bikanir.
-7	1		•••	Hyderabad, Deccan		Major G. A. L. Carew, D.S.O.,
-7	o <u>ł</u>	8 31/2		?		and Capt. G. L. Holdsworth. Maharaja of Bikanir.
7	0			?		Major A. E. Ward.
-6	10	7 10	140	Bignor District .		SurgCapt. E. McK. Williams.
-6	10	•••	•••	Chaubattia		Capt. R. B. Fell.
-6	8	•••	110	?		M. Loam.
- 96	5			Chaubattia		Capt. R. B. Fell.

Speared by Capt. H. Hoare (5th D.G.), Capt. J. G. Rotton (R.A.), and W. Gillman (R.H.A.)
 Major A. C. King and some officers of the 5th Lancers, when out with the Muttra Tent Club.

LEOPARD (Felis pardus). Indian Specimens—continued.

Len bek skin	ore	Total dres	length sed.	Weigh.	Local	lity.			Owner.
ft.	in.	ft.	in.	lbs.					
	••	7	8 1	•••	Kuch Behar	•	•	•	Maharaja of Kuch Behar.
- 8 6	2	6	5		Gwalior .	•			Viscount Edmond de Poncins.
- 9 5	11	•	••		Chaubattia.	•	•		Capt. R. B. Fell.
- 8 5	10	6	81	96	Bignor Dist	rict			SurgCapt. E. McK. Williams.
- 8 5	8	6	4	80	Do.				Do.
		7	8	•	Kuch Behar	•		•	Maharaja of Kuch Behar.
1.	••	7	8		?)			Capt. A. G. Ferguson.
		7	6 <u>1</u>	140	Kuch Behar	•			Maharaja of Kuch Behar.
		7	41/2		?	•			Col. Howard.
	••	7	4	• • •	Nepal .				H. L. Heber-Percy.
		7	I	•••	Do.	•	•		Sir E. L. Durand, Bart., C.B.
		6	8	110	Madras .				M. Loam.

b. AFRICAN SPECIMENS.

			«· ши ши опи	 D-0-1-10.
Length skint ft.		Weight.	Locality.	Owner.
-7	31		?	Count J. Potocki.
-7	3	•••	Wadelai, E. C. Africa	Capt. W. P. Pulteney.
-7	2	•••	Somaliland	Count Scheibler.
-7	2	•••	South-East Africa .	F. Vaughan Kirby.
-7	11/2	•••	?	C. V. A. Peel.
-7	0		Somaliland	J. Johnston-Stewart.
- 6	11		E. C. Africa	James J. Harrison.
-6	10		Somaliland	Count Scheibler.
-6	10	140	Nyasaland	A. White.
-6	3	•••	Somaliland	Norman B. Smith.
1-6	2	•••	Jebba, Nigeria .	The late Capt. H. W. Baker.
-6	0	•••	Somaliland	Count E. Hoyos.
- ♀ 6	0		Do	C. V. A. Peel.
² -5	7		Nigeria	LieutCol. T. D. Pilcher.

1 Height at shoulder, 2 ft. 11 in. 2 Height at shoulder, 22 in.; girth, 211 in.

The following specimens were shot and measured by F. Vaughan Kirby:—

(a) Nguanetsi River, S.E.A. (b) Matamiri Bush, S.E.A.

Lengt tip in a li	h, st ine.	raight	over 'sport	ngth all— sman's' rement.	Shou heig		Girth of neck.	Girth behind shoulder.	Girth of forearm.	Length of skull.	Zygomatic width.	Clea weig	ined ght.
ſ	t.	in.	ft.	in.	ft.	in.	in.	in.	in.	in.	in.	lbs.	oz.
(a) (6	II	7	2	2	7	22	35½	121	9 7	6 1	I	8
(b) (6	9	7	2	2	7	20	33 1	111	9	5 1		

Leopard Skulls.

Total length.	Width.	Weight.	Locality.	Owner.			
in. -10 1	in. 6 §	lbs. oz.	Bignor District, N.W.P.	A. M. Markham.			
-10 18	6₺		Gir Forest, Kathiawar .	LieutCol. L. L. Fenton.			
-918		1 5	Nyasaland	Dr. Percy Rendall.			
-9 1	6	clean 	?	Julius Jeppe.			
-9 <u>1</u>		•••	? .	Do.			
-9	51	•••	S.E. Africa	F. Vaughan Kirby.			
-6 7	5		India	W. T. Blanford.			

OUNCE or SNOW-LEOPARD (Felis uncia).

Although the Persian leopard is in some degree intermediate in respect to colour and the length of the coat, the ounce differs from the leopard by the ground-colour of the long and dense fur being dirty white, with the spots on the back, sides, and tail, in the form of large, irregular, ill-defined, and interrupted rings, and by the great thickness of the tail, which scarcely tapers, and is about three-quarters the length of the head and body. Height at shoulder about 2 feet 4 inches. Skull about 6 inches long.

Distribution.—The high ranges of Central Asia, including Gilgit, Hunza, Turkestan, Trans-Baikalia, Ladak, Tibet, Amurland and Western China, extending in the north-west to the Altai, and in the west, it is said, to Persia. In Prince Demidoff's Hunting Trips in the Caucasus a snow-leopard is figured as coming from that range, but

whether the animal found there is F. uncia or F. pardus tulliana may be doubtful. Generally dwelling at elevations of over 8000 feet, descending in Gilgit during winter to 6000 feet.

Length dressed.		Weight.	Locality.	Owner.	
ft. 8	in. 7	•••	Tibet	Rowland Ward.	
-8	6	Skull, 7\frac{3}{4} + 5\frac{1}{4}	Kashmir	R. Rankin.	
8	2 <u>1</u>		Baltistan .	E. Langworthy.	
-7	10		Near Leh .	Major S. Frewen.	
-7	5 1		Pamir	Viscount Edmond de Poncins.	
7	5	•••	?	Major F. J. Harden.	
-7	4	(24 inc	hes at shoulder)	R. A. Sterndale.	
-7	3		?	Major A. E. Ward.	

CLOUDED LEOPARD (Felis nebulosa).

The arimau dahan, as this species is called by the Malays, is the size of a small leopard, with a very long and thickly furred tail, and large upper tusks. The ground-colour of the fur varies from grayish brown to fulvous, upon which are large dark blotches, frequently bordered in part with black. In old specimens the blotches often disappear, leaving only the black borders.

Distribution.—From the Sikim and Bhutan Himalaya, through Assam, Burma, Siam, and the Malay Peninsula, to Sumatra, Java, and Borneo. Represented by a smaller variety (F. nebulosa brachyurus) in Formosa.

Length dressed.		Weight.	Locality.			Owner.	
ft. -6	in. 6	lbs.	Assam			٠	The late B. H. Hodgson.
6	4	441		?			R. A. Sterndale.
5	9			?			B. R. M. Glossop.
5	7 1	•••	Assam	٠			The late B. H. Hodgson.
5	71/2		Do.	•	•	•	P. Russel.
				Sku	LL.		
Basal length from back to front.		Breadth.	L	ocalit	у.		Owner.
6.2 in.		4-75	Assam				The late B. H. Hodgson.

JAGUAR (Felis onca).

This cat may be regarded as the American representative of the leopard, which it fully equals, even if it does not exceed, in size. The colour and markings are generally similar to those of the latter, but the dark rings are larger and arranged more definitely in groups, each ring usually enclosing one or more dark central spots, and the enclosed light area being of the same tint as the general ground-colour of the fur, which is typically of a rich tan. Usually seven or eight more or less distinct longitudinal rows of rosettes may be noticed on each side of the body.

Distribution.—America, from Louisiana, Texas, and Northern Mexico to about the Rio Negro, on the northern confines of Patagonia in lat. 40° S.

Length before skinning.	Length dressed.	Locality.		Owner.
ft. in. -9 3	ft. in. 9 3	Brazil .		Count Henry Coudenhove.
-8 3 3		Do. .		Do.
-6 11	79	Paraguay .		James J Harrison.
	7 7	British Guiana		Sir A. W. L. Hemming, K.C.M.G.

PUMA (Felis concolor).

With the exception of the much smaller and longer-tailed yaguarondi, the puma (pronounced pooma) is the only uniformly coloured cat found in America, where its range extends from British Columbia and Maine in the north to the Strait of Magellan in the south. The size is inferior to that of the jaguar, the height at the shoulder being about 2 feet, and the weight 150 lbs. The general colour of the fur is tawny, tending, like that of the Virginian deer, to reddish in summer, and to grayish in winter, with the middle of the back darker, and a whitish patch on the back of the otherwise black ears. But with such an enormous geographical range, it is inevitable there should be much local variation; and American writers recognise more than one species. F. concolor oregonensis, of the north-west coast, may be admitted as a local race, as may F. concolor puma of South America.

Leng the	th in field.	Weight.	Locality.	Owner.
ft. -8	in. 6	lbs. about 150	•••	A. Pendarves Vivian.
-7	6₹	•••	Brazil	Count Henry Coudenhove.
-7	5		Gallegos River, Patagonia.	W. Moncreiffe.
-7	3		Fraser River, British Columbia.	J. Fannin.
-7	0		Wyoming	J. L. Scarlett.
-5	9		South America	James J. Harrison.

LYNXES (Felis lynx, etc.).

The lynxes form a well-marked group connected with the more typical members of the cat tribe by the jungle-cat (F. chaus), and distinguished by the tuft of long hairs at the summit of the ears, and the absence of the first upper cheek-tooth. From the others the caracal (F. caracal), of Africa and India, is broadly distinguished by its long tail and uniform rufous colour. The short-tailed lynxes are, however, a group in which it is very difficult to determine whether the variations indicate distinct species or local races. In the typical European lynx (F. lynx) the tail is very short, the throat has a ruff of long hair, and the coat is spotted with dark brown. The Tibet lynx (F. lynx isabellina) is certainly only a pale variety. Probably the Canadian lynx (F. lynx canadensis), which may be identical with the lynx of Northern Europe and Asia, should hold the same rank; it ranges in America as far south as California. More doubt exists whether the red lynx (F.rufa), which is also widely extended in America, should be regarded as a separate race, but the so-called plateau lynx (F. baileyi) seems certainly only a variety of the former inhabiting the high grounds of Colorado, Utah, and Arizona. On the other hand, from differences in the form of the skull, the Spanish lynx (F. pardina), which is a fully spotted animal, seems rightly regarded as a distinct species. It ranges over a large portion of Southern Europe, including Turkey, Greece, Sicily, Sardinia, and Spain and Portugal. The ordinary lynx stands from 16 to 18 inches at the shoulder.

a. EUROPEAN LYNX.

Length on the field, nose to root of tail.

Locality.

Owner.

-42·9 in.

. Eastern Carpathians

. Prince Henry of Liechtenstein.

b. TIBET LYNX.

Height at shoulder, 16 or 18 inches.

c. CANADIAN and RED LYNX.

Length on the field, nose to root of tail.	Height at shoulder.	Weight.	Locality			Owner.
1-38 in.	25	60 lbs. (about)	Wyoming .			Capt. G. Dalrymple White.
² –32 ,,	22	40 lbs. (about)	Nova Scotia	•		Do.
		¹ Canadian I	Lynx.		:	Bay Lynx.

d. CARACAL.

Flat skin.	Locality.	Owner.
46½ in.	Nr. Grahamstown, S. Africa	Dr. H. Smith.

HUNTING-LEOPARD or CHITA (Cynælurus jubatus).

Although this animal is commonly called chita (cheetah) by Anglo-Indian sportsmen, that name is at least as often applied in India to the leopard. From all the true cats and lynxes the hunting-leopard differs by the claws being capable of only partial withdrawal into their sheaths, so that their tips are always exposed. The body also is more slender, and the limbs are proportionately longer. The black spots on the skin are small and without light centres, like those on the head of the leopard. Length 7 feet or less, height at shoulder 30 to 39 inches. A specimen speared by Lieut.-Col. L. L. Fenton in Kathiawar measured 6 feet $\frac{1}{4}$ inch in length, the tail being 2 feet $\frac{1}{24}$ inches.

The animal is the "Ihlose" of the Zulus and Swazis, and is distributed sparsely throughout S.E. Africa. They usually hunt in couples, and fairly stalk their game, securing it with a swift rush at the last. Mr. F. Vaughan Kirby says: "I have seen a party of six hunting together and another of eight. Though I have often tried, I have never yet succeeded in running into one on horseback; they are incredibly swift of foot. They invariably kill their prey by strangulation." Distribution.—Africa and South-Western Asia, extending from Persia

to Western Turkestan and the countries east of the Caspian, and eastwards into India; unknown in the latter country on the Malabar coast and to the north of the Ganges, as it also is in Ceylon. The African hunting-leopard is stated to differ by its more woolly coat, and if this be the case, should be known as C. jubatus laneus.

Shot in the Eastern Transvaal by F. Vaughan Kirby.

Total length in straight line.	Do. over all.	Tail.	Vertical height.	Girth of forearm.	Do. shoulders.
6 ft. 8 in.	7 ft. 7 in.	2 ft. 9 in.	2 ft. 11½ in.	8 1 in.	31 in.

	gth.	Length of tail.	Height at shoulder.	Locality.	Owner.
ft. -7	in. O	ft. in. 		India	R. A. Sterndale.
-6	o l	2 24		Near Rajkot, Kathiawar	LieutCol. L. L. Fenton.
6	4 4	2 31/2	28 <u>1</u>	N.E. Transvaal	Dr. Percy Rendall.
-6	4	•••	•••	90 lbs. Somaliland	J. Johnston-Stewart.
-6	1 1/2			Do	C. V. A. Peel.
6	0	•••	•••	Do	Do.

SPOTTED HYÆNA (Hyæna crocuta).

Somali *Uaraba*. Danakil *Yangula*. Abyssinian *Jib*. Setongwani in the Barotse country. Piri in the Lake Ngami country. Chimbwi in the Chilala and Chibisa countries.

The hyænas form a small family of Carnivora allied in some respects to the cats, but distinguished by the form of the skull, the more numerous teeth (which are, however, to a considerable extent cat-like), and the four-toed feet, with non-retractile claws. The spotted hyæna is the largest of the three species, and takes its name from the large dark blotches on its tawny coat. Its carnassial teeth are more cat-like than those of the other species.

Distribution.—Africa, south of the Sahara.

Nose to root of tail in a straight line, 4 feet $9\frac{1}{2}$ inches. Length of tail, 1 foot 1 inch.

Extreme length over all, 6 feet 3 inches.

Vertical standing height, 3 feet.

Girth behind shoulders, $39\frac{1}{2}$ inches.

SKULLS.

Length.	Width.	Weight.	Locality.	Owner.
in. -12‡	in. 7 1	lbs. oz. 2 8	South-East Africa .	F. Vaughan Kirby.
-121	78	2 12	Zomba, B.C.A.	D. MacAlpine.
-11	$7\frac{1}{2}$		Somaliland	Viscount Edmond de Poncins.
- t I	6 7		South-East Africa .	F. Vaughan Kirby.
-10 §	67	2 3	East Africa	Capt. R. A. J. Montgomerie, C.B., R.N.

BEARS 473



Brown Bear.

BEARS (Ursidæ).

Bears are so unlike other animals and so like one another that no one has the slightest difficulty in recognising a member of the group when he sees it. They constitute a family of Carnivora—the *Ursidæ*—and are spread over the greater part of the globe, with the exception of Africa to the south of the Sahara desert, and the Australasian islands. As some of their leading characteristics may be mentioned their large bodily size, clumsy build, shaggy fur, generally uniform coloration, the very short tail, and the application of the whole sole of the foot to the ground in walking. The skull and teeth are likewise very peculiar and

distinctive, although these need not be taken into consideration in this place.

In consequence of the marked similarity to one another of most members of the group, it is a matter of extreme difficulty to come to a definite conclusion as to the number of species of bears—not that this is a matter of very much importance one way or another. The typical member of the group is the familiar brown bear (Ursus arctus) of Europe, whose colour is generally a darker or lighter shade of brown, but occasionally tends to grayish. The Syrian bear (U. arctus syriacus), in which this grayish tinge predominates, may be regarded as a local variety, and the same is the case with the snow-bear of Kashmir (U. arctus isabellinus), in which the colour is generally a light creamy brown. European specimens probably seldom exceed eight feet, but the huge Kamschatkan bear (U. arctus collaris) grows to nine feet. Even more gigantic is the Kadiak bear (U. arctus middendorfi) of Kadiak Island, Alaska; while the Yezo bear (U. arctus yesoensis) of Japan is another large form, with much the external appearance of a grizzly. The Alaskan bear (U. arctus dalli), from the mainland of Alaska, is also a huge animal, slightly smaller than the one from Kadiak Island, with more resemblance to an ordinary brown bear than to a grizzly. The typical grizzly of the Rocky Mountains is a smaller animal, with longer and straighter claws, but scarcely entitled to rank as more than another local race (U. arctus horribilis), next to which comes the Barren Ground bear (U. arctus richardsoni). Finally, the last animal which can be included in this group of the genus is the African bear (U. arctus crowtheri), of North-Western Africa, still imperfectly known. The extinct cave-bear (U. spelæus) is a large species allied to the brown bear.

The little blue bear (*U. pruinosus*) of Tibet, with more or less of white on the head and shoulders, seems to form a distinct species. The same is the case with the American black bear (*U. americanus*), which is generally black, and exhibits distinctive features in the skull and teeth. Not improbably the Himalayan black bear (*U. torquatus*) is a relative of the last-named species; it may be recognised by the conspicuous white gorget on the breast. In Japan it is represented by the Japanese black bear (*U. japonicus*). The smallest species of the genus are the very distinct Malayan bear (*U. malayanus*), and the allied spectacled bear (*U. ornatus*) of the Peruvian Andes, the latter distinguished by the light-coloured rings around the eyes, from which it derives its name.

The most distinct of all the species included in the genus Ursus is

BEARS 475

the Polar bear (*U. maritimus*), so distinct, indeed, that many naturalists think it ought to form a group by itself. Externally its chief characteristics are its white coat, and the presence of a certain amount of hair on the soles of the feet; both these peculiarities being evidently adaptations to the Arctic habitat of the animal. It has been recently stated that very old Polar bears exhibit a tendency to the development of a brownish tinge in the fur. Last of all come the Indian sloth-bear (*Melursus ursinus*), and the parti-coloured bear (*Æluropus melanoleucus*) of Tibet, both of which are so different from the other kinds as each to form a genus by itself. The former is too well known an animal to need description, some of its characteristics being the long and bare snout, the ragged, wiry hair, extensile tongue, small cheek-teeth, and the diminished number of front teeth. From all its kindred the second differs by its pied fur, as it does by its remarkably-formed cheek-teeth, which are broader and shorter than in other bears.

a-SKULLS.

from back to front.	Width across the zygo- matic arches.	Weight cleaned.	Race, or Species.	Locality.	Owner.
in. 183	in. 1 1 🖁	lbs. oz. I2 O	Cave bear	Europe	Sir Edmund G. Loder, Bart.
18	11	10 0	Kamschat- kan	Siberia	Hon. Walter Rothschild.
-16 1	•••		•••	Alaska	H. W. Seton-Karr.
16	98	6 3	Polar .	Arctic Seas .	Rowland Ward.
16	9 7	58	Do	Nova Zembla.	J. Lamont.
15½ (D)	9 1	5 10	Grizzly .	Brit. Columbia	Sir Peter Walker, Bart.
158	94	4 10	Kamschat- kan	Kamschatka .	Lieut. R. E. R. Benson, R.N.
15 2	10½	5 13	Polar .	Polar Seas .	Sir Edmund G. Loder, Bart.
141	. 9	5 14	Do	Franz Josef Land	Jackson-Harmsworth Expedition.
141	8 1	4 8	Grizzly .	Brit. Columbia	S. B. Bennett.
14 8 (C)	81	5 O	Do	New Mexico .	Montague Stevens.
13	9 1		Polar .	Spitzbergen .	The late Capt. Townley Parker.
$-12\frac{7}{8}$	7		Sloth .	Mysore	Capt. M. M'Neill.
121	78		Do	Cent. Prov	C. F Egerton.
I 2 ½	7		Grizzly .	Wyoming .	J. L. Scarlett.
12½ (B)	63		Sloth .	Cent. Prov	Surgeon-Major M. O'C. Drury.
12 8	7₹		Snow .	Kashmir .	A. Ezra.
121	7₩		Grizzly .	Wyoming .	J. L. Scarlett.
12	7 1 8		Sloth .	Cent. Prov	C. F. Egerton.
113	71/2		Snow .	Kashmir .	W. R. Bindloss.
11½ (A) 6 <u>3</u>		Brown .	W. Caucasus.	St. George Littledale.
-10	6.8	٠	Black .	Nepal	W. T. Blanford.
-8.5	8.3	•••	Malayan .	Borneo	Do.

BEARS (Ursidæ)—continued.

b—skins.

nose	th from to tip tail.	Height at shoulder.	Girth.	Weight.	Race, or Species.	Locality.	Owner.
ft. -13	in. 6 ¹	ft. in.	ft. in.	lbs. 1656	Kadiak Island	English Bay, Kadiak Is-	J. C. Tolman.
-11	6 <u>1</u> 1	IO 2 ² (from one front paw		1536	3	land, Alaska Nevada, 1881	W. F. Sheard.
9	101	to other)	• • • •		Alaskan	Alaska	Rowland Ward.
9	81				Do.	Do	Cape Town Museum.
9	5 ¹				Do.	Do	Dublin Museum.
9	5	4 6			Polar .	Baffin Bay .	W. Livingstone Lear-
9	5¹				Alaskan	Alaska	month. Hon. Walter Rothschild.
8	101	4 O (mounted)			Do.	Do	British Museum.
8	7			1600	Polar .	Arctic Seas .	Captain Lyon.
8	7				Do	Spitzbergen .	A. Barclay Walker.
8	6				Do	Franz Josef	Dr. S. H. T. Armitage.
-8	5				Black .	Land Kashmir .	Capt. W. Westropp White.
8	4				Do	?	SurgCapt. W. White.
8	3				Kams- chatkan	Kamschatka .	Dr. F. H. H. Guillemard.
- 8	οł				Grizzly .	Wyoming .	T. W. H. Clarke.
8	0				Polar .	Spitzbergen .	A. Barclay Walker.
8	O(D)	3 11		•••	Grizzly .	Brit. Columbia	Sir Peter Walker, Bart.
-8	o (c)	•••		735	Do	New Mexico	Montague Stevens.
-7	11	•••		800 (about)	Brown .	Norway .	Capt. Gerard Ferrand.
-7	9				Snow .	Kashmir .	Major C. F. Blane.
7	5 ½				Black .	Wyoming .	T. W. H. Clarke.
-7	ī				?	N.W. Terri-	James J. Harrison.
-7	03	••.		680 (about)	Brown .	tory Lithu ani a .	Prince Radziwill.
7	0	3 2			Do	India	Major A. E. Ward.
-6	11		•••	•••	Black .	Wyoming .	James J. Harrison.

¹ Skin measurement,

² Measurement of head in the flesh from tip of nose to base of skull, 29 inches; between ears, 16 inches.

BEARS (Ursidæ). Skins-continued.

nose	h from to tip tail.	Height at shoulder.	Girth.	Weight.	Species.	Locality.	Owner.		
st. -6	in. IO	ft. in. 3 O	ft. in. 4 4½ forearm	lbs. 423	Sloth .	Kuch Behar .	Maharaja of Kuch Behar.		
-6	91		25	•••	Snow .	Kashmir .	S. V. Occleston.		
-6	8 (B)	•••		•••	Snow .	Do	A. Ezra.		
-6	6	•••	•••	•••	Grizzly .	Wyoming .	Count E. Hoyos.		
6	5				Do	Do	J. L. Scarlett.		
-6	41/2		•••	•••	Snow .	?	Capt. H. M. Biddulph.		
-6	3	•••			Grizzly .	Wyoming .	J. L. Scarlett.		
-6	21	•••			Black .	Brit. Columbia	Count E. Hoyos.		
- \$ 6	2			•••	Grizzly .	Cent. Prov	Capt. M. M'Neill.		
-6	11	•••			Sloth .	Mandla, Cent.	Capt. B. H. Boucher.		
-6	ı	•••			Cinnamon	Provinces Wyoming .	James J. Harrison.		
-6	O1	•••			Snow .	Chamba .	Do.		
-6	o <u>ł</u>		•••		Sloth .	Hyderabad .	Count E. Hoyos.		
-6	0	3 0		280	Do	?	The late Col. G. P.		
-5	101	3 0	3 4		Red .	?	Sanderson. Capt. H. M. Biddulph.		
-5	10		••• .	•••	Black .	Kashmir .	Major C. F. Blane.		
-5	9		•••	•••	Sloth .	Cent. Prov	Capt. M. M'Neill.		
-5	8		•••	•••	Do. .	Do.	James J. Harrison.		
-5	5 1	•••	•••	216	Do	?	Capt. H. M. Biddulph.		
-5	5	2 5 4	2 10 ⁸	250	Black .	Brit.Columbia	Count Scheibler.		
- 9 5	21/2	2 112	4 4	about 600	Grizzly .	Do.	Do.		
-5	2 (A))		about	Brown .	W. Caucasus	St. George Littledale.		
5	2	•••	•••		Snow .	Chamba .	Capt. B. H. Boucher.		
-5	1 1/2	•••		200	Brown .	Wyoming .	Count Scheibler.		
- 9 5	1.65	•••		about 	Snow .	Kandgoot .	Viscount Edmond de		
-4	11	, ···			Sloth .	Ganjam Dis-	Poncins. M. Loam.		
-4	9.65				Do	trict Odeypore .	Do.		
-4	7				?	Asia Minor .	H. O. Whittall.		

N.B.—Some of the specimens entered as "Grizzly" may be the barren ground bear.

WALRUS (Odobœnus rosmarus and O. obesus).

The unwieldy seal-like animals commonly known by a corruption of the Scandinavian name valross (whale-horse) form in some respects a connecting link between the true seals and the eared seals, although differing from both in the huge upper tusks which depend from the muzzle of males and females alike, as also by the thick yellow bristles covering the muzzle itself. Like the true seals, walrus have lost all traces of external ears, but, unlike the former, and like the eared seals, their huge hind-flippers are turned forwards beneath the body when on land. The molar teeth, which are adapted for crushing the shells on which these monsters feed, have simple flattened crowns, quite unlike those of seals. Although young and adolescent walrus have a fairly thick coat of yellowish fur, in old individuals the tough hide becomes almost bare, except for the aforesaid bristles. Walruses are estimated to attain a weight of from 2250 to 3000 lbs.

Walruses are exclusively confined to the Arctic seas, where they spend much of their time on the ice. There are two kinds, now generally regarded as separate species; the one confined to the North Atlantic, and the other to the North Pacific. The distribution of these animals is by no means of circumpolar extent, the Atlantic walrus (O. rosmarus) apparently not ranging on the Asiatic coast east of the mouth of the river Lena; while in America they do not appear to inhabit the vast extent of coast lying between the western shore of Hudson Bay and Alaska. The Pacific walrus, which is the larger animal of the two, with considerably the longer tusks, always had a restricted range, and is now becoming very scarce. In European museums it appears to be represented only by skulls and tusks, and even these are rare. Formerly the Atlantic walrus occurred in countless thousands, but in accessible situations its numbers have been greatly reduced, owing to incessant persecution for the sake of its valuable oil and ivory. Between 1870 and 1880 at least 100,000 of these animals are estimated to have been slain.

The largest walrus ever shot by Mr. W. Livingstone Learmonth measured 12 feet 8 inches in length, and the tusks when extracted measured $25\frac{1}{2}$ inches in length and $8\frac{1}{2}$ in circumference at the largest part, but, as is the case with nearly all old bull walrus, the tusks were much broken at the points.

WALRUS (Odobænus rosmarus and O. obesus)—continued. TUSKS.

Total length of tusk.	Weight.	Girth.	Locality.	Owner.
36	lbs. oz.		?	Sir Thos. Hesketh, Bart.
32 8		88	?	Bethnal Green Museum.
-31	6 I	•••		F. Gordon George.
29	•••	9‡	Arctic North America	C. C. Branch.
283		93	Do.	Do.
$-25\frac{1}{2}$		812	Baffin Bay	W. Livingstone Learmonth.
-19	 Length	812	Spitzbergen	Alex. R. Alston.
	from gum. 22 g	7 3	Do	Sir Edmund G. Loder, Bart.
•••	203	71/2	Do	Sir Victor Brooke's Collection.
•••	18}	7⅓	Do	Arnold Pike.
•••	18	7 ½	Do	Earl of Dunmore.
•••	15‡	6	Kolguev	A. Coats.
	121	7 3	Do	Do.

NARWHAL (Monodon monoceros).

The narwhal is the only member of the Cetacean or whale order furnished with tusks. These are present only in the male, and generally the left one alone is developed. Occasionally, however, both grow, as exemplified in a specimen in the British, and a second in the Cambridge Museum. A model of the animal is exhibited in the British Museum.

Distribution.—Arctic Seas.

The following measurements refer to tusks:-

Len	gth.	Circumference.	Weight.	Localit	y.		Owner.		
ft. 8	in. 8 <u>1</u>	8§	lbs.	Arctic Seas			Bethnal Green Museum.		
8	71/2	81	14	Do.			Rowland Ward.		
8	3 1	7 8		Do.			Bethnal Green Museum.		
8	31	7 1		Do.			Hon. Charles Ellis.		
8	3	7 1	1 7	Do.			Rowland Ward.		
7	103	8 1	•••	Do.			Hon. Charles Ellis.		
7	9	83	I 2 ½	Do.			Sir Edmund G. Loder, Bart.		
7	o	6₹		Do.		•	Duke of Westminster.		

SUGGESTIONS FOR RECORDING THE LENGTH OF ANIMALS IN THE FIELD.

As the animal lies on the ground, and where circumstances permit, proceed as follows:-

Length. Pull the nose and the tail so as to get it in as nearly as possible in a straight line. Fix four pegs in, one at the end of nose, one at end of tail, one at root of tail, and one at the nape of neck behind ears.

Height at shoulder. Put the leg or paw in a standing position and place a peg at top of withers and measure carefully standing height (a) with extended paw, (b) with spread paw, as well as length of fore and hind legs from pegs at the stomach line.

The columns in the game-book would thus be as follows:—

A to F. Straight length from nose to | P. Girth of head. tip of tail.

A to E. Length along curve to root of tail.

E to F. Length of tail.

A to B. Length of head to nape of neck.

N. Girth, upper arm.

M. forearm. ,,

O. of body.

D to G. Height of hind-quarters.

C to I. Height at shoulder.

K to L. Length of fore-leg.

H to G. Length of hind-leg.

Weight, cleaned.

not cleaned.

Cleaned skull (length, breadth, height, weight).

The above measurements together with the note of its sex, estimated age, where and when killed, condition of the animal, with any other features of interest, such as colour of the eyes and skin, length of the hoofs, condition of teeth, weapons used and effect of bullets, etc. etc., will be of interest.



Tiger shot by Major H. G. C. Swayne, Central Provinces.

INDEX OF NAMES

Abdy, Captain, 44 Acland, Sir C. T. D., Bart., 21 Acland, Sir T. D., Bart. (the late), 21 Acland-Hood, Sir A., Bart., 21 Adair, Captain F. E. S., 204, 218, 230, 318, 345, 362, 379, 381, 382 Adams, Dr. Leith (the late), 37 Ailsa, Marquis of, 204 Aitchison, Dr. J., 382, 383 Aitkens, F., 122, 161 Akroyd, C. H., 5, 11, 132, 298 Allan, P. S., 345 Allen, Dr. J. A., 4 Allen, J. W., 122, 134, 393 Allen, W., 31 Allgood, Lieut. A. E., R.N., 5 Alston, Alex. R., 273, 284, 301, 394, 421, Altenburg, Prince, 26 American Exhibition Committee, 44 Amsterdam Zoological Gardens, 291 Ancaster, Earl of, 18 Anderson, Sir James, Bart., 74 Andersson, C. J. (the late), 439 Andrassy, Count Gèza, 25 Andrew, H., 127, 207 Angas, G. F., 289 Ansell, C., 215 Apcar, J. G., 36, 38 Arco, Count, 315 Arco-Zinneberg, Count, 29 Arkcoll, J., 274 Armitage, Dr. S. H. T., 476 Arnold, Major A. J., D.S.O., 108, 180, 184, 267, 285, 397

Ashburnham, Captain C., 448, 450

Ashby, Major J. S., 423, 451

Ashby, F., 11

Ashley, W. W., 48, 105, 113, 148, 156, 171, 172, 249, 257, 262, 274, 301, 304, 311, 429, 437, 452 Assheton-Smith, G. W. D., 22 Atholl, Duke of, 16, 17 Atholl, Dowager Duchess of, 57 Atkinson, H., 139, 191 Atkinson, Lewis, 271 Ava, Earl of, 45 Aylmer, G. Percy V., 17, 395 Babington, W. (the late), 172, 232, 235, 300 Bachelors' Club, 373 Baden-Powell, Col. R. S. S., 457 Bagot, Captain Joscelin, 47 Bailey, Abe, 131, 263, 311 Bailey, A. C., 379 Bailey, Captain W., 378 Baillie-Grohman, W. A., 44, 94, 315, 366 Bainbridge, Captain P. A., 424 Bainbrigge, Dr. W. P. Y., 79, 130, 131, 132, 134, 135, 139, 144, 145, 147, 151, 162, 169, 171, 176, 191, 195, 199, 213, 215, 216, 263, 267, 270, 271, 281, 285, 289, 299, 318, 322, 337, 344, 3⁶2, 379, 413, 421, 431, 440, 457 Baker, Capt. H. W. (the late), 466 Baker, Sir Samuel (the late), 44, 46, 423, 448, 450, 461 Banks, G. H. M., 10, 138, 147, 175, 262, 267, Barber, F. H., 108, 116, 118, 125, 128, 138, 139, 143, 144, 165, 169, 173, 174, 185, 186, 187, 191, 194, 208, 214, 215, 216, 237, 249, 256, 262, 269, 270, 279, 283, 284, 288, 289, 294, 297, 299, 310, 391, 392, 427, 428 Barber, H. M., 118, 135 Barclay, Ford G., 46, 94, 105, 113, 139, 176,

248, 254, 257, 273, 305, 352, 359, 386, 387

Barclay, G. N., 140 Barclay, Hugh G., 431 Baring, Captain Hon. E., 332, 413 Barker, W. Burchart, 384 Barnard, J. H., 7, 13 Barnardiston, G., 368 Barnes, Dr. Fancourt, 19 Barnum and Bailey, Messrs., 447 Barry, W., 279 Barttelot, Major E. M. (the late), 350 Bate, Thomas, 11, 59, 60, 327, 367 Bates, Sir E. P., 62 Bayley, Sir Steuart, K.C.S.I., 443 Beavan, Captain R. C. (the late), 71, 323 Beck, G., 412 Beddington, Claude, 135, 148, 169, 171, 284, 298 Beddington, H. M., 135, 148, 169, 171, 284, 298 Bedford, Duke of, 22, 34, 35, 38, 41, 45, 47, 48, 49, 50, 52, 53, 55, 57, 63, 69, 84, 87, 90, 99, 204, 277, 323 Beech, Reginald, 38, 213, 221, 362, 373 Beit, A., 118, 132, 135, 147, 171, 187, 191, 216, 262, 267, 271, 435, 439 Beit, Otto, 18 Belcher, Captain, 67 Benett-Stanford, J., 7, 19, 114, 233, 248, 298, 310, 334, 344, 346, 408, 421, 437 Bengough, C. F., 105 Bennett, S. B., 475 Benson, Lieut. R. E. R., R.N., 475 Beresford, Captain Hon. J. G., 344 Berkeley, E. J. L., 142, 208, 242, 245, 393 Berlin Museum, 107, 115, 128, 181, 182, 262, 291, 313, 385, 435, 440 Berners, Captain Ralph, R.N., 428 Besant, Major W. H., 250, 350 Bessborough, Earl of, 59 Beswick, R. G., 454 Bethnal Green Museum, 262, 411, 419, 443, 449, 479 Betton, C. Steuart, 116, 208, 246, 276 Bevan, Owen, 318 Biddulph, Captain H. M., 204, 213, 218, 337, 345, 378, 408, 477 Biddulph, Colonel J., 41, 63, 64, 79, 218, 223, 229, 324, 325, 332, 335, 341, 343, 344, 346, 352, 362, 370, 372, 381, 383, 384, 385, 409, 411 Biddulph, General Sir R., 385 Bigge, Captain G. O., 74, 445 Bignold, A., 463 Bikanir, Maharaja of, 204, 229, 230, 281, 424, 465

Bird, S. G., 332 Blackwood, Lord Basil, 242 Blane, Major C. F., 47, 113, 156, 301, 329, 372, 476, 477 Blanford, W. T., 42, 49, 148, 227, 238, 321, 467, 475 Blewitt, R., 62 Bloemfontein Museum, 145, 199 Blois-Johnson, G., 341, 382 Blood, Major-General Sir B., K.C.B., 203 Blundell, Colonel H. B. H., C.B., 440 Blyth, Edward (the late), 343 Boas, H. J., 463 Boden, A. D., 418 Bodkin, J., 11 Bombay Natural History Society's Museum, 37, 62, 71, 73, 154, 204, 218, 304, 325, 411, 423, 462, 463 Bombay Natural History Society's Museum, Proceedings of, 411, 462 Bombelles, Count, 28 Bonham, G. L., 161, 176, 195, 208, 285, 393 Bonthron, Barclay, 47, 367 Booth, Sir Henry Gore, Bart., 84, 85 Bosanquet, O. V., 461, 462 Boucher, Captain B. H., 64, 73, 76, 82, 154, 318, 409, 462, 477 Bourke, Hon. Mrs., 20 Bower, Captain H., 372, 373 Bowie, Colonel M. M., 64, 400, 413 Bowker, A., 283 Bowker, M. E., 191 Bowker, W. Russell, 169, 261, 289, 428 Bowring, C. C., 122, 140, 147, 189, 194, 209, 263, 267, 268, 283, 298, 301, 311, 394, 429 Brackley, Lord, 135, 147, 264, 298 Braganza, H.R.H. D. Miguel, Duke of, 15. 25, 26, 315 Braithwaite, J. S., 10 Branch, C. C., 85, 462, 479 Brandon, A. J., 175, 195, 199, 300, 394 Brassey, Albert, 24 Brassey, Hon. T. A., 17, 19, 44 Breadalbane, Marquis of, 19 Breitmeyer, L., 19 Bridgeman, Colonel Hon. Francis C., 332 Brinkley, Captain J. T., 273, 349 British Museum, 1, 2, 3, 5, 7, 10, 11, 13, 28, 33, 35, 37, 38, 41, 42, 45, 51, 52, 53, 57, 58, 59, 62, 63, 65, 66, 67, 68, 69, 71, 73, 74, 75, 76, 77, 78, 79, 82, 83, 85, 87, 90, 91, 92, 97, 98, 99, 100, 106, 108, 109, 116,

Bindloss, W. R., 38, 113, 249, 429, 475

118, 122, 123, 126, 130, 132 134, 135, 145, 147, 148, 151, 152, 154, 155, 158, 161, 162, 163, 164, 165, 166, 169, 171, 174, 176, 178, 179, 181, 182, 184, 185, 187, 188, 189, 191, 193, 196, 197, 203, 204, 205, 207, 208, 210, 211, 213, 215, 216, 218, 219, 221, 225, 227, 229, 230, 238, 239, 240, 242, 246, 251, 253, 257, 262, 264, 267, 268, 271, 276, 277, 279, 281, 283, 285, 286, 287, 289, 291, 293, 294, 301, 302, 304, 305, 310, 313, 315, 316, 317, 318, 319, 321, 322, 323, 325, 327, 329, 332, 337, 339, 341, 344, 346, 347, 350, 352, 353, 355, 357, 359, 361, 362, 363, 364, 366, 368, 370, 372, 374, 376, 378, 379, 382, 383, 384, 385, 386, 387, 388, 390, 392, 393, 395, 396, 397, 399, 400, 401, 402, 404, 406, 407, 408, 409, 412, 413, 415, 418, 419, 431, 432, 436, 437, 439, 440, 444, 450, 476, 479 Brodie, A. W. M., 108, 163, 180, 267, 268, 397 Brogden, Captain J. S., 183 Brooke, A. Basil, 17, 57 Brooke, Sir Douglas, Bart., 20, 23 Brooke, E. Howard, 378 Brooke, E. J., 445 Brooke, E. W. S., 450, 457 Brooke, Captain Harry V., 82, 203, 228, 229, 321, 335, 383 Brooke, H. Brinsley, 85, 316, 355, 387 Brooke, Sir R. M., Bart., 23 Brooke, Sir Victor, Bart. (the late), 5, 7, 11, 12, 23, 37, 48, 52, 56, 57, 60, 63, 69, 71, 73, 74, 76, 77, 79, 82, 85, 92, 96, 97, 99, 105, 109, 116, 118, 130, 132, 134, 148, 155, 164, 174, 182, 205, 210, 215, 225, 227, 229, 230, 238, 239, 240, 244, 250, 256, 263, 270, 281, 284, 287, 289, 291, 294, 298, 313, 316, 322, 329, 332, 335, 337, 341, 343, 344, 353, 355, 359, 361, 367, 374, 379, 381, 382, 383, 384, 385, 387, 388, 393, 397, 402, 412, 419, 443, 479 Browne, Sylvester, 5, 10, 45 Bruce, C. W. A., 71, 412, 417, 418 Bryden, H. A., 113, 118, 139, 171, 215, 271, 289, 301, 311 Buchanan, Surg.-Major J. B., 204 Buck, W. J., 24, 355 Buckler, J. R., 289 Buckley, T. E., 120, 132, 135, 139, 147, 169, 175, 176, 187, 189, 194, 199, 208, 213, 242, 257, 268, 284, 368, 436, 454, 456 Buffon, 183 Burchell, Dr. (the late), 191, 197, 207 Burdon, J. A., 291

Burnett, J., 99 Burns, W. S. M., 18 Burrow, C. G., 23 Burton, Lord, 16, 17 Burton, Major A. (the late), 38 Butter, A. E., 327, 368 Büttikofer, Dr. J., 152 Buxton, E. N., 13, 25, 28, 45, 46, 225, 226, 273, 300, 315, 326, 348, 350, 352, 355, 386, 387 Buxton, Gerald, 366, 367 Buxton, H. G., 205, 281 Caccia, A. M., 64, 76, 154 Cadell, A., 329 Cadell, Lewis J., 7 Caine, L. E., 246 Cairns, Earl, 64, 401 Calcutta Museum (see Indian Museum) Caldwell, C. H. H. B., 465 Caley, Sir Edward, 11 Cambridge Museum, 479 Camden, Marquis, 19, 401 Cameron, A., 134, 148, 162, 166, 176, 194, 199, 299, 394 Cameron, E. S., 93 Campbell, A. C., 130 Campbell, A. S., 432 Campbell, Captain George, 113, 213, 232, 235, 249, 274, 301, 361, 455, 456 Campbell, Colonel J. E., 443 Campbell, Dr., 41 Campbell of Kilberry, J., 343, 347, 360 Campbell, J. McI. M'Iver, 94, 95, 105 Campbell, Hon. K., 123 Campbell, V. L. A., R. N., 6 Campbell, W. Y., 270, 300 Campbell-Orde, Sir A., Bart. (the late), 399 Cape Town Museum, 118, 123, 130, 131, 139, 144, 147, 149, 151, 159, 161, 167, 169, 172, 174, 191, 194, 208, 215, 262, 266, 271, 285, 289, 297, 311, 440, 476 Capell, A. E., 285 Cappel, E. L., 256 Carden, J. R., 346, 362 Carew, Capt. B. L., 248, 274, 302, 361, 383 Carew, Major G. A. L., D.S.O., 261, 344, 378, 379, 413, 461, 462, 465 Carey, -, 408 Carr, Lieut.-Col. E. E., 204 Carr-Glyn, Geoffrey, 20 Carrington, Colonel E. (the late), 451 Carrington, Major-General, Sir F. C., K.C.B., K.C.M.G., 194

Carroll, J. W., 107, 184, 197, 201, 267 Carroll, R. P., 141, 197, 435, 445 Carter, H. R. P., 423 Carter, Rev. V. R., 440 Cathcart, Sir Reginald, Bart., 423 Cator, R., 275 Cavendish, H. S. H., 171, 200, 298, 450, 45 I Cetywayo (the late), 450 Chalmers, Guthrie and Co. (Messrs.), 450 Chamberlain, Col. Neville, 423 Chambers, Major O. A., 229 Chanler, W. Astor, 116, 142, 200, 207, 242, 306, 392, 436, 450, 452 Chapman, Abel, 2, 4, 7, 12, 17, 24, 354, 355 Chapman, G. D. E., 172, 272 Charrington, F., 392 Charrington, Henry, 142, 207, 242, 245, 276, 436 Charteris, Hon. A., 20 Cheetham, G. H., 112, 247, 248, 274, 302, 304 Chelmsford, General Lord, G.C.B., 450 Chillingworth, S., 262, 267, 298, 310, 428 Chisholme, Colonel Scott, 74 Cholmondeley, V., 47, 363 Christian, E. G., 285 Church, P., 213, 223, 339, 362, 373, 408, 412 Clapperton, Captain (the late), 126, 396, 397 Claraz, G., 419 Clarke, T. W. H., 47, 48, 57, 95, 105, 113, 156, 157, 171, 232, 233, 235, 248, 249, 252, 253, 255, 273, 300, 304, 327, 366, 476 Clifton, J. Talbot, 5, 18, 19, 390 Clive, P. A., 107, 180, 397 Cloete, Montrose, 20, 195 Clough, J. W., 79 Coats, A., 479 Cobb, W. H., 74, 232, 253, 257, 300, 304 Cobbold, J. D., 44, 327, 345, 350, 358, 367, 381, 387 Cobbold, P. W., 38, 339 Cobbold, R. P., 346, 372, 373 Coburn, F., 313 Codrington, Captain E. W. (the late), 38, 341, Codrington, Captain H. W., 38, 203, 218, 229, 318, 335, 341, 344, 361, 362, 379, 382, 383 Codrington, W. R., 346 Coke, Lieut.-Col. Hon. W., 5, 13, 45, 47, 105, 108, 115, 139, 162, 169, 175, 179, 181, 194, 199, 239, 250, 266, 275, 284, 289, 300, 302, 332, 367, 395, 435, 440 Colby, J. V., 6, 94, 326

Collett, General Sir Henry, K.C.B., 79, 341 Collins, A. H., 78, 79 Colmore, Q. C., 5 Colville, Major A., 281 Compton, Captain Lord Douglas, 131, 175, 393, 412 Conway, Sir W. Martin, 7 Cookson, Captain F., 18, 19, 92, 94, 105, 130, 169, 209, 215, 216, 271, 302, 311, 366 Cookson, W. R., 19, 105 Cooper, Frank, 44, 368 Cooper, R. A., 147, 167, 172, 175, 187, 194, 216, 263, 271, 284, 293, 301, 310, 311, 394, 419 Coote, Eyre, 399 Copeman, Capt. H. C., 318 Coppinger, A. J., 204, 229 Coryndon, R. T., 122, 148, 161, 187, 189, 194, 208, 260, 261, 294, 311, 439 Cottingham, H. L., 412, 444 Cotton, B., 250 Cotton, Major J. W. M., 74, 204, 382, 414, 460, 463 Coudenhove, Count Henry, 98, 99, 469 Cox, Captain P. Z., 259, 411 Crake, D. H., 94 Crispin, G. C. W., R.N., 6 Crosley, W., 176 Crossman, Major-General Sir Wm., K.C.M.G., 435 Crum, A. S., 318, 361 Culme-Seymour, Admiral Sir Michael, Bart., Cumberland, Major C. S., 11, 38, 47, 64, 74, 76, 82, 213, 221, 339, 346, 352, 359, 361, 372, 373, 376, 378, 382, 383, 384, 409, 413 Cumine, A., 63, 230, 412 Cuninghame, Captain G. R., 249 Cuninghame, R. J., 300, 428 Cuppage, Colonel (the late), 334 Curtis, W. S., 130, 216 Cust, Lieut.-Col. M., 64, 73 Cuthbert, Captain G. J., 226, 364 Czetwertynski, Prince Boris, 171, 235, 249, 272, 289, 392, 429, 436 Dalgety, Captain H. B., 462 Dalgleish, --, 408 Dalgleish, Sir W. Ogilvie, 17 Dallas, A. G., 390 Dalton, Captain E. G., 325 Dalzell, Major Hon. A., 378

Danford, C. G., 33, 315, 353, 384

Darley, J., 44

Darling, Jff., 161, 169 Darrah, H. Z., 38, 212, 213, 218, 335, 342, 345, 408 Dartmouth, Earl of, 261, 270, 301, 435 Darwin, Charles (the late), 99 Dashwood, General R. L., 5, 6, 9, 10 Dauvergne, H., 373 David, Père A., 58, 90, 219, 319, 326 Davies, Digby, 113, 249, 257, 273, 300, 306, 450 Davies, H. E. M., 62 Davies, H. Ll., 418 Davis, A. P., 321 Davis, A. W., 175, 215 Deasy, Captain H. H. P., 359, 409 de Bardi, Henri de Bourbon, Comte, 219, 379 de Barreto, H., 460, 462 de Brazza, S., 293 de Burgh, Hugo, 329 de Bylands, Count J., 113, 114, 233, 248, 253, 256, 274, 305, 437 Decatur, S., 11 Decle, Lionel, 142 Deeble, Surgeon-Major B. W. C., 82 Deeds, W. G., 445 Delamere, Lord, 13, 111, 114, 116, 120, 141, 142, 157, 169, 171, 176, 177, 179, 200, 207, 232, 235, 242, 244, 245, 253, 256, 258, 259, 273, 274, 301, 302, 304, 311, 393, 449, 450, 455, 456, 457 de Lucinge, Prince, 109, 197, 250, 286, 301, de Lucinge, Prince A., 113, 148, 162, 175 Demidoff, Prince, 12, 13, 33, 34, 47, 219, 316, 346, 353, 357, 359, 376, 384, 423, 467 Demidoff, Princess, 376 de Murietta, F., 364 Denham, Colonel (the late), 126, 396, 397, 419 Denman, G. L., 24 Denny, E. M., 24 Denny, P. R., 249, 302, 450, 451 de Nolde, Baron (the late), 460 de Paris, H.R.H. la Comtesse, 22 de Poncins, Viscount Edmond, 64, 113, 114, 148, 157, 162, 172, 176, 196, 204, 225, 226, 229, 232, 235, 237, 248, 249, 257, 258, 259, 273, 274, 281, 286, 303, 304, 316, 334, 346, 364, 372, 374, 423, 424, 428, 429, 448, 449, 451, 462, 466, 468, 472, 477 de Pousargues, E., 35, 50 De Prée, H., 346, 400, 413, 414 Derby, Earl of (the late), 126, 184, 267, 285, 313, 397, 419 de Trafford, Sir Humphrey, Bart., 16, 44

Devas, F., 17 De Weese, Dall, 10 de Winton, F., 101, 102 Dibble, H., 424 Didham, Captain Chambers, 74, 76 Dingwall, K., 381 Dolby, G. A., 431, 445 Donaldson-Hudson, Mrs., 59 d'Orléans, H.R.H. le Duc, 19, 22, 29, 34, 41, 48, 85, 87, 114, 213, 232, 250, 254, 255, 256, 298, 305, 313, 359, 374, 388, 406, 423, 440, 463 d'Orléans, Prince Henri, 218, 221, 223 Douglas, Major George, 113, 205, 274, 327, 343, 345, 382 Down, C. P., 229 Dowson, Captain C. Hutton, 62 Dowson, Staff-Surgeon J., R.N., 98, 123, 176, 289, 393, 421 Drayson, General A. W., 283 Drewitt, Dr. Dawtrey, 237 Drought, J. A. H., 366 Drury, Surgeon-Major M. O'C., 475 Dublin Museum (see Science and Art Museum) Du Cane, A., 85 Du Chaillu, P., 287 Dudley, Countess of, 18 Dufferin and Ava, Marquis of, 281 Dunbar, J. Brander, 134, 157, 171, 194, 232, 235, 248, 253, 285, 300, 306 Dunell, O. R., 122, 131, 147, 149, 151, 161, 165, 169, 174, 187, 191, 194, 199, 208, 215, 263, 266, 271, 283, 288, 293, 297, 310 Dunmore, Earl of, 19, 33, 45, 47, 174, 262, 267, 289, 299, 346, 373, 394, 421, 431, 479 Dunsford, F. B., 263, 298 Durand, Sir E. L., Bart., C.B., 460, 466 Durban Museum, 176, 289, 298, 437

Ebden, Alfred, 215, 216, 293
Ebrington, Viscount, 21
Eden, Sir Ashley, 41
Edinburgh Museum (see Science and Art Museum)
Edmonds, W. A., 194
Edye, Surgeon-Major J. S., 248
Egerton, Captain A., 326
Egerton, C. F., 64, 76, 154, 413, 475
Elliot, Sir C. A., Bart., 431
Ellis, Major-General Sir Arthur, K.C.V.O., 26, 27, 62, 63, 73, 132, 176, 284, 300, 316, 373, 412

East India Club, 334, 349, 381

Ezra, E., 344

Ellis, Hon. Charles, 11, 38, 44, 45, 60, 175, 213, 284, 289, 300, 321, 339, 343, 346, 373, 379, 382, 421, 479 Ellis, Major C. C., 5, 44, 249, 274, 301, 304, 316, 366, 372 Ellis, F. H. B., 367 Elphinstone, Lord, 413 Elwes, H. J., 7, 12, 29, 34, 41, 49, 50, 60, 87, 344, 376 Ennismore, Viscount, 176, 194, 208, 394 Erbach, Count, 26 Erbach-Erbach, Count, 30 Erdödy, Count George, 27 Erdödy, Count Rudolf, 27 Esterhazy, Count Michael, 27 Eustace, C. F., 284 Evans, Vet.-Capt. G. H., 66, 70, 71, 73, 80, 82, 230, 323, 412, 417, 418 Evans, Colonel J., 61, 63, 154, 229 Everett, A. H., 67 Eyre, A., 194 Eyre, H. H. (the late), 171 Ezra, A., 475, 477

Fairholme, Major W. E., 413 Falconer, Dr. H. (the late), 37, 62 Fannin, J. T. (see Victoria Museum) Farquhar, Ernest, 11, 44, 94 Fawcett, Major J. M., 281 Fell, Captain R. B., 154, 465, 466 Fenton, Lieut.-Col. L. L., 63, 73, 82, 154, 204, 230, 281, 322, 352, 384, 411, 412, 414, 423, 455, 461, 462, 463, 464, 465, 467, 471 Fenton, Major W. D. B. (the late), 38, 345 Fenwick, Noel, 316, 400, 460 Ferguson, Captain A. G., 466 Ferguson, Major H. J., 46 Ferrand, Captain Gerard, 7, 11, 476 Ferrers-Stranack, R. H., 189, 209, 294 Festetics, Count Tassilo, 27, 29 Festing, Captain A. H., 107, 108, 163, 183, 201, 397 Ffinch, B. T., 276 Field, The, 24, 423 Fielden, —, 261 Fielden, Colonel W. H., 390 Fife, Duke of, K.T., 20 Fincastle, Viscount (V.C.), 346, 374 Findlay, F. R. N., 131, 161, 169, 193, 199 Findlay, M. le C., 33 Finnie, Captain Robert, 423 Finsch, Dr. O., 221

Firman, H. B., 431 Fitzgerald, Captain G. J., 48, 94, 105, 230, 413 Fitzgerald, G. P., 367 Fitzgerald, Lord Walter, 382 Fitzherbert, J. W., 440 Fitzpatrick, J. P., 261 Flanagan, Lieut.-Colonel J. W. H., 248, 256, 274, 305, 412, 456 Fletcher, E. L., 18 Fletcher, W. A. L., 17, 87 Flower, Lewis, 372 Forbes, A., 399 Forbes, A. J., 271, 302 Forbes, Major P. W., 123, 195, 266, 297, 310, 421 Fordham, W. A., 48 Forsyth, Sir Douglas (the late), 49, 372 Forsyth, Captain J. (the late), 73, 410 Fortescue, Earl, 21 Fortune, J., 444 Foster, J. Kenneth, 48, 85, 113, 156, 249, 256, 302, 305, 436 Foster, Major Kingsley, 38 Fountaine, A. C., 262, 266, 271 Fowler, Sir John, Bart., 17 Fowler, J. K., 18 Fowler, Sir Thomas, Bart., 175, 301 Franklin, Captain Sir J. (the late), 5 Fraser, Captain Hugh, 105, 460, 462 Fraser, Captain Sir K., Bart., 263, 311, 456, 461, 465 Fraser, Major R. H., 413 Frere, Sir Bartle, Bart., 435 Frewen, Moreton, 46, 48, 92, 94, 367, 406 Fulke-Greville, Hon. C. B., 344 Fuller, F. C., 267 Fuller, Captain John, 401, 413 Fünfkirchen, Count Henry, 28 Gage, M. F., 185

Gage, M. F., 165
Gedge, E., 116, 119, 128, 142, 169, 171, 175, 179, 185, 187, 207, 242, 245, 276, 294, 304, 395, 421, 436, 456
George, F. Gordon, 479
German Emperor, H.I.M. the, 29
Ghika, Prince Demeter, 27, 177, 305, 306, 437, 452
Ghika, Prince Nicolas, 29, 114, 248, 273, 301, 437
Gibbons, Major A. St. H., 169, 189, 193, 266, 297, 454, 456
Gilbert, R., 62
Gill, Captain T. W., 17

Gilliat, J. B., 46, 105 Gillman, W., 71, 465 Ging, Captain, 153 Gleichen, Captain A. W. E. Count, 17 Glossop, B. R. M., 73, 113, 232, 233, 248, 253, 268, 274, 301, 318, 322, 428, 468 Glyn, Sir Richard, Bart., 392 Glynn, A. H., 138, 147, 150, 161, 174, 191, 194, 199, 261, 264, 284, 289, 297, 456 Glynn, H. T., 138, 147, 150, 161, 165, 174, 191, 194, 199, 261, 264, 284, 289, 297, 456 Goddard, H. C., 268, 313 Goff, T. C. E., 46 Goldingham, J. D., 411, 443 Goldschmidt, G. A., 315 Goold-Adams, Major H. J., C.B., C.M.G., 122, 149, 151, 166, 167, 169, 187, 189, 262, 266, 270, 271, 283, 289, 293, 294, 297, 310, 311 Gordon, Colonel Evans, 460, 463 Gordon, Captain E. R., 281 Gordon, Captain P. J., 221, 229 Gordon, Colonel T. E., 372 Gordon, Captain W. E., 346 Gordon-Cumming, Colonel, 16, 17, 84, 465 Gordon-Cumming, Roualeyn (the late), 16, 131, 266, 270, 438, 447 Gordon-Cumming, Colonel W. G., 438, 439 Gore, Sir G., Bart., 46 Gore-Graham, W. P., 350 Gossage, W. W., 18 Gough, J. E., 123, 140, 161, 177, 194, 209, 264, 267, 285, 311, 394 Gould, George H., 366 Gould, J., 316 Gouldsmith, C. C., 123, 138, 175, 195, 393 Grace, M. P., 46 Graeme, G. Lloyd, 327 Graham, Lieut.-Colonel, 325 Graham, Captain M. D., 298, 310 Graham, P. N., 6, 11 Grahamstown Museum, 207, 283 Grant, Colonel (the late), 341 Grant, Sir Arthur, Bart., 17, 84 Grant, A. J., 221, 340, 341 Grant, C. Macpherson, 84 Grant, E., 44 Grant, Sir G. Macpherson, Bart., 84 Grant, Major James, 63, 73, 85, 92, 169, 176, 263, 298, 311, 343, 390, 393, 411 Grant, Captain Johnstone (the late), 16 Grantley, Lord, 237 Graves, A. B., 204 Gray, Dr. G. R., 37

Greaves-Bagshawe, M. C., 263, 267, 310 Green, E. H. E., 465 Greenaway, Lieut.-Colonel T., 218, 378 Greenfield, T. W., 113, 245, 249, 253, 274, 275, 302, 305, 332, 345, 373, 409, 411, 449 Greenway, Major, 377 Grenfell, C. A., 17 Grenfell, Captain H. H., R.N., 6 Greville, Hon. C., 263, 298, 311 Grieve, J. W., 431 Grinnell, G. B., 92 Grogan, E. S., 195 Grogan, Sir E. Ion, Bart., 339 Grudzinski, Count, 254, 436 Guides, Officers' Mess, Q.O. Corps of, 38, 329, 335, 341, 343, 344, 382 Guillemard, Dr. F. H. II., 144, 370, 426, 476 Guinness, Major E., 343 Gunnis, F. G., 232, 235, 256, 302, 306 Günther, Dr. A., 241 Gwynne, Captain J. H., 63, 461 Hagenbeck, Carl, 87, 310, 343, 352, 435 Haggard, J. G., 163 Hale, Surgeon-Major G. E., D.S.O., 64, 74, 401 Hall, A. C., 345 Hallowes, Major R., 6 Hamilton, Duke of, 390 Hamilton, Marquis of, 52, 135, 139, 311, 393 Hamilton, A. S., 122 Hamilton, General Douglas (the late), 411, Hanbury, David T., 5, 39, 218, 223, 346, 371, 373, 374, 379, 390 Hanbury, J. M., 47 Hancock, Captain W. W., 76 Handyside, J. S., 31 Hankey, Lieut.-General B., 44 Hanson, Lieut.-Colonel G. W., 444 Harden, Major F. J., 468 Harden, Major J., 345 Hardinge, Hon. C., C.B., 353 Hardwicke, General (the late), 71, 402, 412 Hardy, Hon. Gathorne, 48 Hardy, Gerald, 367 Hare, E. P., 272, 304 Hargreave, John, 17 Harkness, Major T. R., 304, 455 Harris, Lord, G.C.S.I., 454, 457

Harris, Frank, 139, 175, 289, 294, 297

Harris, Sir W. Cornwallis (the late), 103

Harris, L. T., 399

Harrison, Captain Edgar G., 434 Harrison, James J., 6, 11, 44, 45, 73, 92, 94, 123, 134, 139, 140, 147, 150, 169, 175, 176, 194, 204, 208, 263, 270, 284, 300, 311, 327, 393, 421, 428, 431, 436, 443, 445, 447, 448, 451, 452, 454, 455, 460, 461, 465, 466, 469, 476, 477 Hart, W. W., 10, 390 Harting, J. E., 85 Harvey, Colonel C. B., 7 Harvey, Sir Robert, Bart., 38, 62, 63, 76, 115, 125, 128, 141, 148, 154, 158, 169, 177, 196, 197, 208, 213, 230, 242, 244, 275, 276, 285, 305, 329, 339, 344, 362, 379, 381, 392, 408, 436 Hastings-Wood, C., 230 Hatch, G. W., 73 Hawker, R. McD., 113, 232, 235, 248, 253, 273, 301, 304, 305 Hay, Lord A., 33 Hay, A. R., 64, 74, 401 Hayes, Major C. H., 339, 345 Hayes-Sadler, Major R., 118, 138, 271, 298, Hayes-Sadler, Captain W., 335 Heathcoat-Amory, Sir John, Bart., 21 Heber-Percy, Major Algernon, 368, 404 Heber-Percy, H. L., 401, 466 Heber-Percy, Colonel R. J., 82, 205, 213, 230, 322, 329, 345, 355, 379 Hemming, Sir A. W. L., K.C.M.G., 469 Henderson, A., 24 Henniker, Major Hon. A. H., 19 Henry, E. R., 460 Henry, Captain G. F., 131, 134, 176, 195 Herron, G. O. M., 57 Hesketh, Sir Thomas, Bart., 479 Heuglin, —, 181 Hibbert, Captain E. H. R., 64, 281 Hicks, F. R., 179 Hicks-Beach, Captain A., 204, 229, 318, 329, 413 Higgs, Dr. (the late), 108 Hill, Sir Clement, Bart., 28, 316, 421 Hill, Captain G. W., R.N., 390 Hinde, Colonel J., 463 Hinde, S. L., 116, 141, 242, 245, 268, 421 Hoare, Captain H., 465 Hobley, C. W., 115, 207 Hodgson, B. H. (the late), 41, 63, 71, 76, 79, 82, 218, 322, 325, 360, 362, 378, 379, 400, 409, 419, 468 Hogg, Major, 353

Holden, E. F., 345 Holden, H. R., 123, 139, 176, 194, 284, 392 Holdsworth, Captain G. L., 203, 461, 462, 465 Holland, H. C., 431, 445 Holland, W., 460, 463 Hollinek, Eberhard, 315 Holmwood, F. (the late), 435 Holt, P. A., 357 Holub, Dr. Emil, 187 Hopley, Mr. Justice, 108, 120, 122, 128, 130, 132, 138, 144, 147, 149, 151, 154, 161, 166, 167, 169, 171, 174, 184, 187, 189, 191, 194, 199, 208, 237, 242, 248, 251, 253, 256, 261, 266, 271, 284, 293, 297, 310, 311, 393, 440 Hornaday, W. T., 405 Hornby, Colonel J. F., 21 Hosack, D. D. F., 445 Hoselton, Colonel, 10 Howard, Colonel, 315, 466 Hoyos, Count E., 46, 95, 105, 131, 134, 139, 144, 162, 169, 174, 175, 177, 195, 199, 203, 209, 216, 225, 232, 235, 250, 256, 262, 266, 274, 285, 289, 300, 305, 311, 327, 347, 367, 393, 406, 429, 436, 448, 451, 455, 461, 465, 466, 477 Hoyos, Count Joseph, 25 Hoyos, Count Jose, 28 Hoyos, Count Max, 26 Hudson, W. O., 415 Hughes, R., 175, 393 Hulke, Captain L. I. B., 229, 318, 346 Humbert, A. C., 130, 161 Hume, A. O., C.B., 37, 38, 39, 40, 41, 49, 62, 63, 66, 70, 71, 73, 76, 79, 81, 82, 100, 202, 203, 204, 205, 213, 218, 220, 221, 223, 229, 230, 272, 281, 318, 320, 321, 323, 325, 328, 329, 332, 334, 335, 336, 337, 338, 341, 343, 344, 351, 352, 361, 372, 373, 378, 379, 381, 382, 384, 388, 399, 400, 407, 408, 411, 412, 414, 415, 431, 443 Hunt, Commander A. T., 176, 195, 301, 310, 350, 393, 421 Hunter, H. C. V., 64, 71, 76, 115, 124, 125, 127, 128, 141, 158, 177, 197, 207, 213, 218, 229, 242, 244, 245, 257, 275, 276, 284, 305, 372, 381, 383, 392, 408, 417, 418, 435, 452 Hunter, Captain W. H., 462, 463 Hutchinson, Lieut.-Col. R. E., 343 Hutchison, J. R., 17, 18, 19 Hutton, Colonel E. T. H., 203 Ilchester, Earl of, 22

Illingworth, P. H., 216 Indian Museum, Calcutta, 41, 42, 62, 71, 73, 76, 77, 82, 154, 213, 218, 230, 281, 318, 323, 325, 329, 332, 343, 352, 361, 372, 379, 411, 415, 417 Inglefield, Commander G. F., 21 Innes, Mrs., 465 Inverarity, J. D., 62, 63, 73, 74, 76, 153, 154, 213, 248, 253, 257, 306, 332, 352, 383, 399, 411, 412, 413, 428, 429, 456, 461, 463 Italy, H. M. the King of, 347, 374 Italy, H.M. the late Victor Emanuel, King of, 347 Ivrea, Marquis of, 315, 353 Jackson, F. J., C.B., 115, 119, 120, 122, 128, 141, 148, 157, 158, 163, 172, 176, 185, 197, 200, 207, 208, 241, 242, 244, 246, 257, 272, 274, 275, 276, 284, 285, 310, 392, 393, 436, 448, 450, 451 Jackson-Harmsworth Polar Expedition, 475 James, F. L. (the late), 395 James, H. A., 94 James, H. E. M., 87 James, Walter, 326 James, W. D., 7, 109, 110, 114, 177, 233, 249, 250, 268, 274, 286, 301, 304, 395, 421 Jameson, J. A., 139, 262, 266, 300, 367 Jameson, J. S. (the late), 118, 187, 193, 262, 270, 405, 440, 451 Jankovich, G., 28

Jay, P., 63 Jayakar, Surg.-Lieut.-Col. A. S. G., 330 Jenkins, S. E. F., 80 Jenkinson, Major G. S. C., 180, 397 Jeppe, Carl, 184 Jeppe, Julius, 106, 107, 108, 109, 110, 116, 118, 120, 122, 123, 126, 128, 130, 131, 132, 134, 138, 140, 144, 145, 147, 148, 149, 150, 157, 161, 166, 167, 169, 171, 174, 175, 179, 180, 184, 185, 187, 189, 191, 193, 195, 197, 199, 201, 216, 225, 233, 235, 237, 242, 245, 249, 253, 263, 266, 271, 274, 279, 283, 284, 289, 293, 297, 302, 306, 310, 316, 394, 397, 428, 436, 440, 450, 451, 467 Jerdon, Dr. T. C. (the late), 431 Jhalawar, Maharaj Rana Bahadur of, 153, 228 Johnson, C. R., 335, 339 Johnston, Sir H. H., K.C.B., 137, 279 Johnston-Stewart, J., 82, 100, 113, 157, 204, 232, 235, 249, 253, 256, 257, 273, 297, 318,

381, 454, 455, 466, 471

Johnstone, Mrs. Lawrie, 463

Jones, C. T., 130, 132, 135, 147, 149, 151, 161, 166, 167, 169, 175, 187, 191, 194, 199, 216, 263, 271, 284, 285, 289, 293, 297, 300, 302 Jones, Heighway, 62 Jones, Captain Lewis, 74 Jones, Walter, 353 Keglevich, Count B., 28 Keith-Falconer, V. F. A., 345 Kelham, Major H. R., 345 Kemp, H. F., 24 Kempson, T. P., 6, 92, 94, 95, 327, 367 Kennard, C. H., 367 Kennard, Martyn, 38, 332, 335, 339, 343, 373, 381, 412 Kennedy, Lord David, 193 Kennedy, Vice-Admiral Sir William, 5, 69, 97, Keppel, Commander C., C.B., 69 Keppel, G. Roos, 73, 341 Keys, H. W., 412 King, Colonel, 37 King, Captain A., 350 King, Major A. C., 465 King, H. S., 71 Kinloch, Major-General Alexander A. A., 71, 74, 203, 213, 218, 229, 329, 334, 336, 341, 344, 361, 379, 383, 399, 408, 412, 443, 444 Kirby, F. Vaughan, 103, 122, 123, 132, 134, 135, 138, 139, 147, 149, 150, 151, 161, 165, 167, 169, 171, 174, 191, 193, 199, 207, 261, 262, 264, 266, 283, 289, 297, 310, 392, 394, 420, 421, 425, 428, 429, 431, 435, 447, 454, 455, 456, 457, 466, 467, 471, 472 Kirk, Sir John, K.C.B., 116, 122, 128, 158, 164, 196, 197, 283, 300, 304, 305, 394, 421, 435, 448, 450 Kirwan, Major Maitland, 45, 47, 94, 367 Knight-Bruce, J. C. L., 10 Kraus, Dr. A., 151 Kuch-Behar, Maharaja of, 372, 398, 399, 400, 411, 412, 431, 459, 460, 461, 462, 463, 465, 466, 477 Kund, Jeno, 27

Johnstone, R., 218, 361, 379, 382

Labouchere, F. A., 315, 343 Lacy, Lieut. E., R.N., 239 Lambton, Captain Hon. W., 431 Lamont, J., 392, 421, 435, 456, 475 Lane, Captain H. P., 462 Lane, W. H., 362, 378 Langworthy, E., 346, 468 Lansdowne, Marquis of, K.G., K.C.M.G., 334, 343, 372 Larios, Pablo, 355 Lascelles, Hon. Gerald, 85 Law, S. C., 63 Lawley, Major Hon. R. T., 262 Lawrence, C., 19 Lawrence, W., 372, 373 Lawrence, W. R., 339, 378 Lawson, E. St. J., 203 Laycock, J. F., 18 Lean, F., 393 Learmonth, W. Livingstone, 99, 476, 478, Leatham, A. E., 48, 71, 73, 76, 94, 96, 249, 274, 327, 332, 355, 368, 378, 454, 457, 461, 462 Leather, Captain G. F. T., 175, 232, 233, 235, 257, 304 Le Champion, Loftus M., 228, 281 Lee, C. G. R., 368 Lee, Captain Frank, 461, 462 Lee, L. B., 48 Lehmann, Captain F. H., 123, 132, 139, 175, 195, 263, 271, 393 Leigh, J. Hamilton, 19 Le Mesurier, T., 203 Lennard, H., 46, 64, 105, 223, 335, 373, 400 Lennard, H. St., 74 Lethbridge, A. Y., 387 Levita, Captain Cecil, 318, 347 Liddell, C., 157, 232, 256, 301 Liebinger, P., 94, 405 Liechtenstein, Prince Henry of, 25, 26, 27, 47, 162, 177, 207, 238, 242, 245, 250, 267, 300, 393, 404, 406, 436, 454, 455, 470 Lilford, Lord (the late), 24, 385 Lindsay, A. B., 329 Lindsay, H. E., 401, 445 Lister-Kay, Colonel F. C., 74, 218, 337, 361, 381, 400, 408 Littledale, Edmund, 366 Littledale, St. George, 5, 32, 33, 45, 86, 87, 88, 104, 105, 204, 219, 223, 315, 316, 322, 327, 332, 339, 346, 356, 357, 361, 366, 372, 373, 374, 375, 376, 379, 382, 403, 404, 406, 408, 475, 477 Livingstone, David (the late), 186, 308 Livingstone, Captain H. A. A., 131, 132, 191 Llewelyn, Sir R. B., K.C.M.G., 313 Lloyd, Mrs. Graham, 60 Lloyd, Colonel G. E., D.S.O., 350

Lloyd, Surgeon-Major O. E. P., 78 Loam, M., 64, 332, 465, 466, 477 Loder, Sir Edmund G., Bart., 5, 10, 18, 26, 33, 35, 37, 41, 45, 52, 57, 58, 59, 63, 66, 67, 69, 71, 73, 76, 77, 78, 79, 85, 87, 90, 92, 94, 97, 99, 100, 105, 110, 113, 116, 118, 119, 126, 128, 131, 134, 139, 144, 152, 153, 155, 157, 162, 166, 171, 175, 180, 182, 183, 185, 187, 191, 194, 197, 201, 208, 210, 213, 215, 218, 221, 225, 226, 229, 232, 235, 237, 242, 245, 248, 250, 253, 256, 259, 262, 266, 267, 271, 273, 275, 276, 277, 279, 281, 284, 287, 289, 291, 298, 299, 304, 305, 311, 313, 315, 318, 322, 325, 327, 329, 332, 335, 341, 343, 345, 347, 350, 352, 353, 355, 357, 359, 361, 364, 366, 370, 372, 374, 376, 378, 379, 381, 384, 387, 394, 397, 399, 402, 406, 412, 418, 419, 421, 423, 440, 448, 450, 452, 456, 461, 463, 475 Lostus-Tottenham, W. F., 80, 413 Logan, Crawford G., 45, 105 Logan, E. T., 46, 48, 94, 105 Long, A. M., 323 Lonsdale, Earl of, 5, 11, 327, 390 Lort-Phillips, E., 19, 208, 232, 239, 257, 268, 300, 305, 395 Low, H. B., 66, 418 Lucas, C. T., 20, 22 Lucas, W. H., 22 Lucknow Museum, 204, 334, 345, 361, 374, 378, 381, 400, 408 Lugard, Capt. E. J., 215, 284 Lugard, Colonel F. D., C.B., D.S.O., 108, 116, 120, 142, 185, 196, 207, 242, 245, 424, 450, 452 Lumsden, D. M., 400, 444 Lydekker, R., 38, 218, 322, 339 Lyon, Captain, 476 Lyte, W. Maxwell, 19 MacAlpine, D., 161, 283, 297, 472 Macdonald, H. C., 297, 310, 392 Macdonald, Major J. R., 196 Macdonald, R. H., 335 M'Douall, Kenneth, 7 Macintosh, Mrs., 5 Mackay, W. Grant, 366 Mackenzie, A. T., 412 M'Kie, J., 424 M'Lachlan, W. J., 345, 382 Maclagan, Fergus, 288 Maclaine of Lochbuie, The, 94, 144, 298, 352,

366

M'Lean, H., 31 MacLean, H. L. S., 329 MacLeod, Captain J. N., 63 MacMaster, General, 332 M'Micking, Captain H., 412 M'Neill, Captain M., 73, 76, 85, 112, 229, 248, 274, 305, 412, 436, 437, 448, 451, 452, 455, 462, 475, 477 Madan, R. H., 62 Mainwaring, Lieut.-Col. H. G., 256 Makin, Charles, 11, 46 Malet, Major A. G. W., 76 Maloney, Dr., 215 Mann, T. H., 445 Manners, Lord Edward, 176, 195 Marchetti, G., 11 Marker, Captain R. J., 443, 444, 445 Markham, A. M., 460, 463, 467 Marriott, Captain John, 7, 84, 350, 352, 364, Marshall, E. T., 114, 274 Marshall, Lieut.-Col. L., 25 Marsham, Captain Hon. R. H., 461 Marston, Colonel F., 352 Martin, Colonel, 203 Martin, Colonel Cunliffe, 205 Massey, Major C. F., 382 Mathews, General Sir Lloyd William, 122, Mathie, Colonel J., 399 Matschie, Dr. P., 102 Maude, H., 38 Maxwell, Colonel C., 150 Maxwell, Captain J. M'Call, 95, 105, 113, 256, 273, 301, 450, 451 Meerut Tent Club, 423 Meldrum, James, 428 Melliss, Captain C. J., 457 Melville, S., 346 Menges, J., 109, 110, 112, 114, 179, 232, 233, 235, 237, 239, 248, 253, 256, 257, 258, 259, 267, 272, 274, 298, 304, 349, 395 Meran, Count John of, 315 Messel, L. C. R., 92 Methuen, Messrs., 137 Meux, Sir H. B., Bart., 45, 256, 388, 449, 450 Meyer, Dr. A. B., 30 Micklethwait, G., 162, 195, 284 Micklethwait, R. K., 18, 122, 135, 162, 195, 394 Mikhaelovitch, Grand Duke, 33 Millais, J. G., 17, 18, 19, 20, 22, 23, 56, 57, 84, 85, 174, 261, 262, 267, 297, 374

Millbank, P. C., 388 Milne, G. B., 213, 408, 409 Milner, Sir A., 191 Mitchell, F. J., 45, 311 Mitchell, Dr. R. P., 138 Mittrovszky, Count, 26 Mockler, Captain G. H., 79 Mockler-Ferryman, Major A. F., 183 Moir, D. F., R.N., 6 Möller, Lieut.-Colonel B. D., 461 Molyneux, Captain F., 6 Monck-Mason, R. H., 397 Moncreiffe, W., 43, 44, 47, 350, 386, 387, Moncrieff, R., 84 Montgomerie, Captain, R. A. J., C.B., R.N., 115, 128, 142, 208, 245, 276, 316, 392, 436, 454, 455, 456, 472 Moray-Brown, J. (the late), 424 Morgan, Rhodes, 332 Morland, Major H. C., 46, 73, 204, 213, 229, 232, 235, 249, 274, 306, 339, 361, 374, 378, 400 Morphy, Captain H. J., 465 Morris, Sir John, K.C.S.I., 62, 64 Morris, Colonel W. J., 62 Morse, I., 329, 339, 408 Moseley, A., 118, 134, 298 Mosse, Charles B., 163 Muir, J. Gardiner, 115, 171, 176, 208, 242, 244, 276 Mumford, T., 444 Murdoch, A. M., 463 Murphy, Captain M., 335, 339 Murray, A., 5 Murray, H., 388, 435 Murray, H. W., 413 Murray, M. (the late), 186 Murray, W. S., 73, 444 Muttra Tent Club, 465 Nádasdy, Count Fery, 26, 28 Napier, D. R., 322

Napier, D. R., 322
Nash, R. M., 62
Naval and Military Club, 38
Naylor, A. M., 64, 123, 139, 263, 282, 283, 297
Neilson, A., 122, 271, 298, 394
Nelder, C., 21
Nelson, H. C., 47
Neumann, A. H., 101, 103, 111, 127, 128, 145, 166, 175, 199, 206, 207, 242, 244, 246,

272, 273, 274, 283, 289, 433, 435, 436, 447, 448, 449, 450

Nevin, Rev. Dr. R. J., 134, 139, 169, 263, 267

Newnham, F. J., 139, 151, 165, 207, 288

Newton, Captain H., 343

Nicolls, J. A. (the late), 138

Nolan, R., 460, 463

Nopcsa, Baron A., 315

Norie, Major-General E. M., 71

Norrie, G. M., 273, 274, 445

North, M. K., 19

Northbrook, Earl of, 49

Norton, Captain C. E. G., 461, 465

Nugent, Major A., 38, 100, 339, 345

Nugent, J., 71

Oakshott, Dr. 151, 191 O'Brien, H. S., 18 Occleston, S. V., 344, 477 O'Donnell, J. C., 400 Ohlsson, A., 108, 118, 123, 130, 132, 139, 145, 147, 149, 151, 159, 161, 166, 167, 171, 174, 180, 187, 189, 191, 193, 199, 208, 215, 242, 249, 253, 263, 266, 270, 271, 289, 293, 297, 305, 310, 393, 397, 419, 440 Oldfield, A. R., 343, 344 Oldfield, C. B., 203 Oldham, L. W. S., 335, 336, 346 Oliver, Captain H. V., 335 Orr-Ewing, Major J. A., 318, 344, 362, 409 Osborne, Lieut. F. C., R.N., 6 Osborne, J. W., 6 Osmaston, B. B., 444, 445 Oswell, W. C. (the late), 73, 186, 215, 419, 440, 447, 450

Page, R. P., 18 Paget, Lord Alexander (the late), 17 Paget, Colonel Arthur, 157, 232, 235, 248, 250, 253, 257, 273, 274, 305, 437, 453, 456 Paget, J. Byng, 113, 249, 274, 436 Paget, Major W. L. H., 274 Palffy, Count T., 179, 266, 302, 315, 395 Pallas, --, 220 Palmer, Major G., 353 Pank, Major P. Dunell, 203 Paris Museum, 5, 11, 35, 48, 49, 52, 58, 67, 87, 90, 108, 113, 118, 119, 123, 126, 142, 179, 182, 185, 187, 213, 218, 219, 221, 223, 251, 256, 276, 291, 293, 302, 306, 313, 319, 326, 402 Partridge, H., 445

Paul, Major E. T., 400, 413, 431 Pausinger, F., 27 Payne, Colonel G. M., 443 Payne-Gallwey, A., 144 Payne-Gallwey, S., 235, 248, 253, 257, 273 Peake, R. C., 271 Pearce, Sir W. G., Bart., 18 Pearse, Arthur, 326 Pearse, Surgeon-Captain A., 218, 361, 400, 413 Pease, A. E., 113, 157, 225, 226, 233, 235, 237, 248, 249, 253, 256, 271, 273, 279, 301, 305, 315, 364, 455 Pease, E. H., 350 Pease, J. A., 264 Pease, W. E., 225, 237, 350, 364, 387 Peel, C. V. A., 19, 85, 113, 114, 157, 171, 232, 233, 235, 249, 253, 257, 274, 299, 301, 305, 353, 437, 455, 457, 466, 471 Peel, Hugh, 47, 368 Pegu Club, Rangoon, 400 Penrice, G. W., 139, 148, 149, 180, 195, 215, 216, 268, 285, 294, 302, 311, 394, 456 Penton, J. E., 341 Perak Museum, 66, 82, 323 Percy, Colonel G. A., 161, 176, 195, 208, 283 Petheram, Sir Comer, 64 Petherick, Consul J. (the late), 126, 181, 182 Phelps, E. L., 38, 39, 213, 221, 318, 329, 335, 339, 345, 362, 373, 379, 381, 382, 408 Phelps, J. V., 218, 336, 345, 362, 378, 381 Phillips, Lionel, 147 Phillipps-Wolley, Clive, 48, 96, 223, 326, 359 Phipps, A. B., 293, 294 Pigott, J. R. W., 449 Pike, Arnold, 7, 213, 218, 345, 359, 361, 366, 367, 378, 381, 408, 479 Pike, Warburton, 390 Pilcher, Lieut.-Col. T. D., 268, 466 Pilkington, H. C., 18 Pinney, Capt. C. F., 62, 230, 400, 413 Platt, H. J. H. (the late), 69, 418 Platt, Mrs. J. E., 19 Pless, Prince Hans Henry of, 400 Pole-Carew, Colonel R., C.B., 37, 71, 74, 79, 322, 372, 400 Pollen, A. H., 45, 63, 94 Pollen, S. H., 463 Ponsonby, J., 17, 120, 242 Poole, T., 187 Portland, Duke of, 16, 22, 47, 346, 372, 401, 406, 431

Potocki, Count J., 64, 113, 248, 273, 281, 401,

423, 437, 455, 465, 466

Potter, F. E., 151, 166, 195, 199, 215, 271, 289 Pottinger, Sir H., Bart., 9, 12, 13 Poulett-Weatherley, H., 161, 187, 189 Powell-Cotton, P. H. G., 38, 213, 218, 232, 248, 253, 257, 273, 298, 318, 322, 329, 335, 344, 361, 377, 378, 381, 400, 408, 437 Power, W. S., 6 Powerscourt, Viscount, 10, 12, 14, 16, 17, 18, 20, 23, 25, 26, 27, 28, 29, 30, 44, 45, 46, 51, 52, 59, 60, 62, 63, 79, 84, 87, 373, 400, 406, 411, 443, 444 Prendergast, W. H., 417 Pretoria Museum, 440 Prettyjohn, Col., 37 Pringle, Captain J. W., 142, 242 Proceedings of the Zoological Society, 343, 349, 360, 370, 385, 416, 443 Pryer, W. B., 66, 418 Przewalski, Colonel, 42 Pulley, S., 98, 263, 392, 450 Pulteney, Capt. W. P., 466 Punjab (21st) Infantry Mess, 382 Purvis, Captain J. H., 76, 345

Queen, Her Majesty the, 449 Quicke, Captain F. C., 113, 204, 248, 253, 256, 274, 344, 424, 429 Quilter, Ely, 368 Q.O. Corps of Guides, Officers' Mess (see under Guides)

Radmore, G. R., 79 Radziwill, Prince, 12, 476 Rae, J., 390 Rankin, C. H., 344 Rankin, R., 7, 95, 326, 345, 468 Ratcliff, S., 7, 12, 327, 366 Ratibor, Duke of, 25 Ratibor, Prince Victor, 26 Rattray, Major R. H., 203, 229, 318, 321, 341, 383 Rawlinson, Major Sir H. S., Bart., 381 Rawstorne, L., 394 Reade, F. W., 313 Redde, Charles (the late), 443 Rees, J. D., 412, 444 Rehbock, Th., 118, 139, 148, 215, 216, 271, Reid, Percy C., 193 Rendall, Dr. Percy, 52, 64, 74, 100, 108, 122, 126, 130, 145, 148, 149, 150, 151, 157, 161, 162, 166, 169, 171, 176, 180, 187, 191, 193, 194, 199, 216, 225, 227, 233, 251, 253, 263,

264, 266, 267, 274, 281, 285, 289, 291, 296, 305, 310, 323, 325, 350, 353, 384, 397, 421, 425, 426, 429, 436, 456, 467, 471 Renton, A. Leslie, 74, 204, 249, 257, 274, 346, 372, 374, 381, 437 Reynolds, Captain H., R.N., 19 Rhodes, Cecil, 440 Rhodes, Colonel F., D.S.O., 262, 300 Rice, Major-General W., 413, 453 Richards, G., 118, 134, 139, 147, 151, 162, 166, 167, 169, 172, 187, 191, 194, 264, 266, 284, 289, 293, 297 Richardson, J. J., 113, 248, 256, 274 Rifles (60th) Mess, 382 Ritchie, D. Norman, 176, 263 Rivett-Carnac, J. S., 329 Roberts, Field-Marshal Lord, V.C., 372 Robinson, F. W., 19 Robinson, Captain S. L., 413 Rodney, Lord, 47, 367 Rodon, Major G. S., 332 Rogers, A., 11, 44, 367 Rogers, Captain J. M., 20 Rohan, Prince Lulu, 25 Roosevelt, Theodore, 6, 11, 45, 92, 94, 104, 326, 406 Root, W. H., 405 Rose, J. (the late), 256 Rotheram, A. M., 461 Rothschild, IIon. Walter, 5, 7, 38, 41, 42, 46, 55, 59, 63, 74, 79, 82, 90, 98, 107, 108, 109, 110, 111, 113, 118, 122, 125, 128, 130, 132, 134, 142, 145, 149, 150, 151, 152, 154, 158, 159, 162, 163, 166, 174, 179, 180, 181, 184, 187, 191, 194, 199, 200, 201, 204, 207, 208, 209, 210, 213, 218, 223, 233, 237, 242, 245, 250, 251, 253, 257, 262, 266, 267, 268, 270, 277, 279, 284, 289, 291, 294, 298, 305, 310, 313, 316, 323, 325, 326, 329, 332, 335, 339, 341, 346, 350, 352, 355, 357, 358, 362, 368, 370, 373, 379, 381, 384, 385, 387, 390, 392, 397, 399, 406, 408, 413, 435, 440, 475, 476 Rotton, Captain J. G., 465 Roxburghe, Duke of, 169, 263 Royal Artillery Mess at Attock, 382 Royal Artillery Mess, Woolwich, 382 Rube, C., 118 Rudd, C. D., 118, 139, 145, 166, 193, 215, 262, 271, 284, 289, 298, 310, 393, 436, 440 Rüppell, Dr. (the late), 148, 196 Russel, P., 82, 468 Russell, Major B. B., 461

Russell, E., 179 Russia, H.I.M. the Czar of, 404 Russia, H.I.M. the Empress of, 372 Ryan, J., 64 Rylands, W. P., 300 Ryley, A., 271

Sagar-Musgrave, A. M., 195 Sagar-Musgrave, R. M., 301 Salt, H. (the late), 419 Salt, Captain T. A., 339, 381 Sandbach, Major A. E., 455 Sanders, R. A., 21 Sanderson, Colonel G. P. (the late), 443, 477 Sangster, Guy H., 201, 291 Saunders, J. Carr, 10, 57, 71, 74, 77, 79, 92, 105, 116, 128, 189, 267, 277, 313, 352, 361, 373, 378, 381, 399 Sauter, F., 405 Sawyer, R. H., 132, 367 Saxe-Coburg and Gotha, H.R.H. the Duke of, 7, 11, 16, 25, 26, 27, 28, 29, 30, 33, 47, 63, 73, 76, 84, 132, 154, 175, 204, 284, 301, 315, 329, 341, 361, 362, 378, 394, 406, 448 Saxe-Coburg and Gotha, Prince August Leopold of, 29 Saxe-Coburg and Gotha, Prince Philip of, 25, 26, 27, 28, 29 Saxony, H.M. the King of, 29, 30 Scarlett, J. L., 48, 94, 368, 469, 475, 477 Scheibler, Count, 9, 12, 45, 74, 105, 116, 176, 225, 232, 242, 249, 256, 273, 276, 302, 306, 316, 326, 327, 367, 392, 400, 428, 435, 436, 443, 448, 455, 457, 461, 464, 466, 477 Scheibler, Countess, 431, 437 Schopf, Dr. A., 270 Schoverling, Daly, and Gales (Messrs.), 44 Science and Art Museum, Dublin, 7, 11, 20, 46, 57, 60, 74, 79, 82, 105, 118, 123, 126, 145, 172, 176, 179, 208, 211, 262, 281, 298, 313, 316, 339, 346, 347, 370, 372, 383, 385, 390, 400, 401, 402, 411, 476 Science and Art Museum, Edinburgh, 306, 387 Sclater, P. L., 228, 349 Scott, E. D., 289, 294, 301, 393 Secretan, B., 138 Seely, C. H., 38, 73, 230 Selous, F. C., 18, 33, 34, 48, 91, 92, 94, 95, 103, 105, 118, 122, 123, 130, 132, 133, 134, 135, 138, 139, 145, 146, 147, 149, 151, 161, 166, 167, 169, 171, 174, 175, 187, 189, 191, 194, 199, 208, 215, 216, 261, 262, 263, 264, 265, 266, 267, 270, 271, 284, 285, 289, 293,

294, 297, 298, 302, 307, 310, 316, 352, 392, 393, 421, 428, 436, 440, 441, 448, 450, 451, 453, 454, 455, 456 Seton-Karr, H., 19, 24, 44, 46, 48, 366 Seton-Karr, H. W., 457, 475 Severtzoff, Dr., 49 Sharp, A. H., 64, 457 Shaw, F. G., 138 Shaw, Otho, 6, 10, 45, 47, 48, 74, 104, 105, 204, 213, 335, 339, 341, 344, 361, 366, 378, 379, 381, 412, 460 Sheard, W. F., 10, 44, 365, 366, 405, 476 Sherwill, W. C., 431 Shields, G. O., 47 Shields, J. O., 366 Shirres, Major J. C., D.S.O., 333, 334, 382 Simson, F. B., 76 Sing, Rajah Gulab, 37 Singer, Washington, 74 Sitwell, Major C. G. H., 179 Sitwell, Lieut.-Colonel W., 271, 299 Skeffington-Smyth, R., 429 Slaney, Kenyon, 97 Smee, Captain, 455 Smith, Sir Andrew (the late), 118, 134, 425 Smith, Major A. E., 119, 200, 207, 242, 244 Smith, Berthold, 359 Smith, Dr. Donaldson, 175, 273, 305, 450, 451 Smith, G. de H., 413 Smith, G. E., 120, 179, 207, 242, 245, 283, 450 Smith, Dr. H., 470 Smith, Major Hamilton, 150 Smith, Captain J. Manners, V.C., C.I.E., 343, 374 Smith, L. Gisborne, 64 Smith, Norman B., 248, 299, 303, 304, 454, 466 Smith, R. Gordon, 5, 445 Smitheman, F., 122, 134, 147, 161, 189, 194, 262, 267, 293, 294 Smyth, Sir Greville, Bart., 22, 63, 405 Sneyd, Ralph, 20 Somerset, Hon. H. S., 47 South African Museum (see Cape Town Museum) Speke, Captain (the late), 164, 176, 179, 285, Spence, A. H. Ogilvy, 318, 329 Spreckley, Captain J. A., 261 Stalkartt, Surgeon W. H. S., R.N., 283 Stanley, C. Sloane, 386, 387 Staples, F. S., 392

Stephen, H. L., 108 Stephenson, Guy, 19 Sterndale, R. A., 332, 463, 468, 471 Stevens, Montague, 475, 476 Stewart, Major Colvin, 229 Stewart, C. M. D., 444 Stirling, W., 18 Stobart, Captain W. E., 71 Stockley, Major V. M., 437 Stonor, Hon. E., 444, 445 Storey, H., 445 Stracey, Captain C. E., 5 Strachey, Captain, 346 Straker, A. H., 44, 82, 177, 232, 235, 249, 274, 286, 301, 304, 305, 399, 436, 450 Streatfeild, Major Henry, 76, 431 Struben, F., 263, 298, 310, 311 Stuart, G. R., 97 Studholm, E. C., 31 Studley, J. T., 368 Sulivan, Lieut.-Colonel G. D. F., 12, 154, 203, 204, 229, 399, 408 Surtees, H. E., 388 Sutton, Lieut., 461 Swanzy, F., 180 Swayne, Major H. G. C., 112, 113, 157, 170, 172, 177, 227, 232, 235, 250, 253, 256, 274, 286, 300, 302, 304, 436, 480 Swinhoe, R. (the late), 54, 79, 82, 219, 401 Swire, C. M., 135, 392 Szechenyi, Count, 28 Szechenyi, Count Belà, 26, 27

Tait, Dr. Wm., 6 Talbot, Major, 359 Tankerville, Earl of, 390 Tanner, Colonel H. C. B. (the late), 41, 372 Tate, Henry, 17 Tate, Mrs. Henry, 17 Taylor, Major F. H., 335, 382 Taylor, H. Douglas, 74, 76 Taylor, J. B., 440 Taylor, Captain N. C., 230, 322, 346 Taylor, T., 98 Teck, Duke of, G.C.B., 378 Teleki, Count Arpad, 314, 315 Telford, W., 31 Tennant, Sir Charles, Bart., 18 Tennant, Edward P., 41, 55, 373 Ternan, Colonel Trevor, 119, 120, 158, 163, 179, 185, 197, 242, 452 Thatcher, G. G., 379 Thebaw, King (the late), 443

Thellusson, Hon. F., 11, 46, 94, 405 Thomas, Captain E. A. D'Arcy, 463 Thomas, J. H., 7, 24, 225, 237, 279 Thomas, Oldfield, 228, 244, 330 Thomson, A. Y., 413, 431 Thomson, Joseph (the late), 241 Thorold, Dr. W. G., 42 Thurlow, Lord, 6 Thynne, Hon. T., 122, 135, 195, 208, 215, 284, 285, 456 Thynne, U. O., 205 Tichborne, Sir H. D., Bart., 76, 113, 232, 249, 273, 305, 332, 436 Tickell, Captain E. J., 120, 245 Tidswell, Captain E. C., 352 Timins, Captain C. S., 114, 249, 273, 298, 306, 332, 413 Timmins, H., 122, 139, 161, 176, 194, 261, 267, 294 Tolhurst, F. B., 10, 44 Tollemache, Hon. S., 367, 401, 412 Tolman, J. C., 476 Tompson, E. W., 297, 421 Tompson, Captain W., 413 Tonnerre, L. Fabre, 431 Toppin, H., 353 Tottenham, Captain R. L., 329, 345, 382 Townley-Parker, Captain (the late), 475 Townshend, E. Lee, 113, 273 Trapmann, A. G. (the late), 323 Trauttmansdorff, Count Ferdinand, 26, 45 Travancore, Maharaja of, G.C.S.I., 38, 62, 79, 213, 229, 337, 343, 344, 352, 362, 372, 378, 381, 411 Travers, Dr., 423 Trevor, A. S., 235, 253, 273, 437 Trevor, Captain H., 213, 229, 334, 343, 361, 379 Trollope, Major F., 193 Truninger, L., 401, 413 Tulloch, W. A., 46 Turkey, H.I.M. the Sultan of, 460 Turner, Arthur W., 411 Turner-Turner, J., 5, 47, 92, 95, 327, 366 Tweedmouth, Lord, 15, 16, 17 Twopeny, C. D., 64, 154, 204, 281

Ubsdell, J. E. (the late), 204, 329

Vandeleur, Captain C. B., 213, 218, 229, 318, 344, 361, 409 Vandeleur, C. F. S., D.S.O., 185, 208, 456 Vander-Byl, -, 129

Vander-Byl, C. F., 229 Vander-Byl, P. B., 11, 47, 139, 195, 229, 232, 249, 257, 274, 302, 305, 311, 316, 327, 437 Van-Ness, W., 134, 263, 266, 294, 298, 311, 435 Vans-Agnew, A., 98 Vansittart, H., 329 Van-Son, H., 82, 417, 423, 426, 432 Van-Son, J. C., 82 Van Zeller, F., 297 Varndell, C. R. (the late), 439 Vaughan, J., 263, 344 Vaughan, J. E., 318 Veernhof, Colonel, 423 Venables-Kyrke, R. H., 191, 367, 393 Verschoyle, Captain E. G., 46, 47, 367 Vertue, E. B., 421 Victoria (B. C.) Museum, 326, 366, 469 Villiers, Captain C. H., 120, 274, 305 Vincent, B., 232, 253, 274, 346, 381, 399 Vivian, A. Pendarves, 45, 469 Vivian, Colonel Ralph, 46, 47, 85, 105, 106, 109, 155, 169, 176, 239, 250, 267, 286, 301, 395, 406, 437, 452 Von André, A., 26, 27, 29 Von Massow, Baron, 411 Vryburg Club, 215 Wade-Dalton, Lieut.-Col. H., 64, 74, 281, 344, 345, 384, 463 Wahlberg, J., 148 Wahrmann, R., 114, 249, 250, 273, 305, 401, Wait, Capt. A. M'Lean, 289

Waley, A., 71, 82, 412 Walker, Surgeon-General, 153 Walker, A. Barclay, 476 Walker, H. G., 6 Walker, Sir Peter, Bart., 6, 10, 11, 12, 45, 94, 105, 327, 367, 400, 431, 445, 475, 476 Walker, W. H., 16, 18, 445, 463 Wallace, A. R., 69 Walshe, F. W. H., 38, 218, 318, 337, 361, 379, 381 Walter, A. J., 325, 401, 412 Walter, Major R. L., 344 Ward, Major A. E., 37, 62, 76, 82, 213, 318, 321, 329, 335, 341, 343, 361, 379, 381, 465, 468, 476 Ward, Hon. John, 52, 172, 216, 263, 267, 301, 363, 364 Ward, Rowland, 100, 127, 139, 141, 145, 155,

157, 158, 185, 189, 209, 216, 218, 225, 226,

283

235, 237, 253, 256, 270, 279, 291, 294, 311, 368, 370, 388, 393, 399, 408, 421, 445, 451, 456, 463, 468, 475, 476, 479 Ward, Hon. R. A., 52, 135, 171, 208, 226, 237, 263, 264, 267, 279, 364, 373, 378, 387 Warrand, Colonel A. J., 85 Warrand, H. M., 84 Waterford, Marquis of (the late), 443 Watkins, F., 394 Watkinson, F., 122 Webb, W. F. (the late), 215, 300, 374, 392 Webster, G., 18 Welch, Captain N. C., 107, 148 Welch, Lieut. G. H., R.N., 246 Wellby, Captain M. S., 344, 361, 451 Wellcome, H. S., 92, 94 Weller-Poley, E., 17 Wellington, Duke of, 37 Welsh, C. N., 232 Wenlock, Lord, G.C.I.E., 412, 444, 445 Werne, 182 Wernher, Julius, 18 Westmacott, C. R., 31 Westminster, Duke of, 6, 8, 10, 23, 46, 60, 372, 379, 440, 448, 450, 479 Whitaker, J., 7, 57, 71, 104, 169, 281, 399, 428 Whitaker, J. J. S., 25, 225, 226, 237, 278, 279, 364 Whitby, Lieut.-Colonel F. H., 72, 73, 400, 461, 465 White, A., 466 White, Captain E. D., 76 White, Captain G. Dalrymple, 45, 46, 367, 470 White, Surgeon-Captain W., 476 White, Captain W. Westropp, 476 Whitehouse, J. H., 11 Whitehouse, Langford, 423 Whitely, H., 99 Whitestone, Surgeon-Capt. C. W. H., 411 Whitney, H. P., 63 Whittall, F. E., 58 Whittall, H. O., 34, 353, 477 Wilczek, C. H., 347 Wilding, Captain C. A., 108, 397 Wilkin, Rupert, 31 Wilkinson, Capt. P. S., 277 Williams, Surgeon-Capt. E. M'K., 73, 462, 465, 466 Williams, H. H., 175, 284, 394 Williams, J. G., 85 Williams, Capt. L. O., 147, 174, 195, 199,

Williams, Rhys, 386, 387 Williams, Major W. H., 175, 185, 208, 245 Williamson, F. C., 5, 368 Williamson, H., 385 Williamson, Capt. W. H., 235, 306, 361 Willoughby, Sir John, Bart., 134, 141, 169, 176, 194, 208, 244, 245, 257, 266, 275, 285, 300, 311, 392, 393, 436, 437, 454, 457 Willshire, W., 251 Wilson, Mr. ("Mountaineer"), 81, 328 Wilson, A., R. N., 6 Wilson, Captain F. A., 156 Windischgraetz, Prince Hugo, 25 Wing, Major, F. D. V., 71 Wingfield, W. J. R., 463 Winwood, W. Q., 71 Wolverton, Lord, 250, 256, 273, 401, 413, Wood, Major C. B., 400 Wood, Gordon, 19 Wood, Surgeon-Captain H. S., 82, 417, 418

Wood, J. L., 343
Wood, J. W., jun., 368
Wood, Philip, 463
Wood, T. W., jun., 10
Woodd, Basil H., 456
Wordsworth, R., 62
Wormold, F. W., 345
Worthington, F. V., 161, 261, 262, 266
Wray, J., 116
Wrey, G., 45, 96, 367, 406

Yellowby, W., 454 York, H.R.H. the Duke of, 449 Young, J. G. K., 315 Young, J. W. R., 366 Yule, James, 122, 171, 428 Yule, John, 122, 189, 209, 285, 289, 292, 293,

Zamoyski, Count T., 437, 454 Zoological Society (see under Proceedings)

INDEX OF SPECIES

		PAGE	1	PAGE
Abyssinian Buffalo		. 395	Babirusa	. 426
,, Bushbuck		. 286	Babirusa alfurus	426
,, Duiker		. 148	Banded Duiker	. 152
,, Ibex .		. 350	Banting	. 416
,, Oribi .		. 162	Barasingh. See Hangul.	
Addax		. 278	Barbary Red Deer	. 35
Addax nasomaculatus		. 278	Barking Deer	. 31
Addra Gazelle .		. 250	Basilan Sambar	. 67
Æpyceros melampus		. 206	Bavian Deer	. 69
,, petersi .		206, 209	Bay Duiker	. 151
African Elephant .		. 446	Bears	. 473
,, Rhinoceros		433	Beatrix Oryx	. 276
Alaskan Bighorn .		. 368	Bedford's Deer	. 35
Alces machlis .		. 8	Beira	. 258
Alpine Ibex		. 347	Beisa	. 272
Altai Wapiti .		. 49	Bharal	360
American Bighorn		. 365	Bison	. 403
,, Bison		. 405	Blackbuck	. 202
" Deer .		. 91	Black-tailed Deer	. 96
Ammodorcas clarkei		. 252	Blesbok	. 131
Ammon Sheep .		. 375	Blue Duiker	. 149
Ancient British Red De	eer .	. 23	Bohor Reedbuck	. 196
,, Continental Re	ed Deer	. 30	Bongo ,,	. 287
Anoa		. 402	Bontebok	. 129
Antidorcas euchore		. 214	Bos bison	. 405
Antilocapra americana		. 104	,, bonasus	. 403
Antilope besoartica		202	,, bubalis	. 398
Arabian Gazelle .		. 227	,, caffer	. 391
,, Ibex .		. 348	,, ,, aquinoctialis .	. 395
,, Tahr .		. 330	,, ,, brachyceros	. 396
Armenian Muflon.		. 384	,, ,, planiceros	. 396
Arna		398	,, ,, nanus	. 396
Arui		. 363	,, depressicornis	. 400
Asiatic Ibex		. 342	,, frontalis	. 415
Astor Markhor .		. 333	,, gaurus	. 410
Atlas Gazelle .		. 226	,, grunniens	. 407
Aurochs. See Bison.			,, indicus .	. 419
Avie		72	condaicue	. 419

				PAGE	1			PAGE
Bos taurus				419	Cephalo	phus abyssinici	us .	148
Boselaphus tragocamelu	s			280	,,	coronatus		148
Brindled Gnu .				136	,,	doriæ		152
Broad-Horned Antelop	e			287	,,	dorsalis		151
Bubaline Hartebeest				106	1	grimmi		146
Bubalis boselaphus				106	,,	maxwelli		150
,, cama .				117	1 11	monticola		149
,, cokei .				115	,,	natalensi	ς	150
,, jacksoni .				119	,,	sylviculto		152
,, lichtensteini				121	Cervica	pr a arundinun	n.	192
,, major .				107	,,	bohor .		196
,, swaynei .				112	,,	fulvoruful	a	198
,, tora				109	' ,,	,,	chanleri	200
,, ,, neumanni	•			111	,,	,,	subalpina	199
Budorcas taxicolor				325	,,	redun ca		201
Buffaloes				391	Cervuli	is crinifrons		. 83
Buffon's Kob .				183	,,	fea .	•	83
Burchell's Rhinoceros				439	,,	lachrymans		82
Burmese Serow .				323	,,	muntjac.		81
,, Stag .				78	Cervus			40
Bushbuck				282	٠,	albirostris.		41
					,,	axis .	•	72
Cabul Markhor	•	•		340	,,	barbarus .		15
Cape Buffalo	•	•	•	391	,,	canadensis	• •	42
,, Bush-Pig	•	•		425	· ,,	,,	aticus .	· · 49
" Hartebeest	•	•		117	,,		hdorfi .	50
Capra caucasica .	•	•		356	•••	cash m ir i anus	• : .	36
,, cylindricornis	•	•		358	,,		rarcandensis	0., 0,
,, falconeri .	: .	•	•	333	"	dama	• •	56
,, ,, cashmir		•	•	333	,,	duvaucelli.		75
,, ,, jerdoni		•		340	-,,	elaphus .	•	14
,, ,, megacer	*0 5	•	•	340	,,,	,, maral	• •	32
,, hircus agagrus	•	•		353	٠,	eldi		78
., ,, blythi	•	•		351	,,	giganteus .	•	59
,, ibex	•	•	•	347	٠,	hippelaphus		68
,, nubiana .	•	•	•	348	"	,,	oluccensis	69
,, pallasi .	•	•		350	,,	hortulorum		54
,, pyrenaica .	•	•	•	354	,,	kuhli .		69
,, sibirica .	•	•		342	,,,	mesopotamicus		58
,, vali	•	•		350	,,	moluccensis		69
Capreolus pygargus	•			86	,,,	porcinus .		70
,, <i>vulgaris</i> Caribou	•	•	•	83	,,	schomburgki	•	77
Caspian Red Deer	•	•	٠.	I	,,	sica taëvanus .		51
Caucasian Bharal.	•	•		5, 32	,,	unicolor .		53
71	•	•	•	358	,,			61
,, Ibex .	•	•		356	"	,, dejear		67
Cemas argyrochætus caudatus .	•	•		319	,,	,, equir		65
	•	•	•	319	"	,, nigri		67
• •	•	•		319	,,		opinus . hoci	. 67
,, goral,	•	•		317	,,	,,	wet .	
,, griscus . Central African Buffalo	•	•		319 306	Cevlon	xanthopygus Buffalo		
Contai Automi Dullillo	,			· · ·	· CEAIOII	Lunaio .		. 401

					PAGE	1	PAGE
Ceylon Sambar .					64	Elaphurus davidianus	. 89
Chamois	•	•	•	•	•	•	_
Chanler's Reedbuck	•	•	•	•	314	Eld's Deer	. 78
	•	•	•	•	200	Elephant, African	. 446
Chiru Chita	•	•	•	•	212	", Indian	. 442
	•	•	•	•	47 I	Elephas africanus	. 446
Chital	•	•		٠	72	,, indicus	- 442
Clarke's Gazelle .	•	•	•	٠	252	,, primigenius	. 452
Clouded Leopard.	•	•	•		468	Elk	. 8
Cobus cob	•	•	•	٠	183	English Park Red Deer	. 22
,, defassa .		٠	. •	•	178	Equine Deer	. 65
,, ,, penrice		•	٠	•	180	European Bison	. 403
,, ,, unctuo		•	•	٠	180	,, Lynx	. 470
,, ellipsiprymnus	•	٠	•	٠	173	,, Muflon	. 386
,, lichi	•	•	•	•	186	,, Roe	. 83
., leucotis .	•		•		182	Extinct Giant Fallow Deer	. 59
,, maria .			•		181	Fallow Deer	. 56
,, thomasi .					185	Felis concolor	. 469
,, vardoni .					188	,, leo	. 453
Coke's Hartebeest	•				115		433
Connochætes gnu .					143		. 468
,, taurinus					136	,,	. 469
,, ,,	albojub	atus			141	4	. 464
Corsican Red Deer					15	· · · · ·	
Cynælurus jubatus					471	,, tigris	. 458
Cyprian Muflon .				•	385	Formosan Sika	. 467
Dama Gazelle .						8 1	. 53
Dama Gazene . Damaliscus albifrons	•	•	•	•	251		. 66
	•	•	•	•	131	Fringe-eared Beisa	275
,, corrigum		•	•	•	126	Four-horned Antelope	. 153
	jimela	•	•	•	127	Gaur	. 410
,, ,, 	tiang	•	•	•	126	Gayal	415
,, hunteri	•	•	•	•	124	Gazella arabica	. 227
,, lunatus	•	٠	٠	•	133	,, bennetti	. 228
,, pygargus	•	٠	•	•	129	,, cuvieri	. 226
Damaraland Duiker	•	•	•	•	159	,, dama	. 251
Decula Antelope .	•	•	٠	•	286	,, ,, mhorr	. 251
Defassa Sing-Sing	•	•	•	•	180	., dorcas	. 224
Derbian Eland .	•	•	•	•	312	· ,, granti	. 243
Dibatag	•	•	•	•	252	., gutturosa	. 221
Dik-Dik Antelopes	•	•		٠	156	,, isabella	. 238
Domestic Goat .	•	•	•	•	353	· ,, leptoceros	. 236
,, Oxen	•		•	•	418	A.J. J	. 234
,, Sheep .	•				387	A.4	. 246
Oorcas Gazelle .	•	٠	•		224	Adadama Juan	. 217
Dorcatragus melanoti	s.				258	A	. 219
Duiker	•				146		. 250
Owarf Antelopes .	•				156		. 240
,, Buffalo .					396		
Oybowski's Sika .					54	.4.1.2	. 247
East Caucasian Tur					250		231
Edmi Gazelle .	•	•	•	•	358	,, subgutturosa	. 222
Sami Gazene .	•	•	•	•	226	,, thomsoni	. 241

					PAGE		PAGE
Gemsbuck					269	Indian Sambar	61
Gerenuk					255	Inyala. See Nyala.	
German Red Deer					25	Irish Elk	59
Giraffa camelopardalis					101	,, Red Deer	20
,, capensis .					102	Isabella Gazelle	238
Giraffe					101		
Goa					217	Jackson's Hartebeest	119
Goat					351	Jaguar	469
Goral					317	Japanese Sika	51
Grant's Gazelle .					243	" Serow or Goral	319
Gray Rhebok .					190	Javan Ox	416
Gray's Waterbuck					181	,, Rhinoceros	432
Greater Kudu					296	,, Rusa	68
Grysbuck:					167		
•					•	Kamschatkan Bighorn	319
Haggard's Oribi .					163	Kansu Takin	326
Hairy-fronted Muntjac					83	Kashmir Stag	36
Hangul					36	Kirk's Dik-Dik	158
Haploceros montanus					326	Klipspringer	170
Harnessed Antelope.	See E	Bushb	uck.		3	Kob	185
TT					106	Kongoni	115
Hemitragus hylocrius					331	Korrigum	126
					330	Kudu	296
,, jemlaicus					328		-90
Hemprich's Dik-Dik	_				159	Leopard	464
Heuglin's Gazelle.		•			239	Lesser Kudu	303
Himalayan Goral .	•	•	•		317	,, Reedbuck	198
,, Ibex .	•	•			342	Lhasa Stag	211
,, Serow	•	•	•	•	320	Lichi	186
T-1-		•	•	•	328	Lichtenstein's Hartebeest	121
Hippopotamus .	•	•	•	•	421	Lion	
Hippopotamus amphibis	•	•	:	•	421	, Skins	453
liberiensi			:	•	421	" or "	454
Hippotragus equinus		•	•	•	265	,, Skulls	456
• • •		•	•	•	260		255 165
Hog Deer	•	•	•	•	70	Livingstone's Antelope	_
Hungarian Red Deer	•	•	•	•	•	Luzon Sambar	236
Hunter's Hartebeest	•	•	•	•	25		67
Hunting-Leopard	•	•	•	•	124	Lydenburg Reedbuck	199
~ .	•	•	•		471	Lynx	470
Hyæna Hybrid Hartebeest	•	•	•	•	472	16-1	
riyona riantebeest	•	•	•	•	120	Madoqua damarensis	159
Ibex						,, guentheri	156
	•	•	•	•	354	,, hemprichiana	159
Impala	•	•	•	•	206	,, kirki	158
Indian Antelope .	•	•	•	٠	202	,, phillipsi	156
,, Buffalo .	•	•	•	•	398	,, saltiana	155
,, Elephant .	•	•	•	•	442	,, swaynei	156
,, Gazelle .	•	•	•	٠	228	Malayan Sambar	65
,, Leopard .	٠.	•	•	٠	464	Mammoth Elephant	452
,, Hunting-Leopar	ď	•	•	٠	47 I	Manchurian Wapiti	50
-	•	•	•	•	81	Maral	32
,, Rhinoceros					430	Marco Polo's Sheep	37 I

				PAGE	P	AGE
Markhor				333	Oryx leucoryx	277
Marsh-Deer				97	Ovibos moschatus	389
Maxwell's Duiker .				150	Ovis ammon	375
Masama americana				91	., ,, hodgsoni	377
,, ,, mex	cicana			92	,, aries	387
,, antisiensis				99	,, canadensis	365
,, bezoartica		•		99	., ,, dalli	368
,, columbiana				96	,, ,, nivicola .	369
,, dichotoma				97	,, lervia	363
,, hemionus				93	,, musimon	386
,, nemorivaga				100	,, nahura	360
Mesopotamian Fallow I	Deer .			58	,, orientalis	384
Mexican Deer .				92	., ,, ophion	385
Mhorr				251	,, poli	371
Moluccan Rusa .				69	,, vignei .	380
Mongolian Argali .		-		379	,, ,, cycloceros	384
,, Gazelle		•		221	,, ,, ,,	3.4
Moose			•	8	Pampas Deer	99
Moschus moschiferus	•		•	100	Para	70
Mountain Reedbuck		•	•	198	Pantholops hodgsoni	212
Muflon	•	•	•	384	Pekin Sika	54
Mule-Deer		•	•	93	Pelea capreolus	190
Muntjac		•		93 81	Pelzeln's Gazelle	234
Musk-Deer		•	•	100	Penrice's Waterbuck	180
_		•	•	389	Père David's Deer	80
" Ox		•	•	309	Persian Gazelle	222
Maran Dandhuala						
Nagor Reedbuck .	• •	•	•	201	,,	351
Narwhal		•	•	479	Peruvian Guemal	99
Natal Duiker .	•	•	•	150	Peters's Gazelle	246
Nemorhædus bubalinus			•	320	Phacocharus athiopicus	427
·· · · · · · · · · · · · · · · · · · ·	sumat	rensis	•	323	Philippine Deer	67
,, crispus			•	319	Pir Panjal Markhor	333
Nesotragus livingstonia			•	165	Polar Bear	475
,, moschatus		•	•	164	Potamocharus charopotamus	425
Neumann's Hartebeest				111	,, porcus .	425
New Zealand Red Deer	r.	•	٠	31	Prong-horn	104
Nilgai		•	٠	280	Przewalski's Gazelle	219
Nilgiri Tahr .			•	331	Puku	188
Norwegian Red Deer				24	Puma	469
Nyala				288	Pyrenean Ibex	354
Nyasaland Gnu .				138		
					Kangifer tarandus	1
Oreotragus saltator				170	Red Deer	14
Oribi				160	,, Duiker	150
Oribia haggardi .				163	Red-fronted Gazelle	240
,, montana .				162	Red River-Hog	425
,, nigricaudata				163	Reedbuck	192
,, scoparia .				160	Reindeer	Ī
Oryx beatrix .				276	Rhaphiceros campestris	168
,, beisa				272	,, melanotis	167
,, callotis				275	Rhinoceros bicornis	433
,, gazella				269	,, simus	439
-					•	

					PAGE		PAGE
Rhinoceros sondaicus			•		432	Tahr	328
,, sumatrensis					432	Takin	325
,, unicornis					430	Taurotragus derbianus	312
Roan Antelope .					265	,, oryx	308
Rocky Mountain Goat					326	Tenasserim Muntjac	83
Roe Deer					83	Tetel Hartebeest	109
Rooi Rhebok .					198	Tetraceros quadricornis	153
Rupicapra tragus .					314	Thameng	78
1 1 0					٠.	Thomson's Gazelle	241
Sable Antelope .					260	Thorold's Deer	41
Saiga					210	Tiang	٠.
Saiga tatarica .					210	Tibet Antelope	212
Salt's Dik-Dik .					155	,, Argali	377
Sambar Deer .		•	-	•	61	,, Gazelle	217
Sassaby	•	:			133	" Muntjac	82
Schomburgk's Deer	•		•	•	*33 77	Tiger	458
Scotch Red Deer .	•	•	•	•	16	,, Skins	460
Senegal Gazelle .		•	•	•		Cl11.	•
Senegar Gazene . Senegambian Eland	•	•	•	٠	240	m''.	463
^ -	•	•	•	•	312	Tr	127
	•	•	•	•	323	Tragelaphus angasi	,
Shapu	•	٠	•	•	380		288
Sheep	•	٠	•	٠	360	,, decula .	286
Shou	•	•	•	٠	40	,, euryceros	287
Siberian Argali .	•	•	•	٠	375	,, gratus	290
", Roe .	٠	•	•	•	86	,, scriptus	282
Sig	•	٠	•	•	112	,, spekei	292
Sing-Sing Waterbuck	•	•	•	•	178	,, sylvaticus	283
Situtunga	•				292	Tusks, Boar	423
Snow Leopard .	•	•			467	,, Elephant .	443
Sæmmerring's Gazelle					247	,, Hippopotamus	421
Somali Dik-Diks .					156	., Musk-Deer .	100
,, Gazelles .			_		∫ 231	Udad	
	•	-	•	•	L 235	Uganda Kob	363
Spanish Tur	•	•			354	TT-1.1	185
,, Red Deer					24	77 .23	380
Speke's Gazelle .	•				231	Ursidæ	473
Springbuck					214	Vaal Rhebok	7.00
Steinbuck					168	Vissisian Dans	190
Strepsiceros imberbis					303	Virginian Deer	91
,, kudu .					296	Wallachian Sheep	388
Suleman Markhor					358	Waller's Gazelle	255
Sumatran Rhinoceros					432	Walrus	478
" Serow .					323	Wapiti	42
Sus charopotamus.		. •			425	Wart-Hog	427
" cristatus .					422	Waterbuck	
, porcus					425	317-4 AC' D CC 1	173
" scrofa				Ċ	422	Dt. D'	396
,, scrofa					75	77 1 4 1	425
Swayne's Dik-Dik					156	77 1	290
,, Hartebeest		•	•	•	112	0.111	107
Swift Gazelle .				•	251	W . C	163
Szechuan Sambar.			•	•	67	West Caucasian Tur West of England Red Deer	356
					J/ 1	THE OF LIBRARY INCUITAGE .	21

				PAGE			:	PAGE
White-bearded Gr	nu			141	Yak			407
White-eared Kob				182	Yarkand Gazelle .			222
White Oryx .				277	,, Stag .			39
,, Rhinoceros	;			439	Yellow-backed Duiker			152
White-tailed Deer	:			91				
White-tailed Gnu				143	Zanzibar Antelope			164
Wild Boar .				422	Zebu			418
Wood Brocket	٠.			100	Zubr			403

THE END

ROWLAND WARD

LIMITED

Practical Taxidermists

By Special Appointment to the Courts of Europe

"THE JUNGLE" 166 PICCADILLY, LONDON, W.

TELEPHONE 3644

PRACTICAL AND ARTISTIC TAXIDERMISTS, Designers of Trophies of Natural History, Preservers and Adapters of all Specimens of Animal Life. Natural Features of Animals adapted in Original Designs for Decorative Purposes and Every-day Uses. Furriers and Plumassiers, and Collectors in Natural History.

NOTICE.—ROWLAND WARD is the only member left in the profession of the Ward Family, long unrivalled for their accumulated experience and their skill in Practical Taxidermy, especially in its artistic department.

MEDALS AND DIPLOMAS OF HONOUR FOR ARTISTIC WORK

London International Exhibition, 1862. Paris International Exhibition, 1862. Vienna International Exhibition, 1873. London International Fisheries, 1883. Calcutta International Exhibition, 1883-84. London International Health Exhibition, 1884.

London Colonial and Indian Exhibition, 1886.

The Anglo-Danish Exhibition, South Kensington, 1888.

The Royal Military Exhibition (Army Medical Department), 1890.

THE COLONIAL AND INDIAN EXHIBITION, 1886

THE JUNGLE

AND INDIAN ANIMAL LIFE

WAS DESIGNED AND ARRANGED, AND THE ANIMALS MODELLED, BY

ROWLAND WARD, F.Z.S.

WHAT THE PRESS SAID:

"But everything else here is likely to be forgotten in presence of the wonderful jungle scene which Mr. Rowland Ward has constructed. . . . This will certainly be the first of the many attractions to which visitors will turn. . . . They will find themselves in presence of a scene which is likely to keep their gaze for some time. Mr. Ward has made the most of his limited space, into which he has collected the scenery and life which, in reality, is found scattered over an area of many thousand square miles. On the right we have a trophy from Kuch Behar, formed by His Highness the Maharajah, the most prominent feature of which is a tiger hunt. We see a great group in the deep grass jungle. . . . Adjoining this are trophies designed to represent generally the Fauna and Flora of India, by representative animals and birds, picturesquely grouped in illustration of their life-habits." — Times.

"The witters were lost in admiration of Mr. Rowland Wardt materly designs modellings and general

"The visitors . . . were lost in admiration of Mr. Rowland Ward's masterly designs, modellings, and general arrangement. The novelty is already known as 'the Jungle.' . . The deep grass jungle is occupied necessarily by many creatures which would not in their native wilds be found in such close companionship. . . . The scene is rendered with true tragic power."—Daily News.

"These numerous beasts . . . seem to illustrate the Fauna of India in a most vivid manner, and are very artistically prepared and arranged. . . . The entire trophy has been prepared by Mr. Rowland Ward. This group will unquestionably be one of the leading attractions of an exhibition which is already full of marvellous things."—Morning Post.

"Fitted up with the most perfect completeness—a jungle—the work of Mr. Rowland Ward. . . . The whole scene depicted is so life-like that one is startled by its vivid realism. . . . This jungle alone is almost enough to make an exhibition. . . Besides, Mr. Rowland Ward has designed and arranged such other scenes in connection with several Colonial Courts."—Daily Chronicle.

"Mr. Rowland Ward, of Piccadilly, provides what will probably prove the most attractive feature of the exhibition, in the form of a series of picturesque trophies representing India, Ceylon, South Africa, Canada, and Queensland."—Sportsman.

"Perhaps the first place must be accorded to the jungle scene of Mr. Rowland Ward, which stands at the head of the Indian Courts, and which will certainly prove one of the favourite sights of the vast show."—Daily Telegraph.

EMPIRE OF INDIA EXHIBITION, 1895-96

THE JUNGLE

- "A veritable triumph of the taxidermist's art—a tableau of jungle life which is entirely fresh and in every way remarkable."—Daily Telegraph.
- "A series of scenes illustrative of jungle life, admirable alike in its artistic effect and fidelity to nature."—
 Morning Advertiser.
 - ""The Jungle' will give the visitor vivid notions of Indian life."—Times.
- "Will draw all eyes—gentle and simple, town-bred and country-bred; is a wonderful exhibit. . . . such wealth of pelt and plumage, such glories of shikah . . . the very combined essence of all jungles."—Daily Chronicle.
- "A specially fine representation of an Indian jungle, with its characteristic vegetation and animals and wild scenery, to which Mr. Rowland Ward has contributed all his knowledge as a naturalist and his unrivalled skill as a taxidermist."—Standard.
- "Most attractive . . . a comprehensive representation of animal life in the jungle and on the mountains of India . . . surpasses all former efforts . . . most realistic."—Sporting Life.
 - "Entirely fresh, and in every way remarkable."-Graphic.
- "Rowland Ward's Jungle is the finest thing of the kind ever seen in this country, and should not be missed by any one."—Court Journal.
- "A realisation of nature in its wildest and most tragic moods . . . provides instruction and amusement for the thousands in whose breast the love of nature and animal life is implanted."—(Globe.
- "Grand grouping of tropical life. Scrupulous attention to detail. . . . The hoarse coughing roar of the tiger closely imitated."—South Africa.
- "Surpasses in interest any of the excellent exhibitions of the kind previously shown."—Manchester Guardian.
 "The number of persons who visited Mr. Rowland Ward's Jungle was 10,500 making over 200,000 since the opening."—Times (August 6, 1895).

LONDON: ROWLAND WARD, LIMITED, "THE JUNGLE," 166 PICCADILLY.

A SELECTED LIST OF BOOKS

OF

SPORT, TRAVEL

AND

NATURAL HISTORY



THE JUNGLE

AT THE INDIAN AND COLONIAL EXHIBITION, 1886,
DESIGNED AND THE ANIMALS MODELLED
BY ROWLAND WARD, F.Z.S.

PUBLISHED BY

ROWLAND WARD, LIMITED 166 PICCADILLY, LONDON

One Vol., Post 8vo, bound in leather. Price 3s. 6d. net

Great Game Shooting and Practical Taxidermy

THE

SPORTSMAN'S HANDBOOK

TO PRACTICAL COLLECTING, PRESERVING, AND
ARTISTIC SETTING-UP OF TROPHIES AND SPECIMENS'
TO WHICH IS ADDED A SYNOPTICAL GUIDE TO
THE HUNTING GROUNDS OF THE WORLD

WITH MANY ILLUSTRATIONS

By ROWLAND WARD, F.Z.S.

"With this in his portmanteau, no one fond of shooting and collecting need any longer lament his inability to preserve his trophies, since the directions given for skinning and preserving animals of all kinds are extremely clear and simple, and rendered all the more intelligible by the wood engravings by which they are accompanied. Quadrupeds, birds, fishes, reptiles, and insects are all dealt with in turns, and directions given not merely for skinning them, but also for mounting them, if desired, a year or two (it may be) after they have been procured."—Field.

LONDON: ROWLAND WARD, LIMITED "THE JUNGLE," 166 PICCADILLY

One Vol., Square 8vo, pp. viii. and 264. Illustrated. Price 21s. net

HORN MEASUREMENTS

WEIGHT, LENGTH, ETC.

OF THE GREAT GAME OF THE WORLD

BEING A RECORD FOR THE USE OF SPORTSMEN AND NATURALISTS

By ROWLAND WARD, F.Z.S.

"To sportsmen the utility of these voluminous records can hardly be over-estimated. In the majority of cases the accounts of the antiers and horns are illustrated by engravings; and photographs of many of the animals yielding them, with the sportsmen grouped around, are interspersed throughout the volume. We have thus the African elephant, the Java ox (Bos banting), the Cape buffalo, and the Thibet and Pallah antelopes."—Field.

LONDON: ROWLAND WARD, LIMITED "THE JUNGLE," 166 PICCADILLY

With about 150 Illustrations. One Vol., uniform with "Horn Measurements" 342 pp. Price 30s. net

RECORDS OF BIG GAME

CONTAINING

AN ACCOUNT OF THEIR DISTRIBUTION
DESCRIPTIONS OF SPECIES, LENGTHS, AND WEIGHTS

MEASUREMENTS OF HORNS

AND

FIELD NOTES

For the use of Sportsmen and Naturalists

By ROWLAND WARD, F.Z.S.

AUTHOR OF 'THE SPORTSMAN'S HANDBOOK,' ETC.

"In its present form, the work deals with nearly all the species of horned and antlered game, as well as with elephants, rhinoceroses, hippopotami, and wild boars, together with some of the larger carnivora, such as lions, tigers, leopards, and bears . . . the figures are admirable. With such an up-to-date and authentic record before them, sportsmen will in future have an easy task in deciding whether or no their trophies are record specimens. We may also call attention to the careful attention that has been paid to the proper scientific nomenclature."—The Asian.

LONDON: ROWLAND WARD, LIMITED "THE JUNGLE," 166 PICCADILLY

One Vol., Royal 8vo, 477 pp. Price 25s. net

TRAVEL AND ADVENTURE SOUTH-EAST AFRICA

Being the Narrative of the last eleven years spent by the Author on the Zambesi and its Tributaries; with an Account of the Colonisation of Mashonaland and the Progress of the Gold Industry in that Country

By F. C. SELOUS

GOLD MEDALLIST OF THE ROYAL GEOGRAPHICAL SOCIETY AUTHOR OF 'A HUNTER'S WANDERINGS IN AFRICA'

WITH NUMEROUS ILLUSTRATIONS AND MAP

"It is impossible even to indicate all the points of interest with which Mr. Selous deals. Illustrations are both numerous and excellent."—Times.

LONDON: ROWLAND WARD, LIMITED "THE JUNGLE," 166 PICCADILLY

SUNSHINE AND STORM IN RHODESIA

BEING A NARRATIVE OF EVENTS IN MATABELELAND BOTH BEFORE AND DURING THE RECENT NATIVE INSURRECTION

By F. C. SELOUS

GOLD MEDALLIST OF THE ROYAL GEOGRAPHICAL SOCIETY AUTHOR OF 'TRAVEL AND ADVENTURE,' ETC.

FULLY ILLUSTRATED

"It is a plain tale which Mr. Selous has to tell, and he tells it in a straightforward manner. This volume deserves to be widely read, not only because it is the earliest book on the recent rebellion, but also because it records facts without favour, and facts intensely interesting to Englishmen in all parts of the Empire."—Morning Post.

LONDON: ROWLAND WARD, LIMITED "THE JUNGLE," 166 PICCADILLY

Royal 8vo, about 400 pp. Price 18s. net

SEVENTEEN TRIPS THROUGH SOMALILAND

A RECORD OF EXPLORATION AND BIG GAME SHOOTING

1884 TO 1893

By CAPTAIN H. G. C. SWAYNE, R.E.

FELLOW OF THE ROYAL GEOGRAPHICAL SOCIETY CORRESPONDING MEMBER OF THE ZOOLOGICAL SOCIETY

WITH NUMEROUS ILLUSTRATIONS AND MAPS

"Will be carefully studied by those who have been there, and those—and there are many—who have an eager desire to go. It is full of thrilling episodes. . . . The last chapter . . . contains some highly interesting notes on the wild fauna of the country. The appendices, which deal with the fitting out of Somali expeditions and with the physical geography, have a distinct value."—Times.

LONDON: ROWLAND WARD, LIMITED "THE JUNGLE," 166 PICCADILLY

THE

ENGLISH ANGLER

IN

FLORIDA

WITH SOME DESCRIPTIVE NOTES OF THE GAME, ANIMALS, AND BIRDS

RV

ROWLAND WARD, F.Z.S.

AUTHOR OF 'RECORDS OF BIG GAME,' 'SPORTSMAN'S HANDBOOK,' BTC.

WITH NUMEROUS ILLUSTRATIONS

"Mr. Rowland Ward, so widely known as a sporting publisher, has published a capital little book of his own writing. The English Angler in Florida gives an entertaining and eminently practical account of an angling trip he made to Florida last spring. Punta Gorda, on the western shore of the Peninsula, may be reached in nine days from Piccadilly, and, now that salmon fishings command fancy prices at home, English fishermen have been turning their attention to the tarpon. The tarpon is coming into notoriety, and promises to become the fashion, but it is only within the last ten or fifteen years that he has been persecuted. His flesh is worthless; even the negro will not eat it, and nothing was suspected of the grand sport he gives with rod and line. As he runs ordinarily from 100 lbs. to 200 lbs., and is at least as vigorous in proportion as the salmon, the first consideration is a first-class outfit. Mr. Ward tells where it is best procured, and describes the short, stiff rod, the reel, the line, etc., and notably 'the friendly socket,' to be strapped round the waist, in which the butt of the rod is inserted; for the strain upon back and arms is tremendous when fighting a 150 lb. fish for an hour or more. The tarpon is taken by gorge bait, and, unfortunately, the hook has yet to be invented which might be available for successful fly-fishing. For only here and there is the interior of the mouth vulnerable; elsewhere, in Mr. Ward's words, it is impenetrable as a stone wall. He went in for his expedition almost regardless of expense, but he points out how it can be done as well and more economically; and in particular he recommends chartering a sailing schooner instead of a steam launch. Sharks infest the fishing-grounds, snapping off the baits; 'the slimy and filthy catfish 'swarm, and are even more troublesome; and there is an interesting chapter on other eccentric marine monsters. He heard doubtful talk of a devil's ray, said to be 20 feet in diameter; there is the sawfish, which with its formidable serrated pr

LONDON: ROWLAND WARD, LIMITED

"THE JUNGLE," 166 PICCADILLY

ELEPHANT-HUNTING

IN

EAST EQUATORIAL AFRICA

BEING

AN ACCOUNT OF THREE YEARS' IVORY-HUNTING UNDER MOUNT KENIA AND AMONG THE NDOROBO SAVAGES

OF THE LOROGI MOUNTAINS, INCLUDING A TRIP TO THE NORTH END OF LAKE RUDOLPH

BY

ARTHUR H. NEUMANN

WITH NUMEROUS ILLUSTRATIONS BY J. G. MILLAIS, E. CALDWELL AND G. E. LODGE

COLOURED PLATE AND MAP

LONDON: ROWLAND WARD, LIMITED

"THE JUNGLE," 166 PICCADILLY

[&]quot;It may be said that Mr. Neumann's volume is one of the most interesting books of African sport, adventure, and natural history that has appeared since that of Mr. Selous, and the illustrations are worthy of the text."—Scotsman.

[&]quot;There is no lack of adventures throughout the book, and accounts of many narrow escapes from elephants. In addition to this, the author gives many interesting notes on the country traversed, and the manners and customs of the people with whom he came in contact. . . . Mr. Neumann may be congratulated on having produced a book which will be read with interest by sportsmen and geographers.

[&]quot;The illustrations which the book contains are worthy of special commendation, and the map with which it is furnished will enable the reader to follow Mr. Neumann in his wanderings through a country very little of which, indeed, is known."—Field.

[&]quot;This handsome volume, well written, well printed, well illustrated, and filled with such experiences of wild-forest life as few other men could boast, will delight the sportsman, geographer, the ethnologist, and the general reader. . . . He took his full share of fevers, thirst, lost roads, dangers by field and flood, and even as regards his principal prey, the elephant, was once as near to a dreadful death as any man who ever lived to tell the tale of having been charged and caught by a wounded elephant."—Daily Telegraph.

SPORT IN THE HIGHLANDS

OF

KASHMIR

BEING

A NARRATIVE OF AN EIGHT MONTHS' TRIP IN BALTISTAN AND LADAK, AND A LADY'S EXPERIENCES IN THE LATTER COUNTRY; TOGETHER WITH HINTS FOR THE GUID-ANCE OF SPORTSMEN

RV

HENRY ZOUCH DARRAH

INDIAN CIVIL SERVICE

WITH FIFTY-TWO ILLUSTRATIONS (FROM PHOTOGRAPHS BY THE WRITER) AND TWO MAPS

"Mr. Darrah's narrative of his visit to Kashmir and its outlying provinces Baltistan and Ladak has many merits. It has the excellent foundation of notes 'generally recorded on the evenings of the days on which the events described occurred,' and of a diary kept whilst memory was fresh, the result being a book faithful and accurate in description of country and people . . . eminently rustworthy of the sport which may be enjoyed."—Athenaum.

"Mr. Henry Zouch Darrah's sumptuous volume Sport in the Highlands of Kashmir is the best book we have seen on sport beyond the Himalayas since General Macintyre's, nearly twenty years ago."—Times.

LONDON: ROWLAND WARD, LIMITED, "THE JUNGLE," 166 PICCADILLY.

HUNTING TRIPS THE CAUCASUS

BY

E. DEMIDOFF

PRINCE SAN DONATO

"Hunting Trips in the Caucasus, by Prince Demidoff, will be of interest not only to sportsmen, but to the general reader. In his preface the author, as a foreigner, appeals to English sportsmen for indulgence. We do not, however, think that there is any necessity for him to do so, as, in our opinion, the most interesting portion of the book is contained in his chapter on the game of the Caucasus, and the description of the first hunting trip, both of which have been written by himself. The book is well got up, contains numerous excellent illustrations, and is furnished with a good map."—Field.

WITH 96 ILLUSTRATIONS AND MAP

ROWLAND WARD, LIMITED
166 PICCADILLY

THE DEER OF ALL LANDS

A HISTORY OF THE FAMILY CERVIDÆ LIVING AND EXTINCT

By R. LYDEKKER

Illustrated by Twenty-four Hand-Coloured Plates, drawn by J. Smit; also a number of Process-Drawings of Horns, as well as
Photographic Reproductions of living Deer in the
Woburn and other Collections

"Mr. Lydekker's beautiful volume will, of course, find its way into the hands of every sportsman who can afford to indulge in the fascinating pursuit of 'big game' shooting, but it will also form an authoritative work of reference for naturalists of all countries, who will find in its pages a fund of useful information."—

LONDON: ROWLAND WARD, LIMITED, "THE JUNGLE," 166 PICCADILLY.

Uniform with the above. Price £5:5:0 net.

WILD

OXEN, SHEEP, AND GOATS OF ALL LANDS

By R. LYDEKKER

Illustrated by Twenty-eight Coloured Plates, drawn by J. SMIT and JOSEPH WOLF; and a number of Drawings of Horns, as well as Photographic Reproductions of living Specimens

LONDON: ROWLAND WARD, LIMITED, "THE JUNGLE," 166 PICCADILLY.

THE

GREAT & SMALL GAME OF AFRICA

Being a full Account of their Distribution, Habits, and Natural History, with Numerous Experiences of their Pursuit in the Field

WITH NUMEROUS HAND-COLOURED AND OTHER ILLUSTRATIONS

CONTRIBUTORS

MAJOR A. J. ARNOLD
H. A. BRYDEN
T. E. BUCKLEY
T. W. H. CLARKE
LORD DELAMERE
DR. D. G. ELLIOTT
CAPT. B. T. FFINCH
H. C. V. HUNTER
J. D. INVERARITY
F. J. JACKSON
SIR HARRY JOHNSTON
F. VAUGHAN KIRBY
R. LYDEKKER

CAPT. JOHN MARRIOTT
A. H. NEUMANN
A. E. PEASE
G. W. PENRICE
VISCOUNT EDMOND DE PONCINS
DR. PERCY RENDALL
F. C. SELOUS
ALFRED SHARPE
LT.-COL. WILLIAM SITWELL
A. H. STRAKER
MAJOR H. G. C. SWAYNE
POULETT-WEATHERLEY

General Editor .- H. A. BRYDEN

LONDON

ROWLAND WARD, LIMITED

"THE JUNGLE," 166 PICCADILLY

